# Southeast Alaska chum salmon (Oncorhynchus keta) thermal mark identification and agreement

2012 Pink and Chum salmon workshop, Juneau AK

Lorna Wilson

Bev Agler

ADF&G Thermal Mark Lab

Juneau, AK

## **Introduction: Thermal marks**

Salmonid otoliths are mass thermally marked

Temperature cycles → banding pattern Applications:

- In-season sockeye management
- Hatchery return rates
- Hatchery stray rates
- Evaluate rearing strategies
- Validate model parameter estimates

# Introduction: Detection and Identification

Mark detection

Presence of thermal mark

Hatchery vs. Wild

Mark identification

Hatch code → unique group

E.g. 1,4,3H chum means:


Thermal mark ID: NEETSBAY05SUM

Brood Year: 2005


Agency: SSRAA

Number released: 8,409,868

Stock: Neets Bay



## Introduction: Variants



How accurate are chum salmon thermal mark detections and identifications?

# Introduction: Mark accuracy

ADFG laboratory data quality assurance methods: Study and use known reference collection:

- 1) Mark variations
- 2) Mark measurements: to mark, between bands

Multiple independent reads to identify issues


But... true reader error rate is unknown!

# **Objective**

Assess accuracy of thermal mark detection and identification among areas and labs

## Study area

- Southern Southeast
   Northern Inside
- 2) Lynn Canal and Stephens Passage
- 3) Chatham and Icy straits
- 4) Northern Outside



#### **Methods: Reads**

2009 and 2010 recoveries

Read by:

AK Department of Fish and Game Thermal Mark Lab

ADF&G

Southern SE Regional Aquaculture Association SSRAA

Douglas Island Pink and Chum Inc.

DIPAC

For:

Readability

Presence

Identification

#### **Methods: LCM**

Latent class models (LCM) estimate reader ability:

to detect marked fish when it is marked (H | H)

to detect wild fish when it is wild (W | W)

Using:

- Strata: area (e.g. Northern Inside waters)
- Reader pairs (e.g. ADF&G and DIPAC) to estimate the true reader error rate.

Maximize likelihood function, SE is estimated using jackknife method.

# Methods: Kappa

Agreement between readers on two or more identifications

Accounts for agreement that occurs by chance:

$$\kappa = \frac{p_o - p_e}{1 - p_e}$$

$$p_o = observed$$
,

$$p_e = expected$$



$$SE(\kappa)$$

Overall  $\kappa$  = weighted average ( $\kappa$ )


# Methods: Assumptions

- 1. Independent readings
  - Dependence is caused by:
  - a) Otolith preparation not independent.
  - b) Marking process affecting readability of a mark grouping.


Somewhat resolved with more strata

2. Accuracy rate is greater than error rate.


#### **Results: Mark detection**




#### **Results: Mark detection**




#### **Results: Mark Identification**



#### **Results: Mark Identification**





#### **Discussion**

Accuracy > error rate assumption met.

LCM and Kappa values are appropriate for assessing reader accuracy.

Reader ability to detect a mark higher in 2009 than 2010, but individual mark identification was lower. This possibly due to:

Fewer marked fish in 2009
Better mark quality in 2010
Increased reader ability
Sample loss due to over-grinding



## **Discussion**

Overall, reader ability to detect and identify chum salmon thermal marks is high.

Second reads are essential for assessing reader accuracy.

Reader ability to detect and identify marks increases with:

Mark assignment
Mark quality
Training



# Acknowledgements

DIPAC otolith lab
SSRAA otolith lab
ADF&G Thermal Mark Lab
Megan Lovejoy
Joe Cashen
John Baker



Dion Oxman and Ron Josephson for guidance