

**PACIFIC SALMON COMMISSION JOINT CHINOOK
TECHNICAL COMMITTEE REPORT**

**ANNUAL REPORT ON CATCH, ESCAPEMENT,
EXPLOTATION RATE ANALYSIS AND MODEL CALIBRATION
OF CHINOOK SALMON UNDER
PACIFIC SALMON COMMISSION JURISDICTION, 2006**

REPORT TCCHINOOK (07)-1

January 30, 2007

TABLE OF CONTENTS

| | <u>Page</u> |
|--|-------------|
| Table of Contents..... | ii |
| Membership of the Chinook Technical Committee..... | iv |
| List of Acronyms with Definitions | v |
| List of Tables | vi |
| List of Figures..... | vii |
| List of Figures..... | vii |
| List of Appendices | viii |
| List of Appendices | viii |
| Executive Summary..... | 1 |
| 1 Chinook Catch..... | 7 |
| 1.1 Review of AABM Fisheries..... | 7 |
| 1.1.1 Southeast Alaska Fisheries | 8 |
| 1.1.2 British Columbia Fisheries | 9 |
| 1.1.2.1 NBC Troll Fishery Harvest..... | 10 |
| 1.1.2.2 NBC and CBC Recreational Fishery Harvest..... | 11 |
| 1.1.2.3 WCVI AABM..... | 11 |
| 1.2 Estimates of Incidental Mortalities in AABM Fisheries..... | 13 |
| 1.2.1 SEAK Fisheries..... | 13 |
| 1.2.2 British Columbia Fisheries | 14 |
| 1.2.2.1 NBC Fisheries..... | 14 |
| 1.2.2.2 WCVI Fishery | 15 |
| 1.3 Review of ISBM Fisheries..... | 16 |
| 1.3.1 Canadian ISBM Fisheries | 16 |
| 1.3.2 Southern U.S. Fisheries Harvest..... | 18 |
| 1.3.2.1 Strait of Juan de Fuca and the San Juan Islands | 18 |
| 1.3.2.2 Puget Sound | 18 |
| 1.3.2.3 Washington Coast | 18 |
| 1.3.2.4 Columbia River..... | 19 |
| 1.3.2.5 Ocean Fisheries, Cape Falcon to Humbug Mountain..... | 19 |
| 1.4 Estimates of incidental mortality for Southern U.S. Fisheries..... | 19 |
| 2 Escapements through 2005..... | 21 |
| 2.1 Introduction..... | 21 |
| 2.1.1 MSY or Biologically-Based Escapement Goals..... | 21 |
| 2.1.1.1 Origin of Goals | 21 |
| 2.2 Escapement Assessment | 23 |
| 2.3 Stock Specific Graphs and Commentaries..... | 25 |
| 2.3.1 SEAK/TBR Stocks..... | 25 |
| 2.3.2 Canadian Stocks..... | 31 |
| 2.3.3 Fraser River Stocks..... | 38 |
| 2.3.4 Washington, Oregon and Columbia River Stocks | 42 |
| 3 Exploitation Rate Analysis and model calibration..... | 59 |
| 3.1 Introduction..... | 59 |
| 3.2 Methods..... | 59 |

| | | |
|---------|--|----|
| 3.3 | Exploitation Rate Assessment (Through Calendar Year 2004)..... | 60 |
| 3.4 | Model Output..... | 63 |
| 3.4.1 | AABM Abundance Indices and Associated Catches..... | 63 |
| 3.4.1.1 | Model estimate of stock composition of AABM fisheries, 1979-2006 | 65 |
| 3.4.2 | Overages and Underages..... | 68 |
| 3.4.2.1 | AABM Fisheries | 68 |
| 3.4.2.2 | ISBM Indices by Stock | 69 |
| 3.5 | Model Calibration Evaluation..... | 74 |
| 3.6 | Evaluation of mark-selective fisheries..... | 78 |
| | References Cited | 80 |
| | Appendices..... | 83 |

MEMBERSHIP OF THE CHINOOK TECHNICAL COMMITTEE

Canadian Members

Dr. Rick McNicol, Co-Chair, CDFO
Mr. Richard Bailey, CDFO
Dr. Gayle Brown, CDFO
Mr. Roger Dunlop, FNC
Mr. Ryan Galbraith, CDFO
Mr. Wilf Luedke, CDFO
Mr. Chuck Parken, CDFO
Ms. Teresa Ryan, FNC
Mr. Julian Sturhahn, CDFO
Dr. Arlene Tompkins, CDFO
Mr. Ivan Winther, CDFO
Mr. Howie Wright, FNC

United States Members

Mr. C. Dell Simmons, Co-Chair, NMFS
Mr. John Carlile, Co-Chair, ADF&G
Dr. Dave Bernard, ADF&G
Mr. Ethan Clemons, ODFW
Dr. John H. Clark, ADF&G
Mr. Gary Freitag, SSRAA
Mr. Ed Jones, ADF&G
Dr. Robert Kope, NMFS
Mr. Brian Lynch, ADF&G
Ms. Marianne McClure, CRITFC
Mr. Scott McPherson, ADF&G
Dr. Gary Morishima, QIN
Mr. James Packer, WDFW
Mr. Rishi Sharma, CRITFC
Dr. Brad Thompson, WDFW
Mr. Alex Wertheimer, NMFS
Mr. Henry Yuen, USFWS

LIST OF ACRONYMS WITH DEFINITIONS

| | | | |
|------------------|---|------------------------|---|
| AABM | Aggregate Abundance Based Management | MSH | Maximum sustainable harvest |
| AC | Allowable Catch | MSY | Maximum Sustainable Yield for a stock, in adult equivalents |
| AI | Abundance Index | MSY ER | Exploitation Rate sustainable at the escapement goal for a stock, in AEQs |
| ADF&G | Alaska Department of Fish & Game | NBC | Northern British Columbia Dixon Entrance to Kitimat including Queen Charlotte Islands |
| AEQ | Adult Equivalent | NA | Not Available |
| Agreement | June 30, 1999 PST Annex and the related Agreement | NBC | Northern British Columbia Dixon Entrance to Kitimat including Queen Charlotte Islands |
| AUC | Area Under the Curve | NM | Nautical Mile |
| AWG | Analytical Working Group of the CTC | NMFS | National Marine Fisheries Service |
| BCAFC | British Columbia Aboriginal Fisheries Commission | NOC | Oregon Coastal North Migrating Stocks |
| BTR | Base Terminal Run | NPS | North Puget Sound |
| C&S | Ceremonial & Subsistence | NPS-S/F | North Puget Sound Summer/Fall Chinook stock |
| CBC | Central British Columbia Fishing area – Kitimat to Cape Caution | NR | Not Representative |
| CCMP | Comprehensive Chinook Management Plan | NWIFC | Northwest Indian Fisheries Commission |
| CDFO | Canadian Department of Fisheries & Oceans | ODFW | Oregon Department of Fish & Wildlife |
| CI | Confidence Interval | OTAC | Outside Troll Advisory Committee |
| CNR | Chinook Non-retention | PFMC | Pacific Fisheries Management Council |
| CR | Columbia River | PS | Puget Sound |
| CRITFC | Columbia River Intertribal Fish Commission | PSC | Pacific Salmon Commission |
| CRFMP | Columbia River Fishery Management Plan | PSARC | Pacific Scientific Advice Review Committee |
| CTC | Chinook Technical Committee | PSMFC | Pacific States Marine Fisheries Commission |
| CUS | Columbia Upriver Spring Chinook stock | PST | Pacific Salmon Treaty |
| CWT | Coded Wire Tag | QDNR | Quinault Department of Natural Resources, Division of fisheries |
| DIT | Double Index Tag | QIN | Quinault Nation |
| ESA | U.S. Endangered Species Act | QCI | Queen Charlotte Islands |
| Est+fw | Estuary Plus Fresh Water Area | RER | Recovery Exploitation Rate |
| FL | Fork Length | S_{MSY} | Escapement producing MSY |
| FMP | PFMC Framework Management Plan | SEAK | Southeast Alaska Cape Suckling to Dixon Entrance |
| FNC | First Nations Council | SG | Strait of Georgia |
| FOG | Fisheries Operational Guidelines | SPS | South Puget Sound |
| FR | Fraser River | SSRAA | Southern Southeast Regional Aquaculture Association |
| GCG | Gene Conservation Group | SWVI | Southwest Vancouver Island |
| GW | Gitwinksihlkw | TAC | Technical Advisory Committee |
| GS | Strait of Georgia | TBR | Transboundary Rivers |
| HOR | Hatchery Origin Returns | TTC | Transboundary Technical Committee |
| IDFG | Idaho Department of Fish & Game | UFR | Upper Fraser River |
| IDL | InterDam Loss | UGS | Upper Strait of Georgia |
| IM | Incidental Mortality | USCTC | U.S. members of the CTC |
| ISBM | Individual stock based management | USFWS | U.S. Fish & Wildlife Service |
| LFR | Lower Fraser River | UW | University of Washington |
| LGS | Lower Strait of Georgia | WA/OR | Ocean areas off Washington and Oregon North of Cape Falcon |
| mar | Marine Area | WAC | Washington Coast (Grays Harbor northward) |
| mar+fw | Marine Plus Fresh Water Area | WACO | Washington, Oregon, Columbia River Chinook stock group |
| MOC | Mid Oregon Coast | WCVI | West Coast Vancouver Island excluding Area 20 |
| MRP | Mark-Recovery Program | WDFW | Washington Department of Fisheries and Wildlife |
| MSF | Mark-Selective Fishery | | |

LIST OF TABLES

| | | |
|-------------|--|----|
| Table 1. | Abundance Indices for 1999 to 2006 for the SEAK, NBC, and WCVI AABM fisheries. | 1 |
| Table 2. | Observed catches and postseason allowable catches for 1999 to 2005, and preseason allowable catches for 1999 to 2006, for AABM fisheries. | 2 |
| Table 3. | Deviations in numbers of Chinook salmon and percentages from catch targets derived from the first postseason AI (Table 3.2) for Pacific Salmon Treaty AABM fisheries in 1999 to 2005. | 3 |
| Table 4. | Canadian 2004 ISBM indices based on CWT and the 2006 indices predicted from the PSC Chinook Model. | 4 |
| Table 5. | U.S. 2004 ISBM indices based on CWT and the 2006 indices predicted from the PSC Chinook Model. | 5 |
| Table 1.1. | Annual catches and hatchery add-ons for the AABM fisheries, in thousands of Chinook salmon. The Treaty catches do not include the add-on or exclusions (see Section 1.1.1 and Appendix A.1). Notation is T for Troll, N for Net and S for sport. | 7 |
| Table 1.2. | Harvest of Chinook salmon in SEAK by gear type in 2005. | 9 |
| Table 1.3. | Summary of landed catch by gear for Canadian AABM fisheries in 2005. | 11 |
| Table 1.4. | Fishing periods and Chinook harvested and released during the 2005 accounting year in the WCVI troll fishery. | 12 |
| Table 1.5. | Outer WCVI AABM recreational fishery catches of Chinook by statistical area in 2005 representing catch during the creel survey periods only. | 13 |
| Table 1.6. | Estimated encounters and incidental mortality in SEAK troll, net and sport fisheries for 2003-2005. Mortality estimates of fish released in troll and sport fisheries include drop-off mortality. In the net fishery, 21"-28" fish from both retention and non-retention periods are included in the CNR numbers. | 14 |
| Table 1.7. | Estimated encounters and incidental mortalities (nominal fish) in NBC AABM troll and sport fisheries for 2002-2005. Mortality estimates of fish released in troll and sport fisheries include drop-off mortality. | 15 |
| Table 1.9. | Landed catch and incidental mortalities in Canadian ISBM fisheries for 2005. | 17 |
| Table 1.10. | Estimated incidental mortality in Southern US troll, net, and sport fisheries for 2005. | 20 |
| Table 2.1. | PSC Chinook escapement indicator stocks, where shading indicates that there is not a | 22 |

LIST OF FIGURES

| | | |
|--------------|--|----|
| Figure 1.1. | British Columbia fishery management areas..... | 10 |
| Figure 2.1. | Number and status of stocks with CTC-accepted escapement goals for years 1999-2005..... | 25 |
| Figure 3.1. | Preseason allowable catches (open circles) and postseason catches (diamonds) in Southeast Alaska AABM fisheries, 1999-2005. | 64 |
| Figure 3.2. | Preseason allowable catches (open circles) and postseason catches (diamonds) in Northern British Columbia troll and Queen Charlotte Islands recreational AABM fisheries, 1999-2005. | 65 |
| Figure 3.3. | Preseason allowable catches (open circles) and postseason catches (diamonds) in West Coast Vancouver Island AABM fisheries, 1999-2005. | 65 |
| Figure 3.4. | Total abundance indices for the SEAK troll fishery with annual stock composition indicated by abundance indices for major model stocks from CLB 0604. | 66 |
| Figure 3.5. | Total abundance indices for the NBC troll fishery with annual stock composition indicated by abundance indices for major model stocks from CLB 0604. | 67 |
| Figure 3.6. | Total abundance indices for the WCVI troll fishery with annual stock composition indicated by abundance indices for major model stocks from CLB 0604. | 68 |
| Figure 3.7. | ISBM indices for Canadian fisheries for 1999-2004. The solid horizontal line is an index value of 0.635..... | 70 |
| Figure 3.8. | ISBM indices for U.S. fisheries for 1999-2004. | 71 |
| Figure 3.9. | Estimated CWT (through 2004) and model landed catch fishery indices (through 2005) for the SEAK troll fishery. | 74 |
| Figure 3.10. | Estimated CWT (through 2004) and model total mortality fishery indices (through 2005) for the SEAK troll fishery. | 75 |
| Figure 3.11. | Estimated CWT (through 2004) and model landed catch fishery indices (through 2005) for the NBC troll fishery. | 76 |
| Figure 3.12. | Estimated CWT (through 2004) and model total mortality fishery indices (through 2005) for the NBC troll fishery. | 76 |
| Figure 3.13. | Estimated CWT (through 2004) and model landed catch fishery indices (through 2005) for the WCVI troll fishery. | 77 |
| Figure 3.14. | Estimated CWT (through 2004) and model total mortality fishery indices (through 2005) for the WCVI troll fishery. | 77 |

LIST OF APPENDICES

| | |
|---|-----|
| Appendix A. Landed Chinook catches by region and gear from 1975-2005. | 85 |
| Appendix B. Escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005. | 101 |
| Appendix C. Relationship between exploitation rate indicator stocks, escapement indicator stocks, model stocks, and additional management action stocks identified in the PST annex. | 111 |
| Appendix D. ISBM indices. | 117 |
| Appendix E. Percent distribution of landed catch and total mortality among fisheries and escapement for exploitation rate indicator stocks by calendar year. | 121 |
| Appendix F. Time series of abundance indices from 1979 to 2005 for SEAK, NBC, and WCVI AABM fisheries as estimated by CTC Chinook Model calibration CLB0604. | 181 |
| Appendix G. Model estimates of the stock composition of the AABM, and other troll and sport fisheries for 2005 and the average from 1985 to 2004. | 183 |
| Appendix H. Incidental mortality rates applied in the CTC model. Rates in original model were applied to all years. In the current model, rates in some fisheries vary in accordance to changes in management regulations. | 191 |
| Appendix I. Components of the February 2005 agreement on harvest sharing for Stikine and Taku Rivers Chinook. | 193 |
| Appendix J. Evaluation of mark-selective fisheries. | 195 |

EXECUTIVE SUMMARY

The June 30, 1999, Pacific Salmon Treaty (PST) Annexes and Related Agreements (Agreement) substantially changed the objectives and structure of the Pacific Salmon Commission's (PSC) Chinook salmon fisheries and assessment of Chinook salmon stocks. The Agreement replaced the previous ceiling and pass-through fisheries with Aggregate Abundance Based Management (AABM) and Individual Stock Based Management (ISBM) fisheries. It also assigned the Chinook Technical Committee (CTC) with a number of tasks related to implementation of the Agreement (Appendix to Annex IV, Chapter 3).

In October, 2005, the CTC decided to combine the annual Catch and Escapement and the Calibration and Exploitation Rate Analysis reports into one document due to time constraints resulting from other assignments. In this report, we provide a summary of 2005 fishery catches by region, available estimates of incidental mortality by fishery and limited commentary on fishery catches where needed. Landed catch is reported in the appendices for each geographic area covered under the PST. An assessment of escapement for stocks with CTC accepted goals is included, and escapement data are provided for all escapement indicator stocks. This report also contains the principal results of the annual exploitation rate assessment and the final preseason Chinook model calibration for 2006 (CLB 0604). Results include the Abundance Indices (AIs) for the AABM fisheries and ISBM indices for each party (country).

AABM ABUNDANCE INDICES AND ASSOCIATED CATCHES

The pre- and postseason AIs for the three AABM fisheries, Southeast Alaska All Gear (SEAK), Northern British Columbia Troll and Queen Charlotte Islands Sport (NBC), and West Coast Vancouver Island Troll and Outside Sport (WCVI) are presented in Table 1. The Agreement specifies that the AABM fisheries are to be managed through the use of the AIs. Each calibration provides the first postseason AIs for the previous year and the preseason AIs for the current year. Preseason AIs are used to set total allowable catch limits in the upcoming fishing season. Subsequently, postseason AIs (from the following year's calibration) are used to track catch overage and underage provisions. The first 2005 postseason AIs and the 2006 preseason AIs have now been finalized.

Table 1. Abundance Indices for 1999 to 2006 for the SEAK, NBC, and WCVI AABM fisheries.

| Year | SEAK | | NBC | | WCVI | |
|------|-----------|------------|-----------|------------|-----------|------------|
| | Preseason | Postseason | Preseason | Postseason | Preseason | Postseason |
| 1999 | 1.15 | 1.12 | 1.12 | 0.97 | 0.60 | 0.50 |
| 2000 | 1.14 | 1.10 | 1.00 | 0.95 | 0.54 | 0.47 |
| 2001 | 1.14 | 1.29 | 1.02 | 1.22 | 0.66 | 0.68 |
| 2002 | 1.74 | 1.82 | 1.45 | 1.63 | 0.95 | 0.92 |
| 2003 | 1.79 | 2.17 | 1.48 | 1.90 | 0.85 | 1.10 |
| 2004 | 1.88 | 2.06 | 1.67 | 1.83 | 0.90 | 0.98 |
| 2005 | 2.05 | 1.90 | 1.69 | 1.65 | 0.88 | 0.84 |
| 2006 | 1.69 | | 1.53 | | 0.75 | |

In general, the AIs for 1999 through 2001 are low compared to AIs in the late 1980s and early 1990s but values have increased significantly starting in 2002. The 2006 projected AI values have declined compared to the high values in 2004 and 2005. The Agreement specifies an allowable catch for each AI for each fishery. The maximum allowable Treaty catch (total catch minus any hatchery add-on and exclusion catch) by fishery and year and the actual (observed) catches are shown in Table 2.

Table 2. Observed catches and postseason allowable catches for 1999 to 2005, and preseason allowable catches for 1999 to 2006, for AABM fisheries.

| PST Treaty Allowable and Observed Catches | | | | | | | | | |
|---|-----------------------------|-----------------------------|---------------------------------|----------------------------|-----------------------------|----------------|----------------------------|-----------------------------|----------------|
| Year | SEAK (T, N, S) ¹ | | | NBC (T, S) | | | WCVI (T, S) | | |
| | Pre-season Allowable Catch | Post-season Allowable Catch | Observed Catch | Pre-season Allowable Catch | Post-season Allowable Catch | Observed Catch | Pre-season Allowable Catch | Post-season Allowable Catch | Observed Catch |
| 1999 | 192,800 | 184,200 | 198,842 | 145,600 | 126,100 | 86,726 | 128,300 | 107,000 | 36,413 |
| 2000 | 189,900 | 178,500 | 186,493 | 130,000 | 123,500 | 31,900 | 115,500 | 86,200 | 101,438 |
| 2001 | 189,900 | 250,300 | 186,919 | 132,600 | 158,900 | 43,500 | 141,200 | 145,500 | 117,670 |
| 2002 | 356,500 | 371,900 | 357,133 | 192,700 | 237,800 | 150,137 | 203,200 | 196,800 | 165,036 |
| 2003 | 366,100 | 439,600 | 380,152 | 197,100 | 277,200 | 191,657 | 181,800 | 268,900 | 175,821 |
| 2004 | 383,500 | 418,300 | 428,773 433,446 ² | 243,600 | 267,000 | 241,508 | 192,500 | 209,600 | 216,624 |
| 2005 | 416,400 | 387,400 | 386,707 | 246,600 | 240,700 | 243,606 | 188,200 | 179,700 | 202,662 |
| 2006 | 346,800 | | | 223,200 | | | 160,400 | | |

¹ Nomenclature is T for troll, N for net, and S for sport.

² The lower value resulted from subtracting a disputed terminal exclusion catch for the Stikine River in 2004. Catch accounting has since been defined in the Transboundary Agreement.

Table 3. Deviations in numbers of Chinook salmon and percentages from catch targets derived from the first postseason AI (Table 3.2) for Pacific Salmon Treaty AABM fisheries in 1999 to 2005.

| Year | SEAK | | NBC | | WCVI | |
|------|----------------|--------------------|----------------|--------------------|----------------|--------------------|
| | Number of Fish | Percent Difference | Number of Fish | Percent Difference | Number of Fish | Percent Difference |
| 1999 | +14,642 | +7.9% | -39,374 | -31.2% | -70,587 | -66.0% |
| 2000 | +7,993 | +4.5% | -91,600 | -74.2% | +15,238 | +17.7% |
| 2001 | -63,381 | -25.3% | -115,400 | -72.6% | -27,830 | -19.1% |
| 2002 | -14,767 | -4.0% | -87,663 | -36.9% | -31,764 | -16.1% |
| 2003 | -59,448 | -13.5% | -85,543 | -30.9% | -93,079 | -34.6% |
| 2004 | +10,473 | +2.5% | -25,492 | -9.5% | +7,024 | +3.4% |
| | +15,146 | +3.6% | | | | |
| 2005 | -693 | -0.2% | +2,906 | +1.2% | +22,962 | +12.8% |

¹ The lower value resulted from subtracting a disputed terminal exclusion catch for the Stikine River in 2004. Catch accounting has since been defined in the Transboundary Agreement.

ISBM INDICES

For ISBM fisheries, the Agreement specified that Canada and the United States would reduce base period exploitation rates on specified stocks by 36.5% and 40%, equivalent to ISBM indices of 63.5% and 60% percent, respectively. This requirement is contained in Chapter 3 section 4(d) of the treaty and is referred to as the ‘general obligation’ and does not apply to stock groups that achieve their CTC agreed escapement goals. Estimated ISBM fishery indices are shown in Table 4 for Canadian fisheries and Table 5 for United States (U.S.) fisheries. Both tables present CWT-based indices for 2004, and Chinook model-based indices for 2006. The agreement specifies that the ISBM indices be forecasted preseason and evaluated postseason for each escapement indicator stock listed in Attachments I to V of the Chinook Chapter.

CWT-based Indices in 2004

All Canadian ISBM indices from the CWT-based estimates for 2004 show that exploitation rates were reduced more than required for all stocks or stock groups for which the indices could be calculated. Five of the 16 U.S. ISBM indices for the Coded Wire Tag (CWT) based estimates for 2004 were reduced more than required. Of the 11 U.S. CWT-based ISBM indices that exceeded 0.60, eight (Upriver Brights, Quillayute, Queets, Hoh, Mid-Columbia Summers, Nehalem, Siletz, and Siuslaw) have agreed escapement goals and all eight exceeded their goals in 2004.

Predicted ISBM Indices for 2006

Eight of the 20 ISBM indices for Canada in 2006 based on outputs from calibration 0604 are above the allowable value of 0.635 for Canadian ISBM fisheries. None of these stocks have CTC agreed escapement goals. Eleven of the 24 U.S. ISBM indices for 2006 based on calibration 0604 are above the allowable limit of 0.60 for U.S. ISBM fisheries. Ten of the 11 have CTC agreed escapement goals: Queets, Hoh, Quillayute, Upriver Brights, Lewis, Harrison, Mid-Columbia Summers, Nehalem, Siletz, and Siuslaw.

Table 4. Canadian 2004 ISBM indices based on CWT and the 2006 indices predicted from the PSC Chinook Model.

| Stock Group | Escapement Indicator Stock | Canadian ISBM Indices | |
|---|--|-----------------------|------------------------|
| | | CWT Indices for 2004 | Model Indices for 2006 |
| Lower Strait of Georgia | Cowichan ² | 0.284 ^{1,4} | 0.590 |
| | Nanaimo | NA ⁵ | 0.590 ⁶ |
| Fraser Late | Harrison River ² | 0.032 ⁷ | 0.294 |
| North Puget Sound Natural Springs | Nooksack | NA | 0.993 |
| | Skagit | NA | 0.993 |
| Upper Strait of Georgia | Klinaklini, Kakweikan, Wakeman, Kingcome, Nimpkish | 0.018 | 0.584 |
| Fraser Early (spring and summers) | Upper Fraser, Mid Fraser, Thompson | NA | 0.610 |
| West Coast Vancouver Island Falls | WCVI (Artlish, Burman, Kauok, Tahsis, Tashish, Marble) | 0.488 ⁸ | 1.082 |
| Puget Sound Natural Summer / Falls | Skagit | NA | 1.092 |
| | Stillaguamish | 0.027 | 1.116 |
| | Snohomish | NA | 1.101 |
| | Lake Washington | NA | 0.914 ⁹ |
| | Green River | 0.162 | 0.914 ⁹ |
| North / Central B. C. | Yakoun, Nass, Skeena, Area 8 | NA | 0.626 |
| Washington Coastal Fall Naturals ³ | Hoko, Grays Harbor, Queets ² , Hoh ² , Quillayute ² | NA | 0.363 |
| Columbia River Falls ³ | Upriver Brights ² | NA | 0.523 |
| | Deschutes | NA | 0.523 |
| | Lewis ² | NA | 0.315 |
| Columbia R Summers ³ | Mid-Columbia Summers ² | NA | 0.335 |
| Far North Migrating OR Coastal Falls ³ | Nehalem ² , Siletz ² , Siuslaw ² | NA | 0.515 |

¹ Not available (NA) because of insufficient data (lack of stock specific tag codes, base period CWT recoveries, etc).

² Stock or stock group with a CTC agreed escapement goal.

³ Stock group listed in Annex 4, Chapter 3, Attachment V.

⁴ An inconsistency was discovered between the approaches used to calculate the model-based and CWT-based indices. The former included harvest rates for terminal sport while the latter did not. Terminal sport harvest rates are now included in the calculation of both indices. Further review is yet required to determine whether the base period terminal sport harvest rates obtained from analyses of Big Qualicum CWT recoveries adequately represent impacts that would have occurred on Cowichan Chinook.

⁵ Several problems have been identified in the approach previously used to calculate the CWT-based indices for Nanaimo Chinook. Until these problems are resolved, indices for this stock will not be reported.

⁶ Although model-based indices were previously calculated separately for Cowichan and Nanaimo, these did not adequately represent impacts on either LGS stock because the model-based data represent an aggregate of the two stocks and methods do not currently exist to correctly disaggregate these data for calculation of the ISBM values. Until such methods are developed, a single index value will be reported for both stocks.

⁷ The terminal sport harvest rates for Chilliwack Hatchery Chinook, the indicator stock, were removed from the calculation for the Harrison River naturals because sport harvest has been essentially zero on the natural population.

⁸ An inconsistency was discovered between the approaches used to calculate the model-based and CWT-based indices. The former included harvest rates for terminal sport while the latter did not. Terminal sport harvest rates are now included in the calculation of both indices. A more extended review of the indices for WCVI Chinook will be carried out to determine whether they adequately represent impacts on the WCVI wild aggregate.

⁹ For Canadian ISBM fisheries, Lake Washington and Green River the same distribution and Index value are assumed.

Table 5. U.S. 2004 ISBM indices based on CWT and the 2006 indices predicted from the PSC Chinook Model.

| Stock Group | Escapement Indicator Stock | U.S. ISBM Indices | |
|--|--|----------------------|------------------------|
| | | CWT Indices for 2004 | Model Indices for 2006 |
| Washington Coastal Fall Naturals | Hoko | NA ¹ | 0.442 |
| | Grays Harbor | 0.530 | 0.544 |
| | Queets ⁴ | 0.840 | 1.022 |
| | Hoh ⁴ | 1.220 | 1.493 |
| | Quillayute ⁴ | 1.150 | 0.673 |
| Columbia River Falls | Upriver Brights ⁴ | 1.740 | 0.814 |
| | Deschutes | 0.510 | 0.437 |
| | Lewis ⁴ | 0.170 | 1.861 |
| Puget Sound Natural Summer / Falls | Skagit | NA | 0.258 |
| | Stillaguamish | 0.10 | 0.493 |
| | Snohomish | NA | 0.199 |
| | Lake Washington | NA | 0.613 |
| | Green R | 1.010 | 0.361 |
| Fraser Late | Harrison River ⁴ | 0.320 | 0.787 |
| Columbia R Summers | Mid-Columbia Summers ⁴ | 2.690 | 0.696 |
| Far North Migrating OR Coastal Falls | Nehalem ⁴ | 1.800 | 1.912 |
| | Siletz ⁴ | 2.290 | 1.237 |
| | Siuslaw ⁴ | 1.030 | 1.095 |
| North Puget Sound Natural Springs | Nooksack | NA | 0.121 |
| | Skagit | NA | 0.161 |
| Lower Strait of Georgia ³ | Cowichan, | 7.250 | 0.271 |
| | Nanaimo | 7.250 | 0.271 |
| Upper Strait of Georgia ³ | Klinaklini, Kakweikan, Wakeman, Kingcome, Nimpkish | NA | NC ² |
| Fraser Early (spring and summers) ³ | Upper Fraser, Mid Fraser, Thompson | NA | 0.214 |
| West Coast Vancouver Island Falls ³ | WCVI (Artlish, Burman, Kauok, Tahsis, Tashish, Marble) | NA | 0.128 |
| North / Central B. C. ³ | Yakoun, Nass, Skeena, Area 8 | NA | NC |

¹ Not available (NA) because of insufficient data (lack of stock specific tag codes, base period CWT recoveries, etc).

² NC means that the current model assumes the stock is not caught in U.S. ISBM fisheries.

³ Stock group listed in Annex 4, Chapter 3, Attachment IV.

⁴ Stock with a CTC agreed escapement goal.

ESCAPEMENTS THROUGH 2005

The escapements of 50 naturally spawning escapement indicator stocks/stock aggregates are reviewed annually. Biologically-based escapement goals have been accepted by the CTC for 24 of the 50 escapement indicator stocks/stock aggregates. For 12 of these, the agency escapement goal is defined as a range; for the remaining 12, the escapement goal is the point estimate of S_{MSY} (escapement producing maximum sustained yield). In 2005, for stocks with an escapement goal range, escapements were within the range for eight stocks, above the range for one stock, and below the range for three stocks. For stocks with point estimate goals, escapements were above the goal for all 12 stocks. Data for stocks without accepted goals are presented to illustrate

trends in escapement. The CTC will continue to review escapement goals, as they are provided to the committee.

EXPLOITATION RATE ANALYSIS

There have been mark-selective fisheries (MSF) for Chinook salmon in the Strait of Juan de Fuca Washington sport fishery since 2003, in the Columbia River net fisheries since 2002, and in Columbia River spring Chinook sport fisheries since 2000. Double index tag (DIT) groups are used as a monitoring tool to test the hypothesis that there are differences between the marked and unmarked tagged groups due to MSFs and also to estimate mortalities of unmarked fish in MSFs.

A significant change in the ratio of unmarked to marked DIT groups at hatchery escapement can indicate that mark-selective fisheries have differentially impacted DIT pairs. Statistical Z-tests were used to compare the return rate of the marked and unmarked brood-age groups for seven Puget Sound DIT groups subject to MSFs in 2003 and 2004. Of the 52 tests for brood-age differences between marked and unmarked returns to the hatchery, only 6 were significant (Figure 3.15), and the actual calculated differences were small. This indicates that the Area 5 and 6 MSF did not result in significant differences in hatchery escapement of DIT groups. For this reason, the estimates of exploitation rate of marked tagged groups were used in CTC analyses this year.

1 CHINOOK CATCH

The June 30, 1999, Pacific Salmon Treaty (PST) Annexes and Related Agreements (Agreement) substantially changed the objectives and structure of the Pacific Salmon Commission's (PSC) Chinook salmon fisheries. The Agreement eliminated the previous ceiling and pass-through fisheries and replaced them with Aggregate Abundance Based Management (AABM) and Individual Stock Based Management (ISBM) fisheries. Chinook catches for the AABM fisheries are summarized in Tables 1.1-1.4, as well as Appendix A, and the ISBM catch in Appendices A.1-A.14.

Starting with the report CTC (2004a), the Chinook Technical Committee included estimates of incidental mortalities associated with landed catch for each component of each AABM fishery and most ISBM fisheries (CTC 2004b). Limited commentary on both AABM and ISBM fisheries is also provided.

1.1 REVIEW OF AABM FISHERIES

AABM fisheries for Chinook are managed to achieve a target catch corresponding to a target exploitation rate index and each year's abundance index (AI) in Table 1 of the Agreement. AABM fisheries are mixed stock salmon fisheries that intercept and harvest migratory Chinook from many stocks. The AABM fisheries (Annex IV, Chapter 3, paragraph 2) are:

- 1) Southeast Alaska (SEAK) All Gear,
- 2) Northern BC (NBC) Troll and Queen Charlotte Islands (QCI) sport, and
- 3) West Coast Vancouver Island (WCVI) Troll and Outside Sport.

Catches for these three fisheries are reported in Table 1.1.

Table 1.1. Annual catches and hatchery add-ons for the AABM fisheries, in thousands of Chinook salmon. The Treaty catches do not include the add-on or exclusions (see Section 1.1.1 and Appendix A.1). Notation is T for Troll, N for Net and S for sport.

| Year | SEAK (T, N, S) | | | NBC (T), QCI (S) | | WCVI (T, S) | |
|------|--------------------|--------------------------|--------------------|--------------------|----------|--------------------|----------|
| | Treaty Catch | | Hatchery Add-on | Treaty Catch | | Treaty Catch | |
| | Limit ¹ | Observed | | Limit ¹ | Observed | Limit ¹ | Observed |
| 1999 | 184.2 | 198.8 | 47.7 | 126.1 | 86.7 | 107.0 | 36.4 |
| 2000 | 178.5 | 186.5 | 74.3 | 123.5 | 31.9 | 86.2 | 101.4 |
| 2001 | 250.3 | 186.9 | 77.3 | 158.9 | 43.5 | 145.5 | 117.7 |
| 2002 | 371.9 | 357.1 | 68.2 | 237.8 | 150.1 | 196.8 | 165.0 |
| 2003 | 439.6 | 380.2 | 57.2 | 197.1 | 191.7 | 268.9 | 175.8 |
| 2004 | 418.3 | 428.8/433.4 ² | 72.0 | 267.0 | 241.5 | 209.6 | 216.6 |
| 2005 | 387.4 | 386.7 | 64.1 | 240.7 | 243.6 | 179.7 | 202.7 |

¹ Allowable treaty catches correspond to the postseason AIs for 1999-2005.

² The value on the left does not account for a terminal exclusion for the Stikine River, whereas the value on the right includes such terminal exclusion catch.

1.1.1 Southeast Alaska Fisheries

The SEAK Chinook fishery has been managed to achieve the annual all gear PSC allowable catch through a plan established by the Alaska Board of Fisheries. Once the all gear allowable catch is determined from the preseason AI each spring, this plan establishes gear quotas for the troll, net, and recreational fisheries. The allocation plan reserves 4.3% of the total PSC catch for purse seine, and 8,600 fish for combined set and drift gillnet fisheries. After the net quotas are subtracted, 80% of the remainder is reserved for troll gear and 20% for the recreational fishery. The recreational fishery is managed in-season with bag-limits and other constraints. Regulatory history and maps for each SEAK fishery are detailed in CTC (2004b).

In addition, the SEAK fisheries were managed for:

- 1) An Alaskan hatchery add-on estimated from coded-wire-tag (CWT) sampling, minus 5,000 base-period Alaska hatchery harvest. As a risk adjustment to account for sampling error, the lower bound of the 90% confidence interval is used as the estimate of Alaska hatchery harvest.
- 2) An exclusion of Situk stock catch in District 108, and exclusions of wild Chinook originating from the Taku and Stikine Rivers.
- 3) Compliance with provisions established by the National Marine Fisheries Service in accordance with the United States (U.S.) Endangered Species Act (ESA).
- 4) Consistency with the provisions of the PST as required by the Salmon Fishery Management Plan of the North Pacific Fishery Management Council that was established by the U.S. Magnuson-Stevens Act.

The all gear harvest in SEAK in 2005 was similar in magnitude to those for 2002 to 2004. The pre-season AI of 2.05 allowed an initial all-gear catch of 416,408 fish per the Agreement. The all gear harvest was 495,782 that resulted in a treaty catch of 386,707 (Table 1.1), an add-on of 64,102 and excluded catch of 44,973 Chinook salmon. A breakdown by gear for total catch, Alaskan hatchery contributions and terminal exclusions is detailed in Table 1.2. Historical harvests for 1975-2005 for SEAK are in Appendix A.1.

In February, 2005 an agreement was negotiated between the United States and Canada by the Transboundary Rivers Panel and approved by the PSC for directed harvest of wild Chinook salmon returning the Stikine and Taku Rivers (Annex IV, Paragraph 3). The agreement allowed for harvest sharing and exemption of the catches from harvest quotas above average base catches for the years 1985-2003. The harvest exemptions for transboundary rivers apply only to Stikine and Taku River fish harvested by the United States in Southeast Alaska Management Districts 108 and 111 and by Canada in the in-river fisheries on both rivers. The allowable catch (AC) tables and Base Terminal Run (BTR) calculations for both rivers are given in Appendix I.

Table 1.2. Harvest of Chinook salmon in SEAK by gear type in 2005.

| Gear | Total Harvest | Alaskan Hatchery Harvest | Alaskan Hatchery Add-on | Catch Exclusion¹ | Treaty Catch |
|----------------|----------------------|---------------------------------|--------------------------------|------------------------------------|---------------------|
| Troll | | | | | |
| Winter | 50,461 | 5,474 | 4,696 | 0 | 45,765 |
| Spring | 61,088 | 20,096 | 17,524 | 4,288 | 39,275 |
| Summer | 226,888 | 10,321 | 8,854 | 0 | 218,034 |
| Troll subtotal | 338,437 | 35,891 | 31,074 | 4,288 | 303,074 |
| | | | | | |
| Sport | 84,279 | 24,310 | 21,423 | 771 | 62,085 |
| | | | | | |
| Net | | | | | |
| Set Net | 718 | 0 | 0 | 0 | 718 |
| Driftnet | 52,481 | 5,927 | 5,199 | 39,914 | 7,369 |
| Seine | 19,867 | 6,867 | 6,406 | 0 | 13,461 |
| Net subtotal | 73,066 | 12,794 | 11,604 | 39,914 | 21,548 |
| | | | | | |
| Total | 495,782 | 72,994 | 64,102 | 44,973 | 386,707 |

¹ Exclusion catch claimed in 2005 is for the harvest sharing arrangement on the Taku and Stikine Rivers. There was no catch exclusion claimed on the Situk in 2005 as the catch did not reach the base level.

1.1.2 British Columbia Fisheries

Under the Agreement, the NBC AABM fishery is defined to include troll catch in Management or Statistical Areas 1-5 and sport catch in Areas 1 and 2 (QCI; Figure 1.1). The total AABM catch (troll plus QCI sport) in 2005 was 243,606 (Table 1.3). The WCVI AABM fishery includes the WCVI troll and the outside WCVI Chinook recreational fishery (defined below). The total AABM landed catch (First Nations, troll, and outside tidal sport) in 2005 was 202,662 Chinook (Table 1.3).




Figure 1.1. British Columbia fishery management areas.

1.1.2.1 NBC Troll Fishery Harvest

The NBC troll fishery was opened for Chinook fishing from October 1, 2004 to March 30, 2005 and from June 3 to September 30, 2005. A total of 174,806 Chinook were caught, with 5,043 caught in the winter fishery and 6,953 caught in a traditional style fishery from June 16 to July 17. A demonstration fishery was conducted to examine the application of individual transferable quotas in the troll fishery. A total of 161,321 Chinook were caught under the quota system from June 3 to September 30. A test fishery was also conducted in areas off the west coast of the Queen Charlotte Islands, which harvested 1,489 legal sized Chinook. These catches are included in the total for the troll fishery. The size limit was 67 cm. Barbless hooks and revival boxes were mandatory in the troll fishery. A ribbon boundary around Langara Island and from Skonun Point to Cape Knox on Graham Island excluded the commercial troll fishery from areas within one nautical mile (NM) of the shore June 3 to September 30.

Table 1.3. Summary of landed catch by gear for Canadian AABM fisheries in 2005.

| AABM Fishery | Troll | Sport | Total |
|---------------------|--------------|--------------|--------------|
| NBC | 174,806 | 68,800 | 243,606 |
| WCVI | 148,734 | 53,928 | 202,662 |

1.1.2.2 NBC and CBC Recreational Fishery Harvest

Although CBC is not part of the NBC AABM fishery, the tidal recreational fisheries in NBC and CBC (marine statistical Areas 1-11; Figure 1.1) are managed under one set of regulations (45 cm minimum size limit; two Chinook per day and four in possession; annual bag limit of 30). During the past decade, recreational fisheries in the marine areas of NBC and CBC have expanded substantially. Management of these marine recreational fisheries now recognizes two basic regions: QCI, and the coastal mainland. Only the QCI recreational catch is included in the AABM totals. Since 1995, catch in the QCI recreational fisheries has been estimated by creel surveys (supported by the Haida Nation), lodge logbook programs and independent observations by CDFO staff. Catch for this fishery in 2005 was 68,800 Chinook salmon. Thus, the total NBC AABM catch (troll plus sport) between October 1, 2004 and September 30, 2005 was 243,606 Chinook salmon (Table 1.3).

1.1.2.3 WCVI AABM

Under the 1999 PST Agreement, the WCVI AABM fishery includes the WCVI troll and the outside WCVI Chinook recreational fishery (defined below). The total AABM landed catch (First Nations, troll, and outside tidal sport) in 2005 was 202,662 Chinook (Table 1.3).

1.1.2.3.1 WCVI Troll Fishery Harvest

The AABM troll catch includes the commercial Area G troll catch and First Nations troll caught Chinook in Statistical Areas 21, 23-27, and 121-127 (Figure 1.1). In the 2005 season (October 1, 2004-September 30, 2005), the WCVI troll fishing opportunities were consistent with a CDFO commitment to evaluate winter fisheries as a means to improve the economic base for the fleet and local communities while increasing flexibility in harvest opportunities and reducing the harvest rates on stocks encountered in summer fisheries (Table 1.4). Troll fishery openings were shaped by conservation concerns for upper Fraser River, WCVI and Strait of Georgia (SG) Chinook and upper Fraser River and Thompson River coho. To protect early spring run upper Fraser Chinook, areas where they were known to be present were closed from mid-March to mid-April. To protect Thomson River coho, Chinook troll fisheries were closed after the middle of May. To protect SG Chinook offshore SWVI areas south of Estevan Point (areas 123 to 124) were closed the entire month of March, and from April 1 to 27. All WCVI harvest levels were reduced in May (26,655 in 2005; 51,486 in 2004). These management actions also give some protection to spring run U.S. Chinook stocks which have mature cohorts that may be present in the WCVI fishery in May. In the outside fishery, the southern bank area (Area 121) remained closed in 2005. The minimum size limit for troll-caught Chinook in all periods was 55 cm FL. The majority of catch from November through March came from Areas 23, 123, 125 and 126. WCVI troll fisheries were closed until mid-September to protect local WCVI Chinook stocks. The majority of the catch in September came from Area 126.

Table 1.4. Fishing periods and Chinook harvested and released during the 2005 accounting year in the WCVI troll fishery.

| Areas Open* | Fishing Period | Landed Catch | Sub-legal releases |
|--------------------|----------------------------------|---------------------|---------------------------|
| 123-127 | Oct. 1-2, 2004 | 11,256 | 978 |
| 23-27, 123-127 | Nov 1-4, 2004 | 8,057 | 1363 |
| 23-27, 123-127 | Dec 6, 13, 19, 2004 | 134 | 21 |
| 23-27, 123-127 | Jan 10-31, 2005 | 1,862 | 437 |
| 23-27, 123-127 | Feb 7-12, 20-22, 2005 | 5,650 | 513 |
| 25-27, 125-127 | Mar 1-14, 21-30, 2005 | 16,247 | 1,566 |
| 23, 25-27, 125-127 | Apr 1-27, 2005 | 39,269 | 1,724 |
| 23-27, 123-127 | Apr 28-30, 2005 | 17,794 | 1,102 |
| 23-27, 123-127 | May 1-2, 2005 | 12,197 | 669 |
| 23-27, 123-127 | May 12-13, 2005 | 14,458 | 1,105 |
| 23, 25 | Aug 9, 10, 16, 17, 23-26, 29, 30 | 184 | 0 |
| 26-27, 124-127 | Sep 17-21, 24-30, 2005 | 16,626 | 1,400 |
| | TOTAL | 143,734 | 10,878 |

¹ Troll fisheries were closed mid-May to mid-September to avoid encounters of Upper Fraser and Thompson River coho and WCVI Chinook.

The catch for 2005 Area G troll fisheries between October 1, 2004 and September 30, 2005 was 143,734 Chinook (Table 1.4). With the addition of an estimated 5,000 Chinook caught in WCVI First Nations troll fisheries in 2005, the total WCVI AABM troll catch for 2005 was 148,734 (Table 1.3) with 10,878 sublegal Chinook releases.

1.1.2.3.2 WCVI Recreational Fishery Harvest

The AABM recreational fishery includes all catch in northwest WCVI (Areas 25–27; Figure 1) between October 16 through June 30, and the catch outside one NM offshore from July 1 through October 15, plus all the catch in southwest WCVI (Areas 21–24) between October 16 through July 31, and outside one NM offshore from August 1 to October 15. Catch inside the surf line and outside the AABM periods specified above is included in ISBM fishery catch.

The outer WCVI sport fishery occurs primarily in the Barkley Sound, outer Clayoquot Sound, and Nootka Sound areas. The majority of fishing effort occurs from mid-July to September in north WCVI and August through mid-September in the south WCVI. Creel surveys are generally conducted from late May or early June to September 30. For the outside sport fishery the Chinook daily bag limit was two Chinook salmon greater than 45 cm.

Recreational effort in the AABM portion of the WCVI fishery was estimated at 36,256 boat trips in 2005. The 2005 WCVI AABM sport catch estimate during the creel period was 52,328 Chinook (Table 1.5). Catch rates were estimated from 10,532 interviews (12% of the estimated number of angling parties) at 19 landing sites from June 1 to September 30. No creel surveys occurred between the months of October and May, as effort is relatively low during this period. Catch for this period is estimated to be ~1,600 fish annually. This amount was added to the creel estimate to provide a total WCVI AABM sport catch estimate of 53,928.

Table 1.5. Outer WCVI AABM recreational fishery catches of Chinook by statistical area in 2005 representing catch during the creel survey periods only.

| Statistical areas | | | | | | |
|-------------------|--------|--------|--------|--------|--------|--------|
| 21/121 | 23/123 | 24/124 | 25/125 | 26/126 | 27/127 | Total |
| 11,440 | 29,540 | 6,551 | 1,560 | 1,997 | 1,240 | 52,328 |

1.2 ESTIMATES OF INCIDENTAL MORTALITIES IN AABM FISHERIES

1.2.1 SEAK Fisheries

Estimates of incidental mortality (IM) in SEAK fisheries are shown in Table 1.6. Estimates were available for all SEAK fisheries through 2005, except for the recreational fishery for which 2004 and 2005 data have not yet been tabulated. The IM for the troll and recreational fisheries were estimated from direct fishery observation programs. Estimates for the net fishery included IM for both seine and gillnet fisheries. For the seine fishery, estimates were based on regressions between landed catch in traditional fisheries and IM, from the 1985-1987 purse seine studies (CTC 2004c). For the gillnet fishery, drop-off mortality was estimated as a percentage of the landed catch using the regional-specific drop-off rate for SEAK (CTC 2004c).

Table 1.6. Estimated encounters and incidental mortality in SEAK troll, net and sport fisheries for 2003-2005. Mortality estimates of fish released in troll and sport fisheries include drop-off mortality. In the net fishery, 21"-28" fish from both retention and non-retention periods are included in the CNR numbers.

| Panel A - Troll and Sport Fisheries | | | | | | | | |
|-------------------------------------|------------|-------------------|----------|-------------|----------|-----------------|----------|----------|
| | | Troll | | | | Sport | | |
| | | Retention Fishery | | CNR Fishery | | Retention | Releases | |
| | | Legal | | | | Legal | | |
| Year | | Drop-off | Sublegal | Legal | Sublegal | Drop-off | Legal | Sublegal |
| 2003 | Encounters | NA ¹ | 39,821 | 34,262 | 19,703 | NA ¹ | 25,518 | 57,006 |
| 2003 | IM | 2,646 | 10,473 | 7,503 | 5,182 | 2,497 | 4,057 | 9,064 |
| 2004 | Encounters | NA ¹ | 18,161 | 71,834 | 34,980 | NA ¹ | | |
| 2004 | IM | 2,837 | 4,776 | 15,732 | 9,200 | 3,150 | | |
| 2005 | Encounters | NA ¹ | 31,660 | 49,430 | 24,346 | NA ¹ | | |
| 2005 | IM | 2,707 | 8,327 | 10,825 | 6,403 | 3,034 | | |

| Panel B - Net Fisheries and Total | | | | | | | | |
|-----------------------------------|------------|---------------|-------------|---------|-----------------|----------------------|----------|--|
| | | Net Fisheries | | | | | | |
| | | Seine | | | Gillnet | Total | | |
| | | Retention | CNR Fishery | | Legal | Incidental Mortality | | |
| Year | | < 21" | > 28" | 21"-28" | Drop-off | Legal | Sublegal | |
| 2003 | Encounters | 1,107 | 16,081 | 53,188 | NA ¹ | | | |
| 2003 | IM | 1,107 | 8,202 | 39,093 | 305 | 25,210 | 64,919 | |
| 2004 | Encounters | 591 | 28,700 | 94,922 | NA ¹ | | | |
| 2004 | IM | 591 | 14,637 | 69,767 | 488 | 36,844 | 84,334 | |
| 2005 | Encounters | 663 | 13,225 | 43,841 | NA ¹ | | | |
| 2005 | IM | 663 | 6,760 | 32,223 | 1,064 | 24,390 | 47,616 | |

¹ Drop-off mortality is computed from landed catch times a percentage that incorporates a gear-specific encounter ratio and release mortality rate.

1.2.2 British Columbia Fisheries

1.2.2.1 NBC Fisheries

Table 1.7 summarizes encounter and IM estimates for the NBC AABM fisheries from 2002 to 2005 by size class during retention and Chinook Non-retention (CNR) fishing periods. Encounters for the NBC troll fishery are based on phone-in hails. Encounters for the QCI sport fishery are based on creel survey and logbook programs. The table presents IM estimates using size specific rates from the CTC (1997). The estimated total mortality of Chinook salmon in the NBC AABM fisheries in 2005 was 267,425 nominal fish, including 243,606 fish in the landed catch and 23,819 fish from IM (Table 1.7).

Table 1.7. Estimated encounters and incidental mortalities (nominal fish) in NBC AABM troll and sport fisheries for 2002-2005. Mortality estimates of fish released in troll and sport fisheries include drop-off mortality.

| Year | | Troll | | | | Sport | | Total Incidental Mortalities | |
|------|------------|---------------------------|-------------------|-------------|----------|---------------------------|-----------------------|------------------------------|----------|
| | | Retention Fishery | | CNR Fishery | | Retention | Releases ² | | |
| | | Legal & Sublegal Drop-off | Sublegal releases | Legal | Sublegal | Legal & Sublegal Drop-off | Legal | Legal | Sublegal |
| 2002 | Encounters | NA ¹ | 2,608 | 5,109 | 129 | NA ¹ | 42,226 | | |
| | IM | 1,752 | 618 | 1,032 | 31 | 3,250 | 8,107 | 14,098 | 692 |
| 2003 | Encounters | NA ¹ | 1,721 | 11,798 | 148 | NA ¹ | 47,549 | | |
| | IM | 2,335 | 408 | 2,383 | 35 | 3,747 | 9,129 | 17,566 | 472 |
| 2004 | Encounters | NA ¹ | 2,605 | 31,460 | 489 | NA ¹ | 116,741 | | |
| | IM | 2,848 | 617 | 6,355 | 116 | 5,106 | 22,414 | 36,511 | 725 |
| 2005 | Encounters | NA ¹ | 1,009 | 20,414 | 118 | NA ¹ | 60,987 | | |
| | IM | 2,972 | 239 | 4,124 | 28 | 4,747 | 16,457 | 23,535 | 284 |

¹ Drop-off mortality is computed from landed catch times a percentage that incorporates a gear-specific encounter ratio and release mortality rate.

² Releases are reported as 'mixed' sizes. However, since >90% of such releases are legal-sized, all reported releases were considered to be legal-sized for the purpose of estimating incidental mortality.

1.2.2.2 WCVI Fishery

The estimated total mortality of Chinook salmon in the WCVI AABM fisheries in 2005 was 214,467 nominal fish, including 202,662 fish in the landed catch and 11,805 fish from IM (Table 1.8). The estimated IM included 8,286 legal and 3,445 sublegal fish in nominal numbers of fish. The estimates for the commercial troll fisheries in 2005 are from direct fishery observations programs. Table 1.8 summarizes 2003-2005 encounter and IM estimates for these fisheries by size class during retention. In 2004 and 2005 there were no CNR fishing periods in the AABM fishery.

Table 1.8. Estimated encounters and incidental mortalities (nominal fish) in WCVI troll and sport AABM fisheries for 2003-2005. Mortality estimates of fish released in troll and sport fisheries include drop-off mortality.

| Year | | Troll | | | | Sport | | | Total Incidental Mortalities | |
|------|------------|-------------------|----------|-------------|----------|-----------------|----------|----------|------------------------------|--------------------|
| | | Retention Fishery | | CNR Fishery | | Retention | Releases | | | |
| | | Legal | | Legal | | Legal | Legal | | | |
| | | Drop-off | Sublegal | Legal | Sublegal | Drop-off | Legal | Sublegal | Legal | Sublegal |
| 2003 | Encounters | NA ¹ | 15,479 | 63 | 7 | NA ¹ | 11,016 | 8,073 | | |
| | IM | 2,581 | 3,793 | 13 | 0 | 1,656 | 2,115 | 1,550 | 6,352 | 5,343 |
| 2004 | Encounters | NA ¹ | 10,430 | 0 | 0 | NA ¹ | 16,449 | 5,680 | | |
| | IM | 2,786 | 2,461 | 0 | 0 | 2,723 | 2,023 | 1,091 | 7,532 | 3,510 ² |
| 2005 | Encounters | NA ¹ | 10,878 | 0 | 0 | NA ¹ | 19,319 | 4,571 | | |
| | IM | 2,300 | 2,567 | 0 | 0 | 3,610 | 2,376 | 878 | 8,286 | 3,445 |

¹ Legal drop-off mortality is computed from landed catch, incorporating both an encounter ratio and a mortality rate.

² Sublegal dropoffs are included with sublegal incidental release mortalities

1.3 REVIEW OF ISBM FISHERIES

1.3.1 Canadian ISBM Fisheries

ISBM fisheries include all fisheries that harvest or release Chinook salmon in British Columbia under PST jurisdiction outside areas governed by AABM fisheries. In 2005, 271,821 Chinook were harvested in Canadian ISBM fisheries in British Columbia and Canadian sections of the Alsek, Taku and Stikine Transboundary rivers. Total estimated IM in the Canadian ISBM fisheries in 2005 was 36,873 legal and sublegal sized Chinook. The distribution of the landed catches and estimated incidental mortalities in Canadian ISBM fisheries are presented in Table 1.9. Historical catches in Canadian fisheries may be found in Appendixes A2 through A7.

Table 1.9. Landed catch and incidental mortalities in Canadian ISBM fisheries for 2005.

| Region | Gear Type | Landed Catch | Releases | | Incidental Mortalities ¹ | | Total Nominal Mortality |
|----------------------------|-------------------------|----------------|---------------|---------------|-------------------------------------|-----------------------|-------------------------|
| | | | Legal | Sublegal | Legal ² | Sublegal ³ | |
| Transboundary Rivers | Gillnet | 9,515 | NA | NA | 438 | NA | 9,953 |
| (Taku, Stikine, Alsek) | Recreational | 35 | NA | NA | 2 | NA | 37 |
| | FN | 1,119 | NA | NA | 51 | NA | 1,170 |
| <i>Regional Total</i> | | <i>10,669</i> | <i>0</i> | <i>0</i> | <i>492</i> | <i>0</i> | <i>11,161</i> |
| Northern BC ⁴ | Gillnet | 5,518 | 695 | NA | 879 | NA | 6,397 |
| | Seine | 0 | 4,807 | NA | 3,461 | NA | 3,461 |
| | Tyee Test Fishery | 1,332 | 0 | 0 | 61 | 0 | 1,393 |
| | Tidal Sport | NA | NA | NA | NA | NA | NA |
| | Non-tidal Sport | NA | NA | NA | NA | NA | NA |
| | FSC(Tidal & Non-tidal) | 17,553 | NA | NA | 1,211 | NA | 18,764 |
| <i>Regional Total</i> | | <i>24,403</i> | <i>5,502</i> | <i>0</i> | <i>5,613</i> | <i>0</i> | <i>30,016</i> |
| Central Coast ⁵ | Gillnet | 6,323 | 21 | NA | 310 | NA | 6,633 |
| | Seine | 0 | 15,260 | NA | 10,987 | NA | 10,987 |
| | Tidal Sport | 9,017 | 302 | NA | 659 | NA | 9,676 |
| | Non-tidal Sport | 809 | NA | NA | 56 | NA | 865 |
| | FSC(Tidal & Non-tidal) | 4,180 | NA | NA | 288 | NA | 4,468 |
| <i>Regional Total</i> | | <i>20,329</i> | <i>15,583</i> | <i>0</i> | <i>12,301</i> | <i>0</i> | <i>32,630</i> |
| WCVI terminal | Gillnet | 18,705 | 0 | 213 | 860 | 192 | 19,757 |
| | Seine | 4,894 | NA | 141 | NA | 127 | 5,021 |
| | Tidal Sport | 41,710 | 5,336 | 4,199 | 3,534 | 806 | 46,051 |
| | Non-tidal Sport | 6,225 | NA | NA | 430 | NA | 6,655 |
| | FSC(Tidal & Non-tidal) | 35,000 | NA | NA | 1,610 | NA | 36,610 |
| <i>Regional Total</i> | | <i>106,534</i> | <i>5,336</i> | <i>4,553</i> | <i>6,434</i> | <i>1,125</i> | <i>114,093</i> |
| Johnstone Strait | Seine | 247 | 2,817 | NA | 2,028 | NA | 2,275 |
| | Gillnet | 44 | 10 | 9 | 11 | 8 | 63 |
| | Tidal Sport | 12,009 | 3,606 | 5,916 | 1,272 | 1,136 | 14,417 |
| | FSC(Tidal & Non-tidal) | NA | NA | NA | NA | NA | NA |
| <i>Regional Total</i> | | <i>12,300</i> | <i>6,433</i> | <i>5,925</i> | <i>3,311</i> | <i>1,144</i> | <i>16,755</i> |
| Georgia Strait | Seine | 2 | 6 | NA | 4 | NA | 6 |
| | Tidal Sport | 12,298 | 1,077 | 9,025 | 981 | 1,733 | 15,012 |
| | FSC (Tidal & Non-tidal) | 607 | NA | NA | NA | NA | 607 |
| <i>Regional Total</i> | | <i>12,907</i> | <i>1,083</i> | <i>9,025</i> | <i>985</i> | <i>1,733</i> | <i>15,625</i> |
| Juan de Fuca Strait | Seine | 19 | 615 | 22 | 443 | 19 | 481 |
| | Gillnet | 134 | 50 | 57 | 51 | 51 | 236 |
| | Tidal Sport | 30,480 | 6,908 | 4,949 | 2,953 | 950 | 34,383 |
| | FSC (Tidal & Non-tidal) | NA | NA | NA | NA | NA | NA |
| <i>Regional Total</i> | | <i>30,633</i> | <i>7,573</i> | <i>5,028</i> | <i>3,447</i> | <i>1,020</i> | <i>35,100</i> |
| Fraser River | Gillnet | 5,296 | NA | NA | 244 | NA | 5,540 |
| | Sport (mainstem+tribs) | 21,831 | 10,588 | 2,734 | 2,809 | 525 | 25,165 |
| | FSC (Tidal & Non-tidal) | 26,919 | NA | NA | 1,238 | NA | 28,157 |
| <i>Regional Total</i> | | <i>54,046</i> | <i>10,588</i> | <i>2,734</i> | <i>4,291</i> | <i>525</i> | <i>58,862</i> |
| Grand Total | | 271,821 | 52,098 | 27,265 | 36,873 | 5,547 | 314,241 |

¹ Includes drop-off and release mortalities in both retention and Chinook non-retention fisheries.

² In Chinook non-retention fisheries, all releases were assumed to be legal size as the sizes were unknown. If no release information is available, IM represents dropoff mortality only.

³ Minimum size limits for sport catch were 45 cm in Juan de Fuca Strait and 62 cm elsewhere.

⁴Includes areas 1-5

⁵Includes areas 6-10

1.3.2 Southern U.S. Fisheries Harvest

Southern U.S. fisheries of interest to the PSC, generally those north of Cape Falcon, Oregon, are managed in accordance with legal obligations stemming from treaties between Indian tribes and the United States. In 1974, *U.S. v Washington* set forth sharing obligations to meet Treaty fishing rights in western Washington. Treaty rights of Columbia River tribes were defined by *U.S. v Oregon*, and the Columbia River Fisheries Management Plan was implemented in 1977. In reporting these fisheries, fishermen are termed “treaty” if they are fishing under the Native Treaty fishing rights and “non treaty” otherwise. As specified in the 1999 agreement, all southern U.S. fisheries are ISBM fisheries. Historical catches in these fisheries may be found in Appendices A.8 through A.14.

1.3.2.1 Strait of Juan de Fuca and the San Juan Islands

The preliminary estimate of the 2005 Chinook catch in Strait of Juan de Fuca tribal net fisheries directed at sockeye salmon is 170. An additional five Chinook were taken during the coho management period. The preliminary estimate of the 2005 Chinook catch in the San Juan Islands tribal net fishery directed at sockeye salmon is 4,306. Non-treaty landings totaled about 162 Chinook. The preliminary estimate of the 2005 Strait of Juan de Fuca treaty troll fishery is 5,344 Chinook through December. The catch estimate does not include catches from Area 4B during the May-September PFMC management period. These are included in the North of Cape Falcon troll summary. Historic catch estimates are provided in Appendices A.8 and A.9 for the Strait of Juan de Fuca and San Juan areas respectively.

1.3.2.2 Puget Sound

The preliminary estimate of the 2005 tribal and non-tribal net fishery harvests in Puget Sound marine areas is 58,174 Chinook, mostly taken in terminal areas where harvestable abundance was identified. Additional tribal net harvest occurred in freshwater fisheries with a preliminary estimate of 19,395. Estimates of the sport catch in 2005 are not yet available. Historic catch tables for Puget Sound exclusive of the San Juans are provided in Appendix A.10.

1.3.2.3 Washington Coast

Tribal commercial and ceremonial and subsistence fisheries harvested a total of 11,709 Chinook in north coastal rivers (Quinault, Queets, Hoh, Quillayute) in 2005. An additional 4,224 Chinook were harvested by the Makah tribal fisheries in the Waatch and Sooes rivers.

Harvest in Grays Harbor includes catch from both the Humptulips and Chehalis rivers. The 2005 tribal net fisheries harvested an estimated 2,493 Chinook. The 2005 non-Indian commercial net harvest in Grays Harbor was only about 91 Chinook. Approximately 6,372 Chinook were harvested by non-Indian commercial net fisheries in Willapa Bay in 2005.

From Grays Harbor north, recreational fisheries were implemented based upon pre-season tribal-state agreements and were subject to in-season adjustment. Estimates of sport fishery catches for Washington coastal terminal fishing areas in 2005 are not available. Historic catch estimates for Washington Coastal inside fisheries are shown in Appendix A.11.

Ocean fisheries off the coasts of Washington and Oregon are managed under regulations recommended by the Pacific Fishery Management Council. The estimated catch of Chinook salmon in commercial troll fisheries from Cape Falcon to the U.S.-Canada border in 2005 was

87,126 for both treaty and non-treaty fisheries combined. Estimated catch in the ocean recreational fishery north of Cape Falcon in 2005 was 40,011 Chinook. Historic catch estimates for U.S. ocean fisheries north of Cape Falcon are shown in Appendix A.12.

1.3.2.4 Columbia River

Chinook from the Columbia River are divided into eight stock groups for management purposes. These groups are delineated by run timing and area of origin: (1) spring run originating below Bonneville Dam; (2) spring run originating above Bonneville Dam; (3) summer run originating above Bonneville Dam; (4) fall run returning to Spring Creek Hatchery; (5) fall run originating in hatchery complexes below Bonneville Dam; (6) wild fall run originating below Bonneville Dam; (7) upriver bright fall run; and (8) mid-Columbia bright fall hatchery fish.

In 2005, the total annual harvest for all fisheries (spring, summer and fall) in the Columbia River basin was 263,615 Chinook, which included non-Indian and treaty-Indian commercial net harvest of 151,846 recreational harvest of 81,725, Indian ceremonial and subsistence harvest of 7,960, and 22,084 non-ticket sales. Historic catch estimates for Columbia River fisheries are found in Appendix A.13.

1.3.2.5 Ocean Fisheries, Cape Falcon to Humbug Mountain

Most harvest in ocean fisheries off Oregon's coast is comprised of a mixture of southern Chinook stocks not included in the PSC agreement. These stocks do not migrate north into the PSC jurisdiction to any great extent. Some stocks originating from Oregon coastal streams do migrate into PSC fisheries, including the North Oregon Coastal (NOC) and Mid-Oregon Coastal (MOC) stock aggregates. The NOC stocks are harvested only incidentally in Oregon ocean fisheries, while the contribution of MOC stocks to Oregon ocean fisheries is believed to be much greater. Catch statistics are readily available only for a terminal area troll fishery on one MOC stock at the mouth of the Elk River. Late season (October-December) troll catch in the Elk River terminal troll fishery in 2005 was 1,956 Chinook.

Recreational catch of these two stock groups occurs primarily in estuary and freshwater areas as mature fish return to spawn and is reported through a "punch card" accounting system. These data are only available more than two years after the current season. Therefore, we can only report the riverine and estuarine sport catch through 2004 for the NOC and MOC groups. The 2004 punch card estimate of estuary and freshwater catch for the NOC and MOC groups is 71,726 Chinook. Historic catch estimates for the Elk River troll fishery and the estuary and freshwater sport fisheries targeting on MOC and NOC stocks are shown in Appendix A.14.

1.4 ESTIMATES OF INCIDENTAL MORTALITY FOR SOUTHERN U.S. FISHERIES

Table 1.10 shows estimates of incidental mortalities for Washington Coastal and Puget Sound fisheries. Sources of estimates are shown in the table footnotes. No estimates of incidental mortalities were provided for 2005 for ocean fishery south of Cape Falcon or Columbia River fisheries.

Table 1.10. Estimated incidental mortality in Southern US troll, net, and sport fisheries for 2005.

| Fishery | Troll | Net¹ | Sport |
|------------------------|---------------------|------------------------|--------------------|
| Strait of Juan de Fuca | 868 ² | 5 | NA |
| San Juan Islands | 0 | 134 | NA |
| Puget Sound | 0 | 2,327 | NA |
| Washington Coast | 0 | 747 | NA |
| North of Cape Falcon | 15,700 ³ | 0 | 5,200 ³ |

¹ Assume 3% net dropout rate.

² Estimates from FRAM.

³ Estimates from direct observations.

2 ESCAPEMENTS THROUGH 2005

2.1 INTRODUCTION

The Agreement (Pacific Salmon Treaty Fishing Annexes & Related Agreements, June 30, 1999) established a Chinook management program that:

“introduces harvest regimes that are based on estimates of Chinook abundance, that are responsive to changes in Chinook production, that take into account all fishery induced mortalities and that are designed to meet MSY or other agreed biologically-based escapement objectives”

This chapter compares annual escapement estimates with maximum sustained yield (MSY) or other accepted biologically-based escapement goals established for Chinook stocks. The CTC has reviewed and accepted escapement goals for 24 stocks included in this report. For these stocks, the CTC can evaluate stock status in relation to the accepted goals. For stocks without accepted goals, the CTC must rely on the time series of escapement data and the agency commentary for the individual stocks to provide a perspective on stock status and escapement trends.

This year the CTC is presenting this information in an abbreviated format. Previous annual reports included a section on the framework used for escapement assessments and for each stock, narratives were included that had a description of escapement methodology, escapement goal basis and agency comments. In this report these narratives have been replaced with a commentary that updates the information with 2005 escapements and any changes from the previous report. For a detailed description of the framework used for escapement assessment and stock narratives, please refer to the 2004 Catch and Escapement Report (TCCHINOOK (05-2)).

2.1.1 MSY or Biologically-Based Escapement Goals

2.1.1.1 Origin of Goals

Escapement goals accepted by the CTC were based on analyses that followed the guidelines developed in the CTC escapement goal report (CTC 1999). In the stock-specific narratives presented with the escapement graphs, the agencies may refer to agency goals, but only CTC-accepted escapement goals and ranges (in gray shading) are shown on the escapement graphs and used for evaluation. Table 2-1 presents the status of escapement goal reviews by the CTC for stocks that have been identified as escapement indicator stocks.

Table 2.1. PSC Chinook escapement indicator stocks, where shading indicates that there is not a CTC accepted escapement goal for PSC assessment of stock status.

| Presence in Treaty Attachments | | | | | Stock Group In Att. I-V | Escapement Indicator | Region | Run |
|--------------------------------|-------------|------|------------|-------------|-------------------------------------|--|--------------|---------------|
| SEAK | NBC/ QCI | WCVI | BC ISBM | SUS ISBM | | | | |
| ✓ | | | | | | Situk | Yakutat | Spring |
| ✓ | | | | | | Alsek | Yakutat | Spring |
| ✓ | | | | | | Taku | TBR | Spring |
| ✓ | | | | | | Stikine | TBR | Spring |
| ✓ | | | | | | Chilkat | N. Inside | Spring |
| ✓ | | | | | | King Salmon | N. Inside | Spring |
| ✓ | | | | | | Andrew Creek | C. Inside | Spring |
| ✓ | | | | | | Unuk | S. Inside | Spring |
| ✓ | | | | | | Chickamin | S. Inside | Spring |
| ✓ | | | | | | Blossom | S. Inside | Spring |
| ✓ | | | | | | Keta | S. Inside | Spring |
| ✓ | ✓ | | ✓ | | Northern/Central B.C. | Yakoun | NBC-Area 1 | Summer |
| ✓ | ✓ | | ✓ | | Northern/Central B.C | Nass | NBC-Area 3 | Spring/Summer |
| ✓ | ✓ | | ✓ | | Northern/Central B.C | Skeena | NBC-Area 4 | Spring/Summer |
| | | | ✓ | | Northern/Central B.C. | Dean | CBC-Area 8 | Spring |
| | | | | | | Rivers Inlet | CBC-Area 9 | Spring/Summer |
| ✓ | ✓ | | ✓ | | WCVI Falls | Artlish, Burman, Kaouk, Tahsis, Tashish, Marble | WCVI | Fall |
| ✓ | ✓ | | ✓ | | Upper Strait of Georgia | Klinaklini, Kakwiekan, Wakeman, Kingcome, Nimpkish | UGS | Sum/Fall |
| | | | ✓ | | Lower Strait of Georgia | Cowichan/Nanaimo ² | LGS | Fall |
| ✓ | ✓ | | ✓ | | Fraser Early ¹ (Spr/Sum) | Fraser Spring 1.3 | Fraser River | Spring |
| ✓ | ✓ | | ✓ | | Fraser Early ¹ (Spr/Sum) | Fraser Spring 1.2 | Fraser River | Spring |
| ✓ | ✓ | | ✓ | | Fraser Early ¹ (Spr/Sum) | Fraser Summer 1.3 | Fraser River | Summer |
| ✓ | ✓ | | ✓ | | Fraser Early ¹ (Spr/Sum) | Fraser Summer 0.3 | Fraser River | Summer |
| | | ✓ | ✓ | ✓ | Fraser Late | Harrison | Fraser River | Fall |
| | | | ✓ | ✓ | N. P.S. Natural Springs | Nooksack | NC/PS | Spring |
| | | | ✓ | ✓ | N. P.S. Natural Springs | Skagit Spring | NC/PS | Spring |
| | | ✓ | ✓ | ✓ | P.S. Natural Summer/Falls | Skagit Summer/Fall | NC/PS | Summer/Fall |
| | | ✓ | ✓ | ✓ | P.S. Natural Summer/Falls | Stillaguamish | NC/PS | Summer/Fall |
| | | ✓ | ✓ | ✓ | P.S. Natural Summer/Falls | Snohomish | NC/PS | Summer/Fall |
| | | ✓ | ✓ | ✓ | P.S. Natural Summer/Falls | Lake Washington | NC/PS | Summer/Fall |
| | | ✓ | ✓ | ✓ | P.S. Natural Summer/Falls | Green | NC/PS | Summer/Fall |

-continued-

Table 2.1. Continued.

| Presence in Treaty Attachments | | | | | Stock Group In Att. I-V | Escapement Indicator | Region | Run |
|--------------------------------|-------------|------|------------|-------------|----------------------------|-------------------------|---------|--------|
| SEAK | NBC/ QCI | WCVI | BC ISBM | SUS ISBM | | | | |
| ✓ | ✓ | | | ✓ | WA Coastal Fall Natural | Hoko | WAC/JDF | Fall |
| | | | | | | Quillayute Summer | WAC/JDF | Summer |
| ✓ | ✓ | | | ✓ | WA Coastal Fall Natural | Quillayute Fall | WAC/JDF | Fall |
| | | | | | | Hoh Spring/Summer | WAC/JDF | Summer |
| ✓ | ✓ | | | ✓ | WA Coastal Fall Natural | Hoh Fall | WAC/JDF | Fall |
| | | | | | | Queets Spring/Summer | WAC/JDF | Summer |
| ✓ | ✓ | | | ✓ | WA Coastal Fall Natural | Queets Fall | WAC/JDF | Fall |
| | | | | | | Grays Harbor Spring | WAC/JDF | Spring |
| ✓ | ✓ | | | ✓ | WA Coastal Fall Natural | Grays Harbor Fall | WAC/JDF | Fall |
| | | | | | | Col. Upriver Spring | CR | Spring |
| ✓ | ✓ | ✓ | | ✓ | Col. Upriver Summers | Mid-Columbia Summers | CR | Summer |
| ✓ | ✓ | ✓ | | ✓ | Columbia River Falls | Col. Upriver Bright | CR | Fall |
| ✓ | ✓ | ✓ | | ✓ | Columbia River Falls | Lewis | CR | Fall |
| ✓ | ✓ | ✓ | | ✓ | Columbia River Falls | Deschutes | CR | Fall |
| ✓ | ✓ | | | ✓ | Far N. Migrating OR Coast. | Nehalem | NOC | Fall |
| ✓ | ✓ | | | ✓ | Far N. Migrating OR Coast. | Siletz | NOC | Fall |
| ✓ | ✓ | | | ✓ | Far N. Migrating OR Coast. | Siuslaw | NOC | Fall |
| | | | | | | Umpqua | MOC | Fall |
| | | | | | | Mid South OR | MOC | Fall |

¹ The escapement indicator stocks listed in the Annex tables for this group are Upper Fraser, Middle Fraser, and Thompson. The Fraser spring/summer group is split into these 4 escapement indicators to represent the stock group by life history type rather than geographically.

² An escapement goal was established for the Cowichan in 2005; a goal for Nanaimo is still pending.

2.2 ESCAPEMENT ASSESSMENT

The Agreement directs the CTC to “report annually on the escapement of naturally spawning Chinook stocks in relation to the agreed escapement objectives referred to below, evaluate trends in the status of stocks, and report on progress in rebuilding of naturally spawning Chinook stocks” (Annex IV, Chapter 3, paragraph 1.b.iii). In this report, escapement assessments include stock specific graphs of escapements and commentary, presented to provide a perspective on stock status and escapement trends through 2005. More detailed commentary for each stock can be found in previous CTC catch and escapement reports, e.g. CTC (2005).

The escapement goals and 2005 escapements for the 24 stocks with CTC accepted escapement goals are listed in Table 2-2. For 12 of these stocks, the agency escapement goal is defined as a range; for the remaining 12 stocks, the escapement goal is defined as a point estimate. In 2005, escapements were within the goal range for eight stocks, above the range or S_{MSY} point estimate for 13 stocks, and below the goal for three stocks.

Table 2.2. Escapement goals and 2005 escapements for PSC Chinook escapement indicator stocks with biologically-based goals accepted by the CTC.

| Stock | Region | Stock Group | Escapement Goal | 2005 Escapement |
|--------------------------|---------------|--------------------|------------------------|------------------------|
| Situk | SEAK | Yakutat | 500-1,000 | 613 |
| Alsek (Klukshu index) | SEAK/TBR | Yakutat | 1,100-2,300 | 963 |
| Chilkat | SEAK | Northern Inside | 1,750-3,500 | 3,366 |
| Taku | SEAK/TBR | TBR | 30,000-55,000 | 69,007 |
| Stikine | SEAK/TBR | TBR | 14,000-28,000 | 44,033 |
| King Salmon | SEAK | Northern Inside | 120-240 | 141 |
| Andrew Creek | SEAK | Central Inside | 650-1,500 | 1,190 |
| Unuk (survey index) | SEAK | Southern Inside | 650-1,400 | 929 |
| Chickamin (survey index) | SEAK | Southern Inside | 450-900 | 924 |
| Blossom (survey index) | SEAK | Southern Inside | 250-500 | 445 |
| Keta (survey index) | SEAK | Southern Inside | 250-500 | 497 |
| Harrison | BC | Fraser River | 75,100-98,500 | 86,730 |
| Cowichan | BC | Lower Georgia St. | 6,500 | 1,572 |
| Mid Col. Upr. Summer | CR | Columbia River | 17,857 | 45,874 |
| Col. Upriver Brights | CR | Columbia River | 40,000 | 112,679 |
| Lewis | CR | Columbia River | 5,700 | 11,348 |
| Quillayute Fall | WAC | WA Coast | 3,000 | 6,406 |
| Queets Spring/Summer | WAC | WA Coast | 700 | 294 |
| Queets Fall | WAC | WA Coast | 2,500 | 2,931 |
| Hoh Spring/Summer | WAC | WA Coast | 900 | 1,193 |
| Hoh Fall | WAC | WA Coast | 1,200 | 4,180 |
| Nehalem | ORC | NOC | 6,989 | 7,038 |
| Siletz | ORC | NOC | 2,944 | 6,426 |
| Siuslaw | ORC | NOC | 12,925 | 16,619 |

The CTC has now assessed the status of stocks with CTC-accepted goals for return years 1999-2005. Over this time period, the number of stocks with CTC-accepted goals has increased from 16 to 24 (Figure 2.1). The percentage of stocks below escapement goals or goal ranges has varied over these years from 4% to 19%, and was 12.5% for 2005 escapements.




Figure 2.1. Number and status of stocks with CTC-accepted escapement goals for years 1999-2005.


2.3 STOCK SPECIFIC GRAPHS AND COMMENTARIES

Graphs of time series of escapements and terminal runs for Chinook stocks are included in sections for Alaska, Canada, and Washington/Columbia River/Oregon. A limited commentary is also provided for each stock; more detail on historical assessments and escapement goals for individual stocks is available in CTC (2005a). Each graph contains the name of the stock and the type of data depicted (total escapement, index counts, terminal runs, etc.). For the graphs that include estimates of the terminal run size, the harvests in terminal runs include both jacks and adults in some cases, whereas the escapement is usually reported in adults. The x-axis represents calendar years. All escapement goals accepted by the CTC are shown except for the LGS stock group because this group includes both the Cowichan and Nanaimo stocks and only the Cowichan has a CTC accepted goal. Historical escapement and terminal run data are provided for SEAK stocks in Appendix B.1, for Canadian stocks in Appendix B.2, for Puget Sound in Appendix B.3, Washington Coastal stocks in Appendix B.4, for Columbia River stocks in Appendix B.5 and Oregon Coastal stocks in Appendix B.6.


2.3.1 SEAK/TBR Stocks

Of the 11 SEAK/TBR stocks included in the escapement assessment, the Situk, Chilkat, Taku, King Salmon, and Stikine rivers and Andrew Creek include estimates of total escapement of


large fish, Chinook salmon > 659 mm mid-eye to fork (MEF) length. Escapement estimates for the Alsek, Unuk, Chickamin, Blossom, and Keta rivers are index counts of large fish from a weir on the Alsek River and foot/aerial helicopter surveys on the other four rivers that represent a fraction of the total escapement. Except for the Chilkat River, survey methods have been standardized for all systems since 1975. The assessment of Chilkat River Chinook salmon was standardized in 1991 as an annual mark-recapture estimate of escapement. Escapement goals have been defined as a range for the SEAK/TBR stocks, shown by the grey shaded area on the graphs.




Commentary The Situk River is a small non-glacial system that supports a moderate run of outside-rearing Chinook salmon. Escapements are based on weir counts minus upstream sport fishery harvests (if any) estimated from an on-site creel survey and a postseason mail-out survey. The weir has been operated annually since 1976, and was also operated from 1928-1955.




Commentary: The Alsek River is large transboundary glacial system that supports a moderate run of outside-rearing Chinook salmon. Since 1976 index escapements (shown above) have been determined using a weir operated at the Klukshu River.




Commentary: The Taku River is a large transboundary glacial system that supports a large run of outside-rearing Chinook salmon. In 1989, 1990, and 1995-2005 escapement was determined using mark-recapture methods. In other years since 1975, aerial counts were expanded by a factor of 5.2, the 5-year average of the ratio of the mark-recapture estimates to aerial survey counts (McPherson et al. 2000).




Commentary: The Stikine River is a large transboundary glacial system that supports a large run of outside-rearing Chinook salmon. From 1975 through 1984 index escapements were made using survey counts and since 1985 counts were made using a weir at the Little Tahltan River. Since 1996 mark-recapture experiments were performed indicating the index escapements represented 17% to 20% of the total escapement (Pahlke and Etherton 1999).




Commentary The Chiklat River is a moderate-sized glacial system moderate run of inside-rearing Chinook salmon. Since 1991, escapements have been estimated using mark-recapture methods (Ericksen and McPherson 2003). The current biological escapement goal of 1,750 to 3,500 was formally accepted by the CTC in 2005.




Commentary: The King Salmon River is a small non-glacial system that supports a small run of inside-rearing Chinook salmon. Escapements are based upon weir counts from 1983 to 1992 and expansions of index counts from 1971 to 1982 and 1993 to 2005. The 10 years of weir data showed that on average the escapement was 1.5 times the index count (McPherson and Clark 2001).




Commentary: Andrew Creek, a tributary of the lower Stikine River, is a small non-glacial system that supports a moderate run of inside-rearing Chinook salmon. Escapements are based upon weir counts from 1976 to 1984 and expansions of index counts in 1975 and 1985 to 2005. Four years of concurrent weir and index count data were used to estimate the expansion factor of 2.0.




Commentary: The Unuk River is a moderate-sized glacial system that supports a moderate run of inside-rearing Chinook salmon. Indices of escapement since 1977 are based on the sum of peak index counts from six main tributaries (Pahlke 2003). Mark-recapture studies were implemented in 1994 and annually since 1997 (Weller and McPherson 2003). The current estimated expansion factor is 5.0 for index counts.



Commentary: The Chickamin River is a moderate-sized glacial system that supports a moderate run of inside-rearing Chinook salmon. Indices of escapement since 1975 are based on the sum of peak index counts from eight main tributaries (Pahlke 2003). Mark-recapture studies were performed in 1995, 1996, and 2001-2005. The current estimated expansion factor is 4.64 for index counts.




Commentary: The Blossom River is a small-sized non-glacial system that supports a small run of inside-rearing Chinook salmon. Indices of escapement since 1975 are based on the sum of peak index counts (Pahlke 2003). Mark-recapture studies were performed in 1998, 2004, and 2005 indicating an estimated expansion factor range of 2.0 to 4.0.




Commentary: The Keta River is a small-sized non-glacial system that supports a small run of inside-rearing Chinook salmon. Indices of escapement since 1975 are based on the sum of peak index counts (Pahlke 2003). Mark-recapture studies were performed 1998 to 2000 (Freeman et al. 2001). The current estimated expansion factor is 3.0 for index counts.

2.3.2 Canadian Stocks

Since the beginning of the Chinook rebuilding program of the 1985 PST, escapement goals for Canadian Chinook stocks were generally based on doubling the average escapements recorded between 1979-1982. The doubling was based on the premise that Canadian Chinook stocks were over-fished and that doubling the escapement would still be less than the optimal escapement estimated for the aggregate of all Canadian Chinook populations (see stock-recruitment curve in “Technical Basis of PSC Catch Ceilings,” Figure 1, Attachment 4, PSC file 72006; PSC Office, Vancouver, BC). Doubling was also expected to be a large enough change in escapements to allow detection of the change in numbers of spawners and the subsequent production. The escapement goals of the Canadian stocks are currently being reviewed.




Commentary: The Yakoun River is the only significant Chinook-producing stream on the Queen Charlotte Islands. Chinook spawn primarily at the outlet of Yakoun Lake and are a summer-run stock. Visual estimates of escapement are made by foot surveys of the system. These estimates are then expanded into a total estimate of spawning escapement in the system. The effort spent on escapement surveys has declined in recent years and their accuracy (i.e. ability to estimate the actual escapement) is unknown.




Commentary: The Nass River is the largest river in Area 3, representing a group of approximately 25 streams. Prior to 1992, CDFO observations of escapement were based on visual counts. Mark-recapture programs have been conducted since 1992 by the Nisga'a Fisheries to estimate total spawning escapement in the Nass River. The Nass mark-recapture program uses two fish wheels at Gitwinksihlkw (GW) in the lower Nass canyon to apply tags and two wheels at Grease Harbour in the upper canyon for recovery. A modified Petersen model, stratified by size category, was used to estimate the total population of Chinook past the tagging location. Tags were also recovered in upriver fisheries and on the spawning grounds. Spawning


escapements were calculated as the estimated Chinook population past GW from the mark-recapture studies, less upriver catches in sport and First Nations fisheries. Three tributaries with Chinook populations enter the Nass River below GW. Visual estimates augmented by fence counts of the Kincolith River in 2001, 2002 and 2005 were used to estimate Nass River Chinook escapements below the fish wheels.



Commentary: The Skeena Chinook escapements above represent 40 streams which are consistently surveyed. The Skeena supports over 75 separate Chinook spawning populations, but three (Kitsumkalum, Morice, and Bear Rivers) account for about 70% of the total abundance. A second group of populations (Ecstall, Kispiox, and Babine rivers) have annual returns ranging from 1,000 to 5,000 spawners, and comprise about 13% of Skeena returns. Escapement estimates are generally based on visual observations from helicopter, fixed wing aircraft and/or from stream walking surveys. The Kitsumkalum River is the exploitation rate indicator stock for the Skeena Chinook complex. Spawning escapements in the Kitsumkalum have been estimated using a mark-recapture program since 1984.






Commentary: The Area 8 Chinook stock consists of seven non-enhanced systems, but the Dean River is the main spawning population. Of all Chinook- producing streams in the Central Coast, the Dean is the best indicator in terms of consistent survey coverage and methodology. Chinook returning to the Dean River have an early-summer timing and most spawn in the lower river by July. Up until 2000, counts of spawning Chinook were made during 1-3 surveys and the peak count used as the escapement index. Survey counts were sometimes expanded to account for sections of the river that could not be surveyed in any year, but the counts were not extrapolated to total escapement of Chinook to the river. Since 2001, the annual number of aerial surveys has increased, allowing the calculation of area-under-the-curve (AUC) escapement estimates. Poor counting conditions were encountered during (Aug.) 2004 and as a result an AUC estimate was not possible. Instead a maximum likelihood estimate was used and yielded a total Chinook return of 3,500 to the Dean River.





Commentary: The Wannock, Chuckwalla, and Kilbella Rivers are the primary Chinook streams in Area 9 (Rivers Inlet area). Small tributaries of Owikeno Lake also contain Chinook but these

populations are much smaller. The Wannock River contains the largest Chinook population, averaging 5,200 Chinook in the 1990s, while the Chuckwalla and Kilbella together averaged around 300. The Wannock River drains Owikeno Lake, is about six kilometers long, and is wide and turbid. The Chuckwalla and Kilbella rivers are much longer, drain from coastal mountains, and their visibility is much more variable depending on local weather (glacial flour to clear). The timing of these stocks also differs: the Wannock has late summer/fall run timing, the other two are early summer Chinook stocks. Escapement estimates in the Chuckwalla and Kilbella rivers are derived from aerial surveys, whereas Wannock escapement is derived from expansions of carcass count to estimate total spawning escapement.






Commentary: The WCVI index represents the sum of escapements for six rivers (Marble, Tahsis, Burman, Artlish, Kaouk, and Tahsish), which were chosen to provide an ‘index’ of escapement for wild WCVI stocks in general. These stocks were chosen based on historical consistency of data quality. CDFO has developed a 14 stream expanded index which includes escapements to the six stream index plus the following WCVI streams: Colonial/Cayegle Creeks (Area 26), Leiner (Area 25), Megin, Bedwell/Ursus, Moyeha (Area 24) and Sarita, Nahmint (Area 23), and San Juan (Area 21). In 2005, the Colonial/Cayegle escapement was not available, and was therefore not included in the 14 stream index.



Commentary: The Upper Georgia Strait (UGS) stock index consists of four river systems (Klinaklini, Kakweiken, Wakeman, Kingcome) in Johnstone Strait mainland inlets and the Nimpkish River on northeast Vancouver Island. The accuracy of escapement estimates in the mainland inlet systems is likely poor due to their glacial nature and remote access. Escapement estimates have primarily been based on aerial counts. Swim surveys and stream walks have been conducted in the Nimpkish River. A fish wheel program implemented on the Klinaklini in 1997 was discontinued in 2004. Based on the portion of the assessment program that continued in 2005, estimated abundance in 2005 was assumed to be the same as in 2004.




Commentary: Lower Georgia Strait (LGS) rivers monitored for naturally spawning fall Chinook escapement are the Cowichan and Nanaimo rivers. Total Chinook returns to the Cowichan and Nanaimo rivers have been estimated since 1975. Prior to 1989, escapement estimates from the Cowichan River were derived from swim surveys and over-flights by Fishery Officers and hatchery staff. This approach was also used for the Nanaimo River prior to 1995. Since 1989 a counting fence has been used in the Cowichan, and since 1995 carcass mark-recapture surveys have been used in the Nanaimo. An escapement goal of 6,500 for the Cowichan was accepted by the CTC in 2005; a goal for the Nanaimo is still pending.

2.3.3 Fraser River Stocks


The Fraser River watershed is the largest Canadian producer of Chinook salmon. Fraser Chinook are comprised of a large number of local populations as described in CTC (2002b).

Much of our understanding of the status of Fraser Chinook is based on spawner escapement data. Most data are from visual surveys, which are generally biased low, although many estimates are considered to be reasonably precise. Visual survey data are generated from aerial over-flight surveys and the escapement estimate is usually obtained by dividing the peak count by 0.65 (Farwell et al. 1999). The CDFO continues to evaluate the appropriateness of this expansion factor and AUC methodology through calibration studies. Counting fences and mark-recapture projects exist for some systems, although most of the time series of escapement data from these projects are relatively short.

For populations other than the Harrison River, habitat-based models are being developed to estimate spawning capacity and spawner abundance producing maximum sustained yield. This habitat-based assessment will initially focus on predictive models based on Chinook stock-recruitment relationships, although other habitat-based approaches will also be considered.




Commentary: This aggregate includes the Upper Pitt River and Birkenhead River stocks in the Lower Fraser, and the spring-run Chinook of the Mid and Upper Fraser, North Thompson, and South Thompson, but excluding those of the Lower Thompson (CTC 2002b). Stocks upstream of Prince George include the McGregor and Torpy River systems. In recent years, fence counts have been employed at the Chilako River in the Upper Fraser and at the Salmon River in Salmon Arm (South Thompson). Fence counts were discontinued at the Salmon River (Prince George) in 1998. Estimates for all other systems were generated from aerial surveys, typically, by dividing the peak count by 0.65.




Commentary: The Fraser Spring-Run Age 1.2 aggregate includes six smaller body size populations that spawn in the Lower Thompson River tributaries, Louis Creek of the North Thompson and the spring-run fish of Bessette Creek in the South Thompson (CTC 2002b). Escapement estimates for each system are generated from visual surveys, either from aerial over-

flights, stream walks or by dividing the peak counts by 0.65. The Nicola watershed is a site for calibrating peak count expansion, AUC, and mark-recapture methods.




Commentary: The Fraser Summer-Run Age 1.3 stock complex includes 11 populations, spawning in large rivers, mostly below the outlets of large lakes. These include the Stuart and Nechako rivers upstream of Prince George, Chilko and Quesnel rivers in the mid Fraser and the Clearwater and North Thompson rivers in the North Thompson watershed (CTC 2002b). Escapement estimates are generated from aerial surveys by dividing the peak count by 0.65, except for the Stuart system where a mark-recapture estimate is generated, and for the Nechako River where multiple aerial counts are analyzed with the AUC method.




Commentary: The Fraser Summer-Run Age 0.3 aggregate includes six populations of Chinook spawning in the South Thompson watershed upstream of Kamloops and one in the lower Fraser. These include the Middle Shuswap, Lower Shuswap, Lower Adams, Little River and the South Thompson River mainstem, in the BC interior, and Maria Slough in the lower Fraser (CTC

2002b). Most escapements are estimated by expanding peak visual survey counts (as in previous three Fraser aggregates). Further, the lower Shuswap River is a site for calibrating peak count expansion, AUC, and mark-recapture methods.



Commentary: The lower Fraser stock is dominated by fall returning Harrison-origin Chinook that includes natural spawners in the Harrison River and Harrison-origin fish that were introduced to the Chilliwack River. Since 1984, mark-recapture studies have been conducted annually to obtain reliable estimates of spawning escapements. Estimates of fall Chinook escapement to the Chilliwack River are based on a procedure long established by the Chilliwack Hatchery staff for expanding the number of carcasses counted in standardized reaches of the river.


Harrison River Fall Chinook Escapement and Total Fraser Late Terminal Run
(Harrison R + Chilliwack R Escapements + Terminal Catch)




2.3.4 Washington, Oregon and Columbia River Stocks

The PSC escapement indicator stocks in Washington, Oregon, and Idaho are separated into five groups: Puget Sound, Washington Coastal, Columbia River, North Oregon Coastal, and Mid Oregon Coastal. The indicator stocks include a variety of run timings and ocean distributions.

Biologically based escapement goals have been reviewed and accepted by the CTC for three fall stocks (Queets, Quillayute, Hoh), two Spring/summer stocks (Queets, Hoh), three Columbia River stocks (Lewis, Upriver Brights and Columbia River summer), and three Oregon coastal stocks (Nehalem, Siletz and Siuslaw).




Commentary: In 2005, the escapement estimate was 2,047 for the North Fork and 130 for the South Fork. However, only 10% of the North Fork escapement is identified as natural-origin spawners, and the bulk of the run is composed of hatchery-origin returns from the supplementation program. The Comprehensive Chinook Management Plan (2004) conservation objective for 2005 was for an Adult Equivalent (AEQ) exploitation rate across all southern U.S. fisheries not to exceed 7%. The state-tribal escapement goal established for this stock is 4,000 spawners. There is a small ceremonial and subsistence directed fishery on the spring Chinook and substantial incidental impacts during the terminal fall Chinook fisheries.




Commentary: Due to changes in spawning index areas, beginning in 1992 for the Cascade stock and 1994 for the Sauk and Suiattle stocks, escapements are not directly comparable to previous numbers. The past state-tribal escapement goal of 3,000 adults was the average of the estimated escapements from 1959-1968 (PFMC 1997). In 2004 and in 2005 the Recovery Exploitation Rate (RER) for Skagit springs was 38%, with 576 spawners as the low abundance threshold. While no postseason estimate is available, the preseason expectation for 2005 was for a total rate


of 29.1% (PFMC 2005). Proposed escapement goals, as stated in the draft Shared Strategy Recovery Plan, are 1,200 Chinook for low marine survival years and 2,100 Chinook for high marine survival years. The 2005 escapement estimate is 1,305 natural spawners.




Commentary: Projects to improve escapement estimates of Skagit summer/fall Chinook have recently been funded through the Letter of Agreement (LOA) process. They included: development of variance estimates, determination of age and sex composition of the escapement, and evaluation of the 21-day redd life assumption and 2.5 fish/redd expansion value. The state-tribal escapement goal for this stock is 14,850, the average of the 1965-1976 escapements (Ames and Phinney 1977). Little terminal harvest has occurred since 1997. In 2005, the Federal Management Plan (FMP) conservation objective for this stock was for a RER across all fisheries not to exceed 50%. The 2005 predicted exploitation rate was 39.6%. The 2005 escapement estimate is 20,803 and the terminal run estimate is 23,396.




Commentary: Natural spawning broodstock are collected annually in the river to maintain a CWT indicator stock program and to augment natural production. From 1989 to 1996, approximately 18% of the escapement was comprised of returns from this program. (1996 to 2005 average is 38% hatchery origin returns) The state-tribal escapement goal of 2,000 fish is the average of the 1973-1976 escapements (Ames and Phinney 1977). There have been no terminal harvests since 1996. The 2005 FMP conservation objective for the combined summer/fall stock was for an AEQ exploitation rate not to exceed 15% in the southern U.S. fisheries. The preseason estimate of the total AEQ exploitation rate was 12.1%. The escapement estimate for 2005 is 963 Chinook (885 for the North Fork and 78 for the South Fork).




Commentary: Some terminal area harvest of Snohomish River Chinook occurs in Area 8 incidental to net and sport fisheries targeting Tulalip Hatchery Chinook salmon. Historic terminal run size and catch estimates derived from run reconstruction are being revised to reflect the results of otolith marking studies. The state-tribal escapement goal for this stock had been 5,250 fish (the average of the 1965-1976 escapements). The FMP conservation objective was for a total AEQ exploitation rate not to exceed 15% in southern U.S. fisheries. The preseason prediction of that rate was 14.9%. The 2005 escapement was estimated at 4,484 natural spawners.




Commentary: Substantial artificial production occurs in Issaquah Creek and at the University of Washington. In 1994, spawning estimates were reviewed, and an attempt was made to find a consistent method to estimate escapement. A state-tribal escapement goal of 1,200 has been established for the Cedar River spawners. The single targeted goal represents an index count for the Cedar River. This objective reflects the average of observed spawning escapements from 1965-1969. It should be noted that although there are no hatchery fish released from the Cedar River, nearly 40% of the spawning fish were of hatchery origin. The FMP conservation objective for 2005 for Lake Washington Fall Chinook was for an AEQ exploitation rate not to exceed 15% in all preterminal southern U.S. fisheries. The preseason expected AEQ exploitation rate was 9.8%. The 2005 escapement was a total of 726 spawners (511 to Cedar and 215 to the north tributaries). Hatchery-origin returns (HOR) into the Cedar was estimated to be about 29%, with the north tributary HOR component about 79% for 2005. There have not been freshwater terminal fisheries on this stock since 1995.




Commentary: There is a large hatchery program in this basin and these fish comprise a large portion of the return. The average is about 52% for the years 1996-2003. Tagging studies were conducted in 1975 and 1976 to estimate numbers of returning adults; results were in close agreement with estimates made from aerial surveys. No attempt is made to adjust the estimate of natural escapement for the presence of hatchery origin fish. Projects to improve escapement estimates of Green River fall Chinook, were recently funded through the LOA process, including evaluation of the spatial and temporal distribution of escapement, alternative methods of estimating escapement, and the validity of the 21-day redd life assumption and 2.5 fish/redd expansion value. The state-tribal escapement goal of 5,750 naturally spawning adults is the average of the 1965-1976 escapements (Ames and Phinney 1977). Beginning in 2003, a new method for estimating natural spawning escapement was employed based on a mark/recapture studies conducted 2000-2002. The estimate of mainstem females was compared to the “adjusted” peak count of visible redds for that year, with the assumption that each female dug a single redd. In 2003, the mean ratio of mainstem females to mainstem adjusted peak redds (3.109) from the three study years was applied to the 2005 adjusted peak redd count to estimate mainstem female spawners. A sex ratio of 1.5 males per female was then used to expand the number of female spawners to total mainstem escapement. The 2005 FMP conservation objectives for this stock was for a total AEQ exploitation rate not to exceed 15% in preterminal southern U.S. fisheries, and an escapement of at least 5,800 adults. The 2005 escapement estimate for natural spawning Chinook was 4,089. The number of hatchery-origin spawners was estimated to be almost 60%.



Commentary: There are no directed fisheries on Chinook returning to rivers entering the Strait of Juan de Fuca. The escapement goal established by state and tribal managers is 850 naturally spawning adults. This single targeted goal was developed as a MSY proxy. The escapement goal was calculated by estimating the amount of available spawning habitat, then expanded utilizing assumed optimal redds per mile and fish per redd values (Ames and Phinney 1977). The escapement and terminal run size estimates for 2003 are 1,100 adults. The 2004 escapement estimate was 1,088, while the 2005 escapement estimate dropped to 283.




Commentary: A summer Chinook hatchery program using native stock operated from the mid-1970s to the mid-1980s. Spring Chinook of non-native origin were introduced in a hatchery program in the early 1970s. CWT analyses since then have demonstrated significant straying of these spring Chinook into the summer Chinook spawning population. Estimates from 1991-1995 averaged 47% hatchery origin strays in the naturally spawning population. In 1996, fry plants were eliminated and the smolt plants were reduced. Summer Chinook are managed for a fixed escapement goal of 1,200 adults and jacks combined (PFMC 2003). The 2005 escapement estimate for summer Chinook is 876. This continues a trend of stable returns near the management goal for this stock.




Commentary: No hatchery production of fall Chinook currently occurs in the Quillayute River basin; the program was discontinued in the late 1980s. Since 1991, the returning run size has fluctuated within a range comparable to run sizes observed prior to 1984. The 2005 escapement estimate is 6,406 with a total terminal estimate of 8,402. Terminal fisheries are managed for a harvest rate of 40%, with an escapement floor of 3,000 fish (PFMC 2003). This objective is

designed to actively probe at and above estimates of escapements that produce maximum sustained harvest (MSH), while minimizing potential detrimental effects of existing fisheries. Stock production analyses of spawning escapements from 1968-1982 were used to determine the initial escapement floor.




Commentary: Similar to many of the other Washington coastal stocks, Hoh River spring/summer escapements have been relatively stable except for much larger returns in 1988, 1989, and 1990. The terminal return for this stock declined from 1997 to 2000, but has since rebounded. Terminal fisheries are managed to harvest 31% of the river run, with an escapement floor of 900 fish (PFMC 2003). This objective is designed to allow a wide range of spawner escapements from which to eventually develop an MSY objective or proxy while protecting the long-term productivity of the stock. Stock production analysis of spawning escapement for brood years 1969-1976 was utilized to determine the initial escapement floor. The 2005 escapement estimate and total run size is 1,193 and 1,389 respectively.




Commentary: The natural escapement estimates include fish taken for broodstock in the 1980s.


This stock is managed to harvest 40% of the terminal run, with an escapement floor of 1,200 spawners (PFMC 2003). This objective is designed to actively probe at and above estimates of the escapements that produce MSH, while minimizing potential detrimental effects of existing fisheries. Stock production analyses of spawning escapements from 1968-1982 were utilized to determine the initial escapement floor. The 2005 escapement estimate is 4,180 and a terminal run size of 5,267.




Commentary: Terminal fisheries are managed to harvest 30% of the river run size, with an escapement floor of 700 fish (PFMC 2003). This objective is designed to actively probe at and above the estimates of escapement that produce MSH. Since 1990, terminal fisheries have had minimal impact on this stock as returns to the river have rarely exceeded the escapement floor in this time frame. Since 2000, sport anglers have been required to release all Chinook during the summer, and tribal fisheries have been limited to one tribal netting day for ceremonial and subsistence purposes. Stock production analysis of spawning escapement for brood years 1969-1976 were used to determine the initial escapement floor. The 2005 escapement estimate is 294, with a terminal run size of 302.




Commentary: The 2005 escapement and total run size is 2,931 and 4,253, respectively. Terminal fisheries are managed to harvest 40% of the river return, with an escapement floor of 2,500 spawners (PFMC 2003). This objective is designed to actively probe at and above estimates of the escapements that produce MSH. Stock production analyses of spawning escapements from 1967-1982 were used to determine the initial escapement floor.



Commentary: The Grays Harbor spring Chinook stock is managed for a fixed natural spawning escapement goal of 1,400 fish (PFMC 2003). This single targeted goal was developed as a MSY proxy. This objective was derived from actual spawning data from the mid- to late 1970s, expanded to include additional habitat not covered by spawner surveys. The 2005 escapement was 2,129 Chinook and the 2005 terminal run was 2,743 Chinook.




Commentary: Grays Harbor fall Chinook are managed for a maximum sustained production escapement goal of 14,600 spawners for the Chehalis and Humptulips systems combined (PFMC 2003). This single targeted goal was developed as an MSY proxy. The objective represents assumed optimal spawner density based on estimated available habitat. The 2005 escapement was 19,249 Chinook and the terminal run was 21,410 Chinook salmon.




Commentary: In 1992, Snake River spring/summer naturally spawning Chinook were listed under the ESA. In past escapement assessments, the CTC used the goal of 84,000 natural spawners passing Bonneville Dam (an estimated 70% wild portion of the 120,000 specified in the original 5-year plan for U.S. v Oregon). The interim management goal for the Columbia River Fish Management Plan (CRFMP 1988) for Columbia River Springs was 115,000 hatchery and wild adult Chinook counted at Bonneville Dam and 25,000 naturally produced plus 10,000 hatchery produced adults counted at Lower Granite Dam. However, the CRFMP is currently being renegotiated. The 2005 escapement was 35,833 natural spawners. Terminal harvests were severely constrained from 1977 until recently, with incidental harvests in lower river fisheries

averaging 2% and total harvest in treaty Indian fisheries averaging 5.5% (TAC 1999). On the recent large returns, the terminal harvest rates have been between 13.5% and 19.0%.




Commentary: Productivity is limited by loss of downstream migrants, habitat degradation, lack of screens on water diversions, high water temperatures, low flows, and sediment-laden irrigation water returns (CBFWA 1990). The 2005 escapement was 39,138 naturally spawning fish. Directed commercial fisheries for upper Columbia River summer Chinook resumed in 2003 above Bonneville Dam and in 2004 below Bonneville Dam when the Columbia Upriver Summers began to exceed the interim management goal of 29,000 hatchery and natural origin adults as measured at the Columbia River mouth. The non-Indian and tribal harvest rates between 2003 and 2005 averaged 3.9% and 10.2%, respectively.




Commentary: The escapement goal is 40,000 naturally spawning fish. The 2002, 2003, and 2004 escapements past McNary dam of 141,682, 179,970, and 168,679 were the largest since the


peak escapement and terminal run in 1987. The 2005 escapement of 134,821 was greater than the 1987-2004 average of 93,350.




Commentary: The escapement goal for the Lewis River is 5,700 naturally spawning fish. Except in 1999, escapements have been above the goal since 1979. The 2002, 2003, and 2004 returns and escapements of Lewis River fall Chinook were the largest since 1990. The estimated escapement in 2005 was 16,767 Chinook.




Commentary: Local management agencies use a goal of 4,000 adult Chinook, which includes 2,000 fish above Sherars Falls. This goal is based on average spawning escapement. The 2002 and 2003 escapements of Deschutes fall Chinook were at least 3 times the management goal, based on either the expansion of escapements above Sherars Falls, or the total river mark recapture estimate. They were also the largest escapements since the peak in 1997. The estimated escapement in 2005 was 13,550 Chinook.




Commentary: Estimated spawner abundance was 7,038 large (adult) Chinook for 2005. Methods used to generate escapement estimates in this basin have not changed since last report in 2005. Punch card data used to estimate the recreational sport catch are unavailable for 2005, hence terminal run sizes are not available for this year.




Commentary: The Siletz River spawner abundance in 2005 is estimated at 6,426 adult Chinook salmon. Methods used to generate escapement estimates in this basin have not changed since last report in 2005. All four standard surveys were conducted in 2005. Punch card data used to estimate the recreational sport catch are unavailable for 2005, hence terminal run sizes are not available for this year.



Commentary: The estimated spawner abundance in 2005 was 16,619 adult Chinook. Methods used to generate escapement estimates in this basin have not changed since 2004. Punch card data used to estimate the recreational sport catch are unavailable for 2005, hence terminal run sizes are not available for this year.



Commentary: Fall run Chinook from the Umpqua River are harvested in PSC fisheries, and should be included as escapement indicator stocks. Four years of LOA funded research has allowed the calibration of the redd counts to derive a fish per redd expansion factor so that annual escapements estimates can be made. The average expansion factor from these studies is 3.69 fish per redd. The coefficient of variation of the expansion factor was found to be 14%, which shows that the average expansion factor is a reliable statistic to use for annual estimates of escapement. The escapement estimate for 2005 was 3,084 based on redd count expansions.



Commentary: The 2005 MOC density index is calculated to be 65 fish/mile. Research funded by the LOA is underway that will provide information to designate the Coquille Chinook production river system as the escapement indicator stock for this stock aggregate.

3 EXPLOITATION RATE ANALYSIS AND MODEL CALIBRATION

3.1 INTRODUCTION

This chapter describes the methods and results of the cohort analysis, used to estimate exploitation rates from CWT data, and the PSC Chinook model calibration. The results of the 2006 preseason calibration (CLB 0604) are based on the exploitation rate analysis using CWT data through 2004, coast-wide data on catch, spawning escapements and age structure through 2005, and forecasts of Chinook returns expected in 2006. This chapter includes:

- 1) estimated postseason abundance indices for 1979 through 2005 and the preseason projection for 2006 for the AABM fisheries,
- 2) estimated non-ceiling indices, referred to as the ISBM indices in this report, for 1999 to 2004 and modeled ISBM projections for the 2006 ISBM fisheries,
- 3) estimated stock composition for 1979 through 2005 and a projection for 2006 for the AABM and other fisheries, and
- 4) estimated harvest rates (fishery indices) for the AABM fisheries.

Appendices C to H summarize the indicator stocks, ISBM indices, distribution of catch and total mortality, abundance indices for the AABM fisheries from CLB 0604, stock composition of AABM fisheries estimated from the PSC Chinook model and incidental mortality rates used.

This chapter is an abbreviated version of TCCHINOOK (05)-2 (CTC 2005b); see that document for a complete description of methods and prior year's results that are not included here.

3.2 METHODS

The exploitation rate assessment is performed through cohort analysis of CWT release and recovery data (CTC 1988). Cohort analysis is the reconstruction of the exploitation history of a given stock and brood year and is used to produce a variety of statistics, including total exploitation rates, age and fishery specific exploitation rates, maturation rates, pre-age 2 recruitment survival indices, and annual distribution of fishery-related mortalities.

Estimates of age and fishery-specific exploitation and maturation rates from the cohort analysis are combined with data on catches, escapements, non-retention, and enhancement to complete the annual calibration of the CTC Model. The calibration procedure estimates pre-age 2 survival to recruitment for the stocks included in the model.

Results from the annual preseason calibration of the Chinook model are used to calculate: 1) AIs for the three AABM fisheries; 2) postseason AIs for the previous year; and 3) preseason and postseason ISBM indices. Projected AIs for 2006 are used to determine the allowable 2006 catch of Treaty Chinook for AABM fisheries. Postseason AIs are used to determine postseason allowable catches and to evaluate compliance for AABM fisheries. For the ISBM fisheries, the Agreement specifies that Canada and the United States will reduce the exploitation rate from the 1979–1982 base period by 36.5% and 40.0%, respectively, on stocks that have not achieved their CTC agreed escapement goals. The ISBM index is used to estimate the annual reduction in

exploitation rates relative to the base period. Postseason ISBM indices for 2004 are computed using results of the exploitation rate analysis. Forecasts of the 2006 ISBM indices are computed using the CTC model. The Agreement specifies that the postseason ISBM indices estimated through exploitation rate analysis of CWT recoveries will be used to assess the ISBM index.

3.3 EXPLOITATION RATE ASSESSMENT (THROUGH CALENDAR YEAR 2004)

The CTC currently monitors 39 exploitation rate indicator stocks that are coded-wire tagged, but only 36 were used for analyses in this chapter (Table 3.1). An exploitation rate indicator stock is not used in the exploitation rate analysis if the number of CWT recoveries is very limited¹ or there is no quantitative estimate of tags in the spawning escapement (see footnotes in Table 3.2). Indicator stocks used for exploitation rate analysis and the type of analysis performed for each are shown in Table 3.2. The relationship between the exploitation rate indicator stocks, model stocks, and PST Annex stocks are shown in Appendix C. Extrapolation of results to similar stocks and/or generalizations about fishery impacts will only be appropriate to the extent that the exploitation rate indicator stocks are representative of the stocks groups they are intended to represent.

¹ 35 estimated recoveries for a given stock and age combination.

Table 3.1. The 39 exploitation rate indicator stocks monitored by the CTC, their location, run type, and smolt age. Stocks in bold, italic text were not used in the exploitation rate analysis.

| Area | Exploitation Rate Indicator Stocks | Location | Run Type | Smolt Age |
|-----------------------------------|---|---------------------------|----------------------|--------------|
| S.E. Alaska | Alaska Spring | Southeast Alaska | Spring | Age 1 |
| British Columbia | Kitsumkalum | North/Central BC | Summer | Age 1 |
| | <i>Atnarko</i> ¹ | <i>North/Central BC</i> | <i>Spring/Summer</i> | <i>Age 0</i> |
| | <i>Kitimat River</i> ¹ | <i>North/Central BC</i> | <i>Summer</i> | <i>Age 0</i> |
| | Robertson Creek | WCVI | Fall | Age 0 |
| | Quinsam | Georgia Strait | Fall | Age 0 |
| | Puntledge | Georgia Strait | Summer | Age 0 |
| | Big Qualicum | Georgia Strait | Fall | Age 0 |
| | Cowichan | Georgia Strait | Fall | Age 0 |
| | <i>Chehalis (Harrison Stock)</i> ¹ | <i>Lower Fraser River</i> | <i>Fall</i> | <i>Age 0</i> |
| | Chilliwack (Harrison Stock) | Lower Fraser River | Fall | Age 0 |
| Puget Sound | Nooksack Spring Fingerling | North Puget Sound | Spring | Age 0 |
| | Nooksack Spring Yearling | North Puget Sound | Spring | Age 1 |
| | Skagit Spring Fingerling | Central Puget Sound | Spring | Age 0 |
| | Skagit Spring Yearling | Central Puget Sound | Spring | Age 1 |
| | Samish Fall Fingerling | North Puget Sound | Summer/Fall | Age 0 |
| | Skagit Summer Fingerling | Central Puget Sound | Summer | Age 0 |
| | Stillaguamish Summer Fingerling | Central Puget Sound | Summer/Fall | Age 0 |
| | Nisqually Fall Fingerling | Central Puget Sound | Summer/Fall | Age 0 |
| | University of Washington Accelerated | Central Puget Sound | Summer/Fall | Age 0 |
| | George Adams Fall Fingerling | Hood Canal | Summer/Fall | Age 0 |
| | South Puget Sound Fall Fingerling | South Puget Sound | Summer/Fall | Age 0 |
| | South Puget Sound Fall Yearling | South Puget Sound | Summer/Fall | Age 1 |
| | Squaxin Pens Fall Yearling | South Puget Sound | Summer/Fall | Age 1 |
| | White River Spring Yearling | South Puget Sound | Spring | Age 1 |
| Washington Coast /Juan de Fuca | Elwha Fall Fingerling | Strait of Juan de Fuca | Summer/Fall | Age 0 |
| | Hoko Fall Fingerling | Strait of Juan de Fuca | Summer/Fall | Age 0 |
| | Sooes Fall Fingerling | North Wash. Coast | Fall | Age 0 |
| | Queets Fall Fingerling | North Wash. Coast | Fall | Age 0 |
| Columbia River | Willamette Spring | Lower Columbia R. | Spring | Age 1 |
| | Columbia Summers | Columbia R. (WA) | Summer | Age 1 |
| | Cowlitz Tule | Columbia R. (WA) | Fall Tule | Age 0 |
| | Spring Creek Tule | Columbia R. (WA) | Fall Tule | Age 0 |
| | Columbia Lower River Hatchery | Columbia River (OR) | Fall Tule | Age 0 |
| | Columbia Upriver Bright | Upper Columbia R. | Fall Bright | Age 0 |
| | Hanford Wild | Upper Columbia R. | Fall Bright | Age 0 |
| | Lyons Ferry ² | Snake River | Fall Bright | Age 0 |
| | Lewis River Wild | Lower Columbia R. | Fall Bright | Age 0 |
| Oregon Coast | Salmon River | North Oregon Coast | Fall | Age 0 |

¹ These stocks are CWT-tagged, but there is no reliable quantitative CWT escapement data and CWT data presented for these stocks is useful for distribution of harvest and mortalities only.

² Subyearlings have been CWT-tagged since brood year 1986, except for brood years 1993 through 1997.

Table 3.2. The 36 CWT exploitation rate indicator stocks used in the exploitation rate analysis and the data derived from them: fishery, ISBM and survival indices, brood exploitation rates (Brood Exp), and stock catch distribution (Dist) with quantitative escapement estimates (Esc) and tagging during the base period years 1979–1982.

| Exploitation Rate Indicator Stocks | Fishery Index | ISBM Index | Brood ¹ Exp | Survival Index | | | Base Tagging |
|--------------------------------------|---------------|--------------|------------------------|----------------|------|------------------|--------------|
| | | | | | Dist | Esc | |
| Alaska Spring | yes | — | Total | yes | yes | yes | yes |
| Kitsumkalum | — | — | Total | yes | yes | yes | — |
| Robertson Creek | yes | yes | Ocean ¹ | yes | yes | yes | yes |
| Quinsam | yes | yes | Total | yes | yes | yes | yes |
| Puntledge | yes | — | Total | yes | yes | yes | yes |
| Big Qualicum | yes | yes | Total | yes | yes | yes | yes |
| Cowichan | yes | yes | Total | yes | yes | yes | — |
| Chilliwack (Harrison Fall Stock) | — | yes | Total | yes | yes | yes | — |
| Nooksack Spring Fingerling | — | — | ⁴ | — | yes | yes | — |
| Nooksack Spring Yearling | — | yes | ⁴ | yes | yes | yes ³ | — |
| Skagit Spring Fingerling | — | — | Ocean | — | yes | yes | — |
| Skagit Spring Yearling | — | — | Ocean | yes | yes | yes ³ | — |
| Samish Fall Fingerling | yes | — | Ocean | yes | yes | yes ³ | yes |
| Skagit Summer Fingerling | — | — | Ocean | — | yes | yes | — |
| Stillaguamish Summer Fingerling | — | yes | ⁴ | — | yes | — | — |
| Nisqually Fall Fingerling | — | — | ⁴ | — | yes | — | yes |
| University of Washington Accelerated | yes | ² | ² | — | yes | yes ³ | yes |
| George Adams Fall Fingerling | yes | ² | ² | yes | yes | yes ³ | yes |
| South Puget Sound Fall Fingerling | yes | yes | Ocean | yes | yes | yes ³ | yes |
| South Puget Sound Fall Yearling | yes | ² | ² | yes | yes | yes ³ | yes |
| Squaxin Pens Fall Yearling | — | ² | ² | yes | yes | yes ³ | — |
| White River Spring Yearling | — | — | ⁴ | yes | yes | yes ³ | yes |
| Elwha Fall Fingerling | — | — | ⁴ | yes | yes | — | — |
| Hoko Fall Fingerling | — | — | Ocean | yes | yes | yes | — |
| Sooes Fall Fingerling | — | — | Ocean | yes | yes | yes | — |
| Queets Fall Fingerling | — | yes | ⁴ | yes | yes | — | yes |
| Willamette Spring | yes | — | Ocean | yes | yes | yes | yes |
| Columbia Summers | yes | yes | Total | yes | yes | yes | — |
| Cowlitz Tule | yes | — | Ocean | yes | yes | yes | yes |
| Spring Creek Tule | yes | — | ² | yes | yes | yes | — |
| Columbia Lower River Hatchery | yes | — | ² | yes | yes | yes | yes |
| Upriver Bright | yes | yes | Total | yes | yes | yes | yes |
| Hanford Wild | — | — | Total | yes | yes | yes | — |
| Lyons Ferry | — | — | Total | yes | yes | yes | — |
| Lewis River Wild | yes | yes | Total | yes | yes | yes | yes |
| Salmon River | yes | yes | Ocean | yes | yes | yes | yes |

¹ For stocks of hatchery origin and subject to terminal fisheries directed at harvesting surplus hatchery production, ocean fisheries do not include terminal net fisheries. Otherwise, total fishery includes terminal net fisheries.

² Hatchery stock not used to represent naturally spawning stock.

³ Only hatchery rack recoveries are included in escapement.

⁴ Insufficient escapement data for exploitation rate analysis

3.4 MODEL OUTPUT

3.4.1 AABM Abundance Indices and Associated Catches

Beginning with the 1999 fishing season, the Agreement specified that the AABM fisheries are to be managed through the use of the preseason AIs, where specific allowable harvest corresponds to a given AI for each fishery. The preseason AIs that were used to establish harvest management targets are listed in Table 3.3. The 2006 preseason AI for the SEAK troll fishery is 1.69, for the NBC troll fishery it is 1.53, and for the WCVI troll fishery is 0.75.

The postseason AI is considered a more accurate estimate of the abundance index for the AABM fisheries, and is used to compute a final allowable catch for each fishery to evaluate overage or underage of the landed catch relative to the harvest objective. Postseason AIs for 1999-2005 are also listed in Table 3.3.

Table 3.3. Abundance indices for 1999 to 2006 for the SEAK, NBC, and WCVI troll fisheries.

| | Calibration | SEAK | | NBC | | WCVI | |
|------|--------------------------|-----------|------------|-----------|------------|-----------|------------|
| Year | Preseason/ Postseason | Preseason | Postseason | Preseason | Postseason | Preseason | Postseason |
| 1999 | 9902 / 0107 | 1.15 | 1.12 | 1.12 | 0.97 | 0.60 | 0.50 |
| 2000 | 0021 / 0107 | 1.14 | 1.10 | 1.00 | 0.95 | 0.54 | 0.47 |
| 2001 | 0107 / 0206 | 1.14 | 1.29 | 1.02 | 1.22 | 0.66 | 0.68 |
| 2002 | 0206 / 0308 | 1.74 | 1.82 | 1.45 | 1.63 | 0.95 | 0.92 |
| 2003 | 0308 / 0404 | 1.79 | 2.17 | 1.48 | 1.90 | 0.85 | 1.10 |
| 2004 | 0404 / 0506 | 1.88 | 2.06 | 1.67 | 1.83 | 0.90 | 0.98 |
| 2005 | 0506 / 0604 | 2.05 | 1.90 | 1.69 | 1.65 | 0.88 | 0.84 |
| 2006 | 0604 | 1.69 | | 1.53 | | 0.75 | |

The Agreement specifies the allowable catch for various values of the AI for each fishery. The allowable treaty catch by fishery and year based on pre- and postseason AIs and the actual (observed) catches are given in Table 3.4 and are shown in Figures 3.1 through 3.3; the solid line represents the relationship between AIs and allowable catch under Table 1 of the annex.

Table 3.4. Observed catches and postseason allowable catches for 1999 to 2005, and preseason allowable catches for 1999 to 2006, for AABM fisheries.

| | PST Treaty Allowable and Observed Catches | | | | | | | | |
|------|---|-----------------------------|---------------------------------|----------------------------|-----------------------------|----------------|----------------------------|-----------------------------|----------------|
| | SEAK (T, N, S) ¹ | | | NBC (T, S) | | | WCVI (T, S) | | |
| Year | Pre-season Allowable Catch | Post-season Allowable Catch | Observed Catch | Pre-season Allowable Catch | Post-season Allowable Catch | Observed Catch | Pre-season Allowable Catch | Post-season Allowable Catch | Observed Catch |
| 1999 | 192,800 | 184,200 | 198,842 | 145,600 | 126,100 | 86,726 | 128,300 | 107,000 | 36,413 |
| 2000 | 189,900 | 178,500 | 186,493 | 130,000 | 123,500 | 31,900 | 115,500 | 86,200 | 101,438 |
| 2001 | 189,900 | 250,300 | 186,919 | 132,600 | 158,900 | 43,500 | 141,200 | 145,500 | 117,670 |
| 2002 | 356,500 | 371,900 | 357,133 | 192,700 | 237,800 | 150,137 | 203,200 | 196,800 | 165,036 |
| 2003 | 366,100 | 439,600 | 380,152 | 197,100 | 277,200 | 191,657 | 181,800 | 268,900 | 175,821 |
| 2004 | 383,500 | 418,300 | 428,773 433,446 ² | 243,600 | 267,000 | 241,508 | 192,500 | 209,600 | 216,624 |
| 2005 | 416,400 | 387,400 | 386,707 | 246,600 | 240,700 | 243,606 | 188,200 | 179,700 | 202,662 |
| 2006 | 346,800 | | | 223,200 | | | 160,400 | | |

¹ Nomenclature is T for troll, N for net, and S for sport.

² The lower value results from subtracting a terminal exclusion catch for the Stikine River in 2004, which is in dispute.




Figure 3.1. Preseason allowable catches (open circles) and postseason catches (diamonds) in Southeast Alaska AABM fisheries, 1999-2005.




Figure 3.2. Preseason allowable catches (open circles) and postseason catches (diamonds) in Northern British Columbia troll and Queen Charlotte Islands recreational AABM fisheries, 1999-2005.




Figure 3.3. Preseason allowable catches (open circles) and postseason catches (diamonds) in West Coast Vancouver Island AABM fisheries, 1999-2005.

3.4.1.1 Model estimate of stock composition of AABM fisheries, 1979-2006

There are 30 model stocks (Appendix C). However, the majority of model catches in AABM fisheries are often composed of a smaller set of major stocks (Figures 3.4 through 3.6). The relative abundance for each major stock is shown in those graphs from CLB 0604. In general, postseason AIs had a peak during the late 1980s and another in 2003 and 2004. For all three AABM fisheries, the postseason 2005 AI and preseason 2006 AI dropped from the higher levels seen in 2003 and 2004.




Figure 3.4. Total abundance indices for the SEAK troll fishery with annual stock composition indicated by abundance indices for major model stocks from CLB 0604.

The major model stocks contributing to the SEAK AIs are: WCVI Natural and Hatchery, Upriver Brights, North/Central BC, and Oregon Coastal (Figure 3.4). The 2006 forecasts for all four of these stock groups are lower than the 2005 returns. The “other” category is primarily driven by Upper Georgia Strait, Columbia River Summers, Mid Columbia River Brights and Fraser Early.




Figure 3.5. Total abundance indices for the NBC troll fishery with annual stock composition indicated by abundance indices for major model stocks from CLB 0604.

The major model stock groups contributing to the NBC AABM fishery AIs are: WCVI Natural and Hatchery, Upriver Brights, Oregon Coastal, North/Central BC, and Washington Coastal Wild and Hatchery (Figure 3.5). The 2006 forecasts for four of these stock groups are lower than the estimated 2005 returns, while that for Washington Coastal is the same. The “other” category is primarily driven by Columbia River Summers, Mid Columbia River Brights and Willamette Springs. For these stocks, the 2006 forecasts are similar in magnitude to the estimated returns in 2005.




Figure 3.6. Total abundance indices for the WCVI troll fishery with annual stock composition indicated by abundance indices for major model stocks from CLB 0604.

The major model stock groups in the WCVI fishery are: Fraser Late, Puget Sound, Upriver Brights, and Columbia River Tules (Figure 3.6). The 2006 forecasts are for lower abundances of Columbia River Tules and Upriver Brights and similar numbers for the other two stock groups, relative to 2005 returns. The “Other” category is comprised primarily of Columbia River Summers and Oregon Coastal fish.

3.4.2 Overages and Underages

Until an approach for full implementation of overage/underage provisions has been developed and accepted by the PSC, the Commissioners have instructed the CTC to track and report overages and underages relative to agreed-upon harvest objectives.

3.4.2.1 AABM Fisheries

Table 3.5 shows the differences between the postseason allowable catches and the observed catches in AABM fisheries for 1999–2004, and the cumulative differential for those years. All three AABM fisheries have cumulative underages. In SEAK, observed catches have been below final allowable catches for four of the seven years; the cumulative differential is –4.7% or -4.5%. In NBC, observed catches have been below the final allowable catches in six of the seven years; the cumulative differential is –30.9%. In WCVI, observed catches have been below allowable catches in four of the seven years; the cumulative differential is –14.8%.

Table 3.5. Deviations in numbers of Chinook salmon and percentages from catch targets derived from the first postseason AI (Table 3.2) for Pacific Salmon Treaty AABM fisheries in 1999 to 2005.

| Year | SEAK | | NBC | | WCVI | |
|------|-----------------------------------|--------------------|----------------|--------------------|----------------|--------------------|
| | Number of Fish | Percent Difference | Number of Fish | Percent Difference | Number of Fish | Percent Difference |
| 1999 | +14,642 | +7.9% | -39,374 | -31.2% | -70,587 | -66.0% |
| 2000 | +7,993 | +4.5% | -91,600 | -74.2% | +15,238 | +17.7% |
| 2001 | -63,381 | -25.3% | -115,400 | -72.6% | -27,830 | -19.1% |
| 2002 | -14,767 | -4.0% | -87,663 | -36.9% | -31,764 | -16.1% |
| 2003 | -59,448 | -13.5% | -85,543 | -30.9% | -93,079 | -34.6% |
| 2004 | +10,473 +15,146 | +2.5% +3.6% | -25,492 | -9.5% | +7,024 | +3.4% |
| 2005 | -693 | -0.2% | +2,906 | +1.2% | +22,962 | +12.8% |
| Cum. | -105,181 -100,508 ¹ | -4.7% -4.5% | -442,166 | -30.9% | -178,036 | -14.9% |

¹ The lower value results from subtracting a terminal exclusion catch for the Stikine River in 2004, which is in dispute.

3.4.2.2 ISBM Indices by Stock

For ISBM fisheries, the Agreement specifies that Canada and the United States will reduce base period exploitation rates on specified stocks by 36.5% and 40%, equivalent to ISBM indices of 63.5% and 60% percent, respectively. This requirement is referred to as the ‘general obligation’ and does not apply to stocks that achieve their CTC agreed escapement goal. Estimated ISBM fishery indices are shown in Table 3.6 for Canadian fisheries and Table 3.7 for U.S. fisheries. Both tables present CWT-based indices for 2004, and Chinook model-based predicted indices for 2006. The agreement specifies that the indices for postseason assessment be assessed using the CWT-based estimates, 2004 is the most recent analysis available. CWT-based indices for 2000-2004 and model-based indices for 2000-2006 are presented in Appendix D.

3.4.2.2.1 CWT-based Indices in 2004

Canadian ISBM indices from the CWT-based estimates for 2004 were reduced more than required under the agreement for all stocks or stock groups (Table 3.6). Several inconsistencies were identified in the way these indices had been computed in the past, as noted in the footnotes 4-9 in Table 3.6. Most of them were inconsistencies between the way indices had been calculated by the model versus in the CWT analysis. However, in the case of Lower Georgia Strait, Nanaimo was dropped from the CWT-based index because of concern about the adequacy of base-period data. In addition, Nanaimo and Cowichan stocks are no longer reported separately in the model-based index because there is no way to split the two stocks in the base period.

Three stocks, Cowichan, Nanaimo, and Green River, exceeded the 0.60 benchmark established under the Agreement (Table 3.7). None of these stocks had escapement goals in 2004. The

Cowichan escapement goal was accepted in 2005. Of the remaining 13 stocks for which the ISBM index was calculated, eight exceeded their escapement goals and five were below the limit. Figures 3.7 and 3.8 show the historical ISBM indices based on CWT recoveries for 1999-2004.




Figure 3.7. ISBM indices for Canadian fisheries for 1999-2004. The solid horizontal line is an index value of 0.635.




Figure 3.8. ISBM indices for U.S. fisheries for 1999-2004.

3.4.2.2.2 Predicted ISBM Indices for 2006

Model projected indices (Table 3.6) show that the Canadian ISBM indices are expected to be below 0.635 for all Canadian stocks other than WCVI fall stocks. Canadian indices are projected to be above 0.635 for Puget Sound stocks and below 0.635 for other U.S. stocks. In the southern U.S. fisheries (Table 3.7) 10 stocks are projected to have ISBM index values over 0.60, but with the exception of Lake Washington, the stocks with projected ISBM indices greater than 0.60 have agreed escapement goals and have been meeting these goals.

Table 3.6. Canadian 2004 ISBM indices based on CWT and the 2006 indices predicted from the PSC Chinook Model.

| Stock Group | Escapement Indicator Stock | Canadian ISBM Indices | |
|---|--|-----------------------|------------------------|
| | | CWT Indices for 2004 | Model Indices for 2006 |
| Lower Strait of Georgia | Cowichan ² | 0.284 ^{1,4} | |
| | Nanaimo | NA ⁵ | 0.590 ⁶ |
| Fraser Late | Harrison River ² | 0.032 ⁷ | 0.294 |
| North Puget Sound Natural Springs | Nooksack | NA | 0.993 |
| | Skagit | NA | 0.993 |
| Upper Strait of Georgia | Klinaklini, Kakweikan, Wakeman, Kingcome, Nimpkish | 0.018 | 0.584 |
| Fraser Early (spring and summers) | Upper Fraser, Mid Fraser, Thompson | NA | 0.610 |
| West Coast Vancouver Island Falls | WCVI (Artlish, Burman, Kauok, Tahsis, Tashish, Marble) | 0.488 ⁸ | 1.082 |
| Puget Sound Natural Summer / Falls | Skagit | NA | 1.092 |
| | Stillaguamish | 0.027 | 1.116 |
| | Snohomish | NA | 1.101 |
| | Lake Washington | NA | 0.914 ⁹ |
| | Green River | 0.162 | 0.914 ⁹ |
| North / Central B. C. | Yakoun, Nass, Skeena, Area 8 | NA | 0.626 |
| Washington Coastal Fall Naturals ³ | Hoko, Grays Harbor, Queets ² , Hoh ² , Quillayute ² | NA | 0.363 |
| Columbia River Falls ³ | Upriver Brights ² | NA | 0.523 |
| | Deschutes | NA | 0.523 |
| | Lewis ² | NA | 0.315 |
| Columbia R Summers ³ | Mid-Columbia Summers ² | NA | 0.335 |
| Far North Migrating OR Coastal Falls ³ | Nehalem ² , Siletz ² , Siuslaw ² | NA | 0.515 |

¹ Not available (NA) because of insufficient data (lack of stock specific tag codes, base period CWT recoveries, etc).

² Stock or stock group with an agreed CTC escapement goal.

³ Stock group listed in Annex 4, Chapter 3, Attachment V.

⁴ An inconsistency was discovered between the approaches used to calculate the model-based and CWT-based indices. The former included harvest rates for terminal sport while the latter did not. Terminal sport harvest rates are now included in the calculation of both indices. Further review is yet required to determine whether the base period terminal sport harvest rates obtained from analyses of Big Qualicum CWT recoveries adequately represent impacts that would have occurred on Cowichan Chinook.

⁵ Several problems have been identified in the approach previously used to calculate the CWT-based indices for Nanaimo Chinook. Until these problems are resolved, indices for this stock will not be reported.

⁶ Although model-based indices were previously calculated separately for Cowichan and Nanaimo, these did not adequately represent impacts on either LGS stock because the model-based data represent an aggregate of the two stocks and methods do not currently exist to correctly disaggregate these data for calculation of the ISBM values. Until such methods are developed, a single index value only will be reported representing the aggregate.

⁷ The terminal sport harvest rates for Chilliwack Hatchery Chinook, the indicator stock, were removed from the calculation for the Harrison River naturals because sport harvest has been essentially zero on the natural population.

⁸ An inconsistency was discovered between the approaches used to calculate the model-based and CWT-based indices. The former included harvest rates for terminal sport while the latter did not. Terminal sport harvest rates are now included in the calculation of both indices. A more extended review of the indices for WCVI Chinook will be carried out to determine whether they adequately represent impacts on the WCVI wild aggregate.

⁹ For Canadian ISBM fisheries, the Lake Washington and Green stocks are assumed to have the same distribution and index value.

Table 3.7. U.S. 2004 ISBM indices based on CWT and the 2006 indices predicted from the PSC Chinook Model. Order of the stock groups correspond to Annex 4, Chapter 3, Attachment V of the PST 1999 Revised Annexes.

| Stock Group | Escapement Indicator Stock | U.S. ISBM Indices | |
|--|--|----------------------|------------------------|
| | | CWT Indices for 2004 | Model Indices for 2006 |
| Washington Coastal Fall Naturals | Hoko | NA ¹ | 0.442 |
| | Grays Harbor | 0.530 | 0.544 |
| | Queets ⁴ | 0.840 | 1.022 |
| | Hoh ⁴ | 1.220 | 1.493 |
| | Quillayute ⁴ | 1.150 | 0.673 |
| Columbia River Falls | Upriver Brights ⁴ | 1.740 | 0.814 |
| | Deschutes | 0.510 | 0.437 |
| | Lewis ⁴ | 0.170 | 1.861 |
| Puget Sound Natural Summer / Falls | Skagit | NA | 0.258 |
| | Stillaguamish | 0.10 | 0.493 |
| | Snohomish | NA | 0.199 |
| | Lake Washington | NA | 0.613 |
| | Green R | 1.010 | 0.361 |
| Fraser Late | Harrison River ⁴ | 0.320 | 0.787 |
| Columbia R Summers | Mid-Columbia Summers ⁴ | 2.690 | 0.696 |
| Far North Migrating OR Coastal Falls | Nehalem ⁴ | 1.800 | 1.912 |
| | Siletz ⁴ | 2.290 | 1.237 |
| | Siuslaw ⁴ | 1.030 | 1.095 |
| North Puget Sound Natural Springs | Nooksack | NA | 0.121 |
| | Skagit | NA | 0.161 |
| Lower Strait of Georgia ³ | Cowichan ⁴ , | 7.250 | 0.271 |
| | Nanaimo | 7.250 | 0.271 |
| Upper Strait of Georgia ³ | Klinaklini, Kakweikan, Wakeman, Kingcome, Nimpkish | NA | NC ² |
| Fraser Early (spring and summers) ³ | Upper Fraser, Mid Fraser, Thompson | NA | 0.214 |
| West Coast Vancouver Island Falls ³ | WCVI (Artlish, Burman, Kauok, Tahsis, Tashish, Marble) | NA | 0.128 |
| North / Central B. C. ³ | Yakoun, Nass, Skeena, Area 8 | NA | NC |

¹ NA means not available because of insufficient data (lack of stock specific tag codes, base period CWT recoveries, etc).

² NC means that the current model assumes the stock is not caught in U.S. ISBM fisheries.

³ Stock group listed in Annex 4, Chapter 3, Attachment IV.

⁴ Stock with an agreed CTC escapement goal.

3.5 MODEL CALIBRATION EVALUATION

Previous reports included evaluations of model performance for the most current model year, including comparisons of model estimates of catch and escapement/terminal run sizes to actual estimates of catch and escapement/terminal run size. This year, the model catches and stock escapements or terminal run sizes estimated by CLB 0604 were evaluated as were other aspects of the calibration. The calibration was distributed to the CTC membership for review and subsequently approved. Correlations between model and CWT fishery indices are normally presented. However, while these comparisons were made as part of the normal calibration checking process, the results are not presented in this report.

Fishery mortality indices generated by CLB 0604 can be compared to the CWT-based exploitation rate analysis. Model and CWT-based fishery mortality indices use the same equation, but the former are derived from model estimates of catch for all model stocks instead of CWT recovery data from specific exploitation rate indicator stocks. The CWT fishery mortality indices are considered to be the most accurate. Two types of fishery indices are presented; reported catch and total mortality. In general, the model results are closely associated with the CWT-based indices and changes in fishery exploitation rates as indicated in Figures 3.9 through 3.14. The SEAK fishery mortality index from the model closely follows the trend of the CWT derived estimate from 1979 through 1989 for both landed catch and total mortality (Figures 3.9 and 3.10). Between 1989 and 2000, the model estimate of both landed catch and total mortality indices is less than the CWT-derived estimate for most years but since 2001, the model estimate is noticeably higher. Since 1990, the model estimates also show less variability compared to the CWT-derived indices.




Figure 3.9. Estimated CWT (through 2004) and model landed catch fishery indices (through 2005) for the SEAK troll fishery.




Figure 3.10. Estimated CWT (through 2004) and model total mortality fishery indices (through 2005) for the SEAK troll fishery.

The model-derived fishery mortality indices for NBC generally follow the same trend as CWT-derived indices (Figures 3.11 and 3.12). However, since 1991, the model-based estimates have exceeded the CWT-derived estimates in all but three years for both landed catch and total mortality indices. Since 2001, this difference has been noticeably large.




Figure 3.11. Estimated CWT (through 2004) and model landed catch fishery indices (through 2005) for the NBC troll fishery.




Figure 3.12. Estimated CWT (through 2004) and model total mortality fishery indices (through 2005) for the NBC troll fishery.

Since the base period, the model-derived landed catch fishery index estimates and trends for the WCVI troll fishery have been similar to those derived from CWTs. However, from 1987 through

1995, the model estimates are consistently greater than the CWT-based estimates (Figures 3.13 and 3.14). Starting in 2000, model and CWT estimates have diverged significantly for both landed catch and total mortality, with CWT indices being consistently higher than model indices.




Figure 3.13. Estimated CWT (through 2004) and model landed catch fishery indices (through 2005) for the WCVI troll fishery.




Figure 3.14. Estimated CWT (through 2004) and model total mortality fishery indices (through 2005) for the WCVI troll fishery.

3.6 EVALUATION OF MARK-SELECTIVE FISHERIES.

There have been mark-selective fisheries (MSF) for Chinook salmon in the Strait of Juan de Fuca Washington sport fishery since 2003, in the Columbia River net fisheries since 2002, and in Columbia River spring Chinook sport fisheries since 2000. Double index tag (DIT) groups are comprised of paired releases of marked and unmarked fish with CWTs. Seven Puget Sound fall Chinook stocks and one Columbia River stock have DIT groups. The DIT is used as a monitoring tool to test the hypothesis that there are differences between the marked and unmarked tagged groups due to MSFs and also to estimate mortalities of unmarked fish in MSFs.

A significant change in the ratio of unmarked to marked DIT groups at hatchery escapement can indicate that mark-selective fisheries have differentially impacted DIT pairs. Statistical Z-tests were used to compare the return rate of the marked and unmarked brood-age groups for seven Puget Sound DIT groups subject to MSFs in 2003 and 2004. Out of 52 tests for brood-ages with marked and unmarked returns to the hatchery, only 6 were significant (Figure 3.15), and the actual calculated differences were small. This indicates that the Area 5 and 6 MSF did not result in significant differences in hatchery escapement of DIT groups. For this reason, the estimates of exploitation rate of marked tagged groups were used in CTC analyses this year.

Although a DIT group is available for Willamette spring Chinook, a similar analysis could not be carried out as all unmarked fish were not sampled upstream of the North Fork trap. All marked fish were taken to the hatchery and sampled or returned to the river. Methods to directly estimate impacts of MSFs on the Willamette spring stock and any stock represented by single index tag (SIT) groups are not available.

Methods for estimating exploitation rates on unmarked fish for DIT and SIT stocks and results of the analyses are presented in Appendix J. Currently the Selective Fishery Evaluation Committee (SFEC) is preparing a report on the impact of MSFs on Chinook and coho salmon DITs.




Figure 3.15. Percent of release returning to hatchery by stock, brood and age for marked and unmarked DIT groups. * indicates a pair where the percent returns were significantly different

REFERENCES CITED

- Ames, J. and D. E. Phinney. 1977. 1977 Puget Sound summer-fall chinook methodology: escapement estimates and goals, run size forecasts, and in-season run size updates. Washington Department of Fisheries Technical Report 29. Olympia, Washington.
- CBFWA (Columbia Basin Fish and Wildlife Authority). 1990. Integrated system plan for salmon and steelhead production in the Columbia River basin. Portland, Oregon.
- CCMP (Comprehensive Chinook Management Plan). 2004. Comprehensive Chinook management plan for Puget Sound Chinook: harvest management component. Northwest Indian Fisheries Commission. Olympia, Washington.
- CRFMP. 1988. Columbia River Fisheries Management Plan.
- CTC (Chinook Technical Committee). 1997. Incidental fishery mortality of Chinook salmon: Mortality rates applicable to Pacific Salmon Commission Fisheries Report TCCHINOOK (97)-1. Vancouver, British Columbia.
- CTC (Chinook Technical Committee). 1999. Maximum sustained yield or biologically based escapement goals for selected Chinook salmon stocks used by the Pacific Salmon Commission's Chinook Technical Committee for escapement assessment. Pacific Salmon Commission, Report TCCHINOOK (99)-3. Vancouver, British Columbia.
- CTC (Chinook Technical Committee). 2002. Catch and escapement of Chinook salmon under Pacific Salmon Commission jurisdiction 2001. Pacific Salmon Commission, Report TCCHINOOK (02)-1. Vancouver, British Columbia.
- CTC (Chinook Technical Committee). 2004a. Annual exploitation rate analysis and model calibration. Pacific Salmon Commission, Report TCCHINOOK (04)-4. Vancouver, British Columbia.
- CTC (Chinook Technical Committee). 2004b. Standardized fishery regimes for Southeast Alaska Chinook fisheries. Pacific Salmon Commission, Report TCCHINOOK (04)-3. Vancouver, British Columbia.
- CTC (Chinook Technical Committee). 2004c. Estimation and application of incidental fishing mortality in Chinook salmon management under the 1999 Agreement of the Pacific Salmon Treaty. Pacific Salmon Commission, Report TCCHINOOK (04)-1. Vancouver, British Columbia.
- CTC (Chinook Technical Committee). 2005a. Catch and escapement of Chinook salmon under Pacific Salmon Commission jurisdiction, 2003. Pacific Salmon Commission, Report TCCHINOOK (05)-2. Vancouver, British Columbia.
- CTC (Chinook Technical Committee). 2005b. Annual exploitation rate analysis and model calibration. Pacific Salmon Commission, Report TCCHINOOK (05)-3. Vancouver, British Columbia.
- Ericksen, R. P. and S. A. McPherson. 2003. Biological escapement goal for Chilkat River Chinook salmon. Alaska Department of Fish and Game, Division of Sport Fish, Fishery Manuscript 03-01, Juneau.
- Farwell, M. K., R. E. Bailey, and B. Rosenberger. 1999. Enumeration of the 1995 Nicola River Chinook salmon escapement. Canadian Manuscript Report Fisheries and Aquatic Science 2491:44p.
- Freeman, G.M., S. A. McPherson and D.L. Magnus. 2001. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Keta River, 2000. Alaska Department of Fish and Game, Division of Sport Fish, Fishery Data Series 01-19, Anchorage, Alaska.
- Joint Coho DIT Analysis Workgroup. 2003. Analysis of coho salmon double index (DIT) data for the brood years 1995-1997. Northwest Fishery Resource Bulletin. Project Report Series No. 12.

- McPherson, S. A., D. R. Bernard, and J. H. Clark. 2000. Optimal production of Chinook salmon from the Taku River. Alaska Department of Fish and Game, Sport Fish Division, Fishery Manuscript 00-2. Anchorage, Alaska.
- McPherson, S. A. and J. H. Clark. 2001. Biological escapement goal for King Salmon River Chinook salmon. Alaska Department of Fish and Game, Regional Information Report 1J01-40. Anchorage, Alaska.
- Pahlke, K. A. 2003. Escapements of Chinook salmon in Southeast Alaska and transboundary rivers in 2001. Alaska Department of Fish and Game, Division of Sport Fish, Fishery Data Series 03-11, Anchorage, Alaska.
- Pahlke, K. A., and P. Etherton. 1999. Chinook salmon research on the Stikine River, 1997. Alaska Department of Fish and Game, Division of Sport Fish, Fishery Data Series No. 99-6, Anchorage, Alaska.
- PFMC (Pacific Fishery Management Council). 2003. Review of 2002 Ocean Salmon Fisheries. Pacific Fishery Management Council. Portland, Oregon.
- PFMC (Pacific Fishery Management Council). 2005. Preseason Report III: Analyses of Council Adopted Management Measures for 2000 Ocean Fisheries. Pacific Fishery Management Council. Portland, Oregon.
- SFEC. 2002. Investigation of methods to estimate mortalities of unmarked salmon in mark-selective fisheries through the use of double index tag groups. PSC TCSFEC(02)-1
- TAC (U.S. v Oregon Technical Advisory Committee). 1999. All Species Review: Columbia River Fish Management Plan.
- Weller, Jan L. and Scott A. McPherson. 2003. A mark-recapture experiment to estimate the escapement of Chinook salmon in the Unuk River, 2002. Alaska Department of Fish and Game, Fishery Data

APPENDICES

Appendix A. Landed Chinook catches by region and gear from 1975-2005.

| | |
|--|----|
| Appendix A.1. Southeast Alaska (SEAK) Chinook catches, 1975-2005. | 86 |
| Appendix A.2. Northern British Columbia (NBC) Chinook catches, 1975-2005. | 87 |
| Appendix A.3. Central British Columbia (CBC) Chinook catches, 1975-2005. | 88 |
| Appendix A.4. West Coast Vancouver Island (WCVI) Chinook catches, 1975-2005. | 89 |
| Appendix A.5. Johnstone Strait Chinook catches, 1975-2005. | 90 |
| Appendix A.6. Strait of Georgia/Fraser Chinook catches, 1975-2005. | 91 |
| Appendix A.7. Canada - Strait of Juan de Fuca Chinook catches, 1975-2005. | 92 |
| Appendix A.8. Washington - Strait of Juan de Fuca Chinook catches, 1975-2005. | 93 |
| Appendix A.9. Washington - San Juan Chinook catches, 1975-2005. | 94 |
| Appendix A.10. Washington – Other Puget Sound Chinook catches, 1975-2005. | 95 |
| Appendix A.11. Washington – Inside Coastal Chinook catches, 1975-2005. | 96 |
| Appendix A.12. Washington/Oregon North of Cape Falcon Chinook catches, 1975-2005. | 97 |
| Appendix A.13. Columbia River Chinook catches, 1975-2005. | 98 |
| Appendix A.14. Oregon Chinook catches, 1975-2005. | 99 |

Appendix A.1. Southeast Alaska (SEAK) Chinook catches, 1975-2005.

| Year | Southeast Alaska | | | | | | |
|------|------------------|--------|--------|---------|--------|--------------------|---------------------|
| | Troll | Net | Sport | Total | Add-on | Terminal Exclusion | Treaty Catch |
| 1975 | 287,342 | 13,365 | 17,000 | 317,707 | - | - | - |
| 1976 | 231,239 | 10,523 | 17,000 | 258,762 | - | - | - |
| 1977 | 271,735 | 13,443 | 17,000 | 302,178 | - | - | - |
| 1978 | 375,919 | 25,492 | 17,000 | 418,411 | - | - | - |
| 1979 | 337,672 | 28,388 | 16,581 | 382,641 | - | - | - |
| 1980 | 303,643 | 20,114 | 20,213 | 343,970 | - | - | - |
| 1981 | 248,782 | 18,952 | 21,300 | 289,034 | - | - | - |
| 1982 | 241,938 | 46,992 | 25,756 | 314,686 | - | - | - |
| 1983 | 269,821 | 19,516 | 22,321 | 311,658 | - | - | - |
| 1984 | 235,622 | 32,405 | 22,050 | 290,077 | - | - | - |
| 1985 | 215,811 | 33,870 | 24,858 | 274,539 | 6,246 | - | 268,293 |
| 1986 | 237,703 | 22,099 | 22,551 | 282,353 | 11,091 | - | 271,262 |
| 1987 | 242,562 | 15,532 | 24,324 | 282,418 | 17,095 | - | 265,323 |
| 1988 | 231,364 | 21,788 | 26,160 | 279,312 | 22,525 | - | 256,787 |
| 1989 | 235,716 | 24,245 | 31,071 | 291,032 | 21,510 | - | 269,522 |
| 1990 | 287,939 | 27,712 | 51,218 | 366,869 | 45,873 | - | 320,996 |
| 1991 | 264,106 | 34,864 | 60,492 | 359,462 | 61,476 | - | 297,986 |
| 1992 | 183,759 | 32,140 | 42,892 | 258,791 | 36,811 | - | 221,980 |
| 1993 | 226,866 | 27,991 | 49,246 | 304,103 | 32,910 | - | 271,193 |
| 1994 | 186,331 | 35,654 | 42,365 | 264,350 | 29,185 | - | 235,165 |
| 1995 | 138,117 | 47,955 | 49,667 | 235,739 | 58,800 | - | 176,939 |
| 1996 | 141,452 | 37,298 | 57,509 | 236,259 | 72,599 | 8,663 | 154,997 |
| 1997 | 246,409 | 25,069 | 71,524 | 343,002 | 46,463 | 9,843 | 286,696 |
| 1998 | 192,066 | 23,514 | 55,013 | 270,593 | 25,021 | 2,420 | 243,152 |
| 1999 | 146,219 | 32,720 | 72,081 | 251,020 | 47,725 | 4,453 | 198,842 |
| 2000 | 158,717 | 41,400 | 63,173 | 263,290 | 74,316 | 2,481 | 186,493 |
| 2001 | 153,280 | 40,163 | 72,291 | 265,734 | 77,287 | 1,528 | 186,919 |
| 2002 | 325,308 | 31,689 | 69,537 | 426,534 | 68,164 | 1,237 | 357,133 |
| 2003 | 330,692 | 39,374 | 69,370 | 439,436 | 57,228 | 2,056 | 380,152 |
| 2004 | 354,664 | 64,038 | 87,505 | 506,207 | 72,025 | 5,409 736 | 428,7731 433,446 |
| 2005 | 338,437 | 73,066 | 84,279 | 495,782 | 64,102 | 44,973 | 386,707 |

Troll, net, sport and total catches include catch of SEAK hatchery-origin fish; catches that count towards the all-gear ceiling (with hatchery add-on subtracted) are shown in the "treaty catch" column.

“-“ = not applicable.

¹ The value on top excludes District 108 Stikine catch above base levels. The value below includes it.

Appendix A.2. Northern British Columbia (NBC) Chinook catches, 1975-2005.

| Year | Northern British Columbia | | | | | | |
|------|--------------------------------|-----------------|-------------------|-----------|---------------------------------|------------------------------|---------|
| | | | Tidal Sport | | | | |
| | Area 1-5 Troll ¹ | Area 1-5 Net | Areas 1,2E, 2W | Areas 3-5 | Area 1-5 Freshwater Sport | Area 1-5 First Nations | Total |
| 1975 | 228,121 | 25,095 | NA | NA | NA | 4,055 | 257,271 |
| 1976 | 190,267 | 16,105 | NA | NA | NA | 2,791 | 209,163 |
| 1977 | 130,899 | 44,196 | 106 | 1,670 | 2,158 | 6,998 | 186,027 |
| 1978 | 146,054 | 27,924 | 125 | 1,668 | 6,610 | 5,363 | 187,744 |
| 1979 | 147,576 | 40,640 | 0 | 2,523 | 1,960 | 5,266 | 197,965 |
| 1980 | 157,198 | 26,895 | 200 | 3,867 | 4,515 | 10,121 | 202,796 |
| 1981 | 153,065 | 41,724 | 184 | 2,760 | 2,613 | 11,115 | 211,461 |
| 1982 | 173,472 | 44,844 | 215 | 3,760 | 2,726 | 13,255 | 238,272 |
| 1983 | 162,837 | 17,134 | 90 | 4,092 | 5,374 | 15,532 | 205,059 |
| 1984 | 185,134 | 31,321 | 171 | 2,300 | 3,426 | 11,408 | 233,760 |
| 1985 | 165,845 | 39,562 | 600 | 3,600 | 3,186 | 15,794 | 228,587 |
| 1986 | 175,715 | 23,902 | 1,153 | 3,950 | 4,410 | 24,448 | 233,578 |
| 1987 | 177,457 | 18,357 | 2,644 | 4,150 | 3,625 | 16,329 | 222,562 |
| 1988 | 152,369 | 31,339 | 7,059 | 4,300 | 3,745 | 21,727 | 220,539 |
| 1989 | 207,679 | 38,623 | 20,652 | 4,150 | 5,247 | 21,023 | 297,374 |
| 1990 | 154,109 | 28,359 | 16,827 | 4,300 | 4,090 | 27,105 | 234,790 |
| 1991 | 194,018 | 40,899 | 15,047 | 4,256 | 4,764 | 23,441 | 282,425 |
| 1992 | 142,340 | 35,716 | 21,358 | 6,250 | 6,182 | 27,012 | 238,858 |
| 1993 | 161,686 | 33,944 | 25,297 | 3,279 | 7,813 | 21,353 | 253,372 |
| 1994 | 164,581 | 22,032 | 28,973 | 3,171 | 3,093 | 15,949 | 237,799 |
| 1995 | 56,857 | 18,076 | 22,531 | 2,475 | 3,503 | 13,635 | 117,077 |
| 1996 | 21 | 28,894 | 670 | 3,382 | 1,250 | 13,345 | 47,562 |
| 1997 | 83,488 | 20,415 | 27,738 | 0 | NA | 14,610 | 146,251 |
| 1998 | 107,837 | 7,144 | 34,130 | 4,750 | NA | 20,622 | 174,483 |
| 1999 | 56,499 | 10,094 | 30,227 | 11,700 | NA | 27,399 | 135,919 |
| 2000 | 9,800 | 22,329 | 22,100 | 8,600 | NA | 23,476 | 86,305 |
| 2001 | 13,100 | 25,424 | 30,400 | 11,000 | NA | 23,508 | 103,432 |
| 2002 | 103,038 | 14,902 | 47,100 | 8,000 | NA | 14,125 | 187,165 |
| 2003 | 137,357 | 14,730 | 54,300 | NA | 5,711 ² | 20,950 | 233,048 |
| 2004 | 167,508 | 16,187 | 74,000 | NA | NA | 20,548 | 278,243 |
| 2005 | 174,806 | 6,850 | 68,800 | NA | NA | 17,553 | 267,770 |

¹ Since 1998, the catch accounting year for troll fisheries was set from October 1-September 30. To make comparisons to previous years more meaningful, the same catch accounting period was applied for years prior to 1998.

² Estimate of lower Skeena River sport catch only.

Note that Troll (Areas 1-5) and Tidal Sport (Areas 1, 2E, 2W) are the components of the NBC AABM fishery. Net catch excludes jacks and small red-fleshed Chinook.

NA=not available

Appendix A.3. Central British Columbia (CBC) Chinook catches, 1975-2005.

| Year | Central British Columbia | | | | | |
|------|--------------------------|--------|-------------|------------------|---------------|---------|
| | Troll ¹ | Net | Tidal Sport | Freshwater Sport | First Nations | Total |
| 1975 | 135,470 | 40,985 | NA | NA | NA | 176,455 |
| 1976 | 145,204 | 32,669 | NA | NA | NA | 177,873 |
| 1977 | 122,689 | 32,409 | 4,773 | 1,544 | 6,972 | 168,387 |
| 1978 | 91,025 | 35,708 | 5,694 | 1,770 | 7,944 | 142,141 |
| 1979 | 107,884 | 50,445 | 5,225 | 1,940 | 7,585 | 173,079 |
| 1980 | 95,377 | 27,715 | 4,802 | 988 | 6,240 | 135,122 |
| 1981 | 69,247 | 18,912 | 3,490 | 1,261 | 5,701 | 98,611 |
| 1982 | 69,748 | 32,419 | 5,419 | 1,293 | 9,112 | 117,991 |
| 1983 | 97,447 | 12,556 | 4,271 | 821 | 6,442 | 121,537 |
| 1984 | 78,120 | 4,630 | 4,354 | 1,332 | 9,736 | 98,172 |
| 1985 | 27,090 | 12,391 | 3,943 | 823 | 6,019 | 50,266 |
| 1986 | 54,407 | 23,032 | 4,566 | 1,245 | 6,353 | 89,603 |
| 1987 | 65,776 | 10,893 | 3,933 | 1,563 | 6,296 | 88,461 |
| 1988 | 36,125 | 12,886 | 3,596 | 1,496 | 6,000 | 60,103 |
| 1989 | 21,694 | 6,599 | 3,438 | 4,526 | 8,992 | 45,249 |
| 1990 | 29,882 | 18,630 | 4,053 | 5,626 | 9,811 | 68,002 |
| 1991 | 29,843 | 15,926 | 4,409 | 3,335 | 8,801 | 62,314 |
| 1992 | 47,868 | 18,337 | 4,891 | 3,204 | 8,533 | 82,833 |
| 1993 | 23,376 | 10,579 | 6,114 | 2,880 | 9,095 | 52,044 |
| 1994 | 18,976 | 14,424 | 4,303 | 973 | 5,383 | 44,059 |
| 1995 | 5,819 | 11,007 | 2,172 | 1,180 | 3,501 | 23,679 |
| 1996 | 0 | 6,829 | 2,936 | 3,986 | 6,922 | 20,673 |
| 1997 | 12,351 | 3,575 | 8,524 | 1,139 | 9,764 | 35,353 |
| 1998 | 2,198 | 5,355 | 5,514 | 779 | 6,671 | 20,517 |
| 1999 | 2,074 | 4,320 | 10,300 | NA ² | 5,440 | 22,134 |
| 2000 | 0 | 3,210 | 7,400 | NA ² | 4,576 | 15,186 |
| 2001 | 0 | 6,462 | 7,650 | 1,024 | 5,435 | 20,571 |
| 2002 | 481 | 4,676 | 7,330 | 723 | 3,292 | 16,502 |
| 2003 | 20 | 2,806 | 8,385 | 491 | 3,173 | 14,875 |
| 2004 | 0 | 6,324 | 10,677 | 524 | 4,003 | 21,528 |
| 2005 | 0 | 6,323 | 9,017 | 809 | 4,180 | 20,329 |

¹ Since 1998, the catch accounting year for troll fisheries was set from October 1-September 30. To make comparisons to previous years more meaningful, the same catch accounting period was applied for years prior to 1998.

² freshwater catch included with tidal catch

Net catch excludes jacks and small red-fleshed Chinook.

NA=not available

Appendix A.4. West Coast Vancouver Island (WCVI) Chinook catches, 1975-2005.

| Year | West Coast Vancouver Island | | | | | | |
|------|-----------------------------|--------|---------------------|-------------|------------------|---------------|---------|
| | | | Tidal Sport | Tidal Sport | | | |
| | Troll ¹ | Net | Inside ² | Outside | Freshwater Sport | First Nations | Total |
| 1975 | 546,214 | 19,233 | NA | - | NA | NA | 565,447 |
| 1976 | 665,010 | 17,492 | NA | - | NA | NA | 682,502 |
| 1977 | 545,742 | 13,745 | NA | - | NA | NA | 559,487 |
| 1978 | 568,705 | 25,143 | NA | - | NA | NA | 593,848 |
| 1979 | 477,222 | 35,623 | 7,964 | - | NA | NA | 520,809 |
| 1980 | 486,303 | 34,732 | 8,539 | - | NA | NA | 529,574 |
| 1981 | 423,266 | 36,411 | 11,230 | - | NA | NA | 470,907 |
| 1982 | 538,510 | 41,172 | 17,100 | - | NA | NA | 596,782 |
| 1983 | 395,636 | 37,535 | 28,000 | - | NA | NA | 461,171 |
| 1984 | 471,294 | 43,792 | 44,162 | - | NA | NA | 559,248 |
| 1985 | 345,937 | 11,089 | 21,587 | - | NA | NA | 378,613 |
| 1986 | 350,227 | 3,276 | 13,158 | - | NA | NA | 366,661 |
| 1987 | 378,931 | 478 | 38,283 | - | NA | NA | 417,692 |
| 1988 | 408,668 | 15,438 | 35,820 | - | NA | NA | 459,926 |
| 1989 | 203,751 | 40,321 | 55,239 | - | NA | NA | 299,311 |
| 1990 | 297,858 | 29,578 | 69,723 | - | NA | 1,199 | 398,358 |
| 1991 | 203,035 | 60,797 | 85,983 | - | NA | 41,322 | 391,137 |
| 1992 | 340,146 | 9,486 | 46,968 | 18,518 | NA | 8,315 | 423,433 |
| 1993 | 277,033 | 28,694 | 65,604 | 23,312 | NA | 5,078 | 399,721 |
| 1994 | 150,039 | 2,369 | 52,526 | 10,313 | NA | 1,515 | 216,762 |
| 1995 | 81,454 | 458 | 21,675 | 13,956 | NA | 5,868 | 123,411 |
| 1996 | 4 | 0 | 2,266 | 10,229 | NA | 4,308 | 16,807 |
| 1997 | 52,748 | 486 | 47,355 | 6,400 | NA | 1,199 | 108,188 |
| 1998 | 2,282 | 1,643 | 55,697 | 4,177 | NA | 1,600 | 65,399 |
| 1999 | 5,307 | 970 | 47,163 | 31,106 | NA | 11,458 | 96,004 |
| 2000 | 63,400 | 100 | 4,468 | 38,038 | NA | 2,396 | 108,402 |
| 2001 | 77,491 | 0 | 6,423 | 40,179 | 6,198 | 930 | 131,221 |
| 2002 | 132,921 | 456 | 36,140 | 32,115 | 77 | 10,893 | 212,602 |
| 2003 | 151,826 | 9,057 | 51,622 | 23,995 | NA | 10,082 | 246,582 |
| 2004 | 174,128 | 12,532 | 61,132 | 42,496 | 26 | 20,000 | 310,314 |
| 2005 | 148,734 | 23,599 | 41,710 | 53,928 | 6,225 | 35,000 | 316,756 |

Troll: Areas 21, 23-27, and 121-127; Net: Areas 21, and 23-27; Sport: Areas 23a, 23b, 24-27

¹ Since 1998, the catch accounting year for troll fisheries was set from October 1-September 30. To make comparisons to previous years more meaningful, the same catch accounting period was applied for years prior to 1998.

² Prior to 1992, catch was not reported as 'inside' or 'outside'. Therefore 'inside' catch for those years represents total tidal sport catch.

³ Including 5,000 First Nations troll catch.

NA=not available; “-“ = not applicable.

Appendix A.5. Johnstone Strait Chinook catches, 1975-2005.

| Year | Johnstone Strait | | | | | |
|------|-------------------------------|--------|----------------|------------------|---------------|--------|
| | Troll ¹ Area 12 | Net | Tidal Sport | Freshwater Sport | First Nations | Total |
| 1975 | 18,065 | 30,295 | NA | NA | NA | 48,360 |
| 1976 | 30,838 | 31,855 | NA | NA | NA | 62,693 |
| 1977 | 26,868 | 49,511 | NA | NA | NA | 76,379 |
| 1978 | 13,052 | 55,148 | NA | NA | NA | 68,200 |
| 1979 | 13,052 | 31,291 | NA | NA | NA | 44,343 |
| 1980 | 11,743 | 30,325 | NA | NA | NA | 42,068 |
| 1981 | 13,035 | 28,620 | NA | NA | NA | 41,655 |
| 1982 | 11,234 | 29,454 | NA | NA | NA | 40,688 |
| 1983 | 14,653 | 28,364 | NA | NA | NA | 43,017 |
| 1984 | 9,260 | 18,361 | NA | NA | NA | 27,621 |
| 1985 | 3,567 | 38,073 | NA | NA | NA | 41,640 |
| 1986 | 3,951 | 17,866 | NA | NA | NA | 21,817 |
| 1987 | 1,780 | 13,863 | NA | NA | NA | 15,643 |
| 1988 | 1,566 | 6,292 | NA | NA | NA | 7,858 |
| 1989 | 1,825 | 29,486 | NA | NA | NA | 31,311 |
| 1990 | 2,298 | 18,433 | NA | NA | NA | 20,731 |
| 1991 | 1,228 | 15,071 | 10,075 | NA | 1,287 | 27,661 |
| 1992 | 2,721 | 9,571 | 14,715 | NA | 29 | 27,036 |
| 1993 | 4,172 | 15,530 | NA | NA | 20 | 19,722 |
| 1994 | 2,231 | 8,991 | NA | NA | 0 | 11,222 |
| 1995 | 4 | 970 | NA | NA | 71 | 1,045 |
| 1996 | 0 | 447 | NA | NA | 107 | 554 |
| 1997 | 1,380 | 819 | NA | NA | 179 | 2,378 |
| 1998 | 990 | 60 | 2,366 | NA | 138 | 3,554 |
| 1999 | 89 | 156 | 7,813 | NA | 469 | 8,527 |
| 2000 | 197 | 220 | 5,719 | NA | 212 | 6,348 |
| 2001 | 500 ² | 200 | 3,759 | NA | 370 | 4,329 |
| 2002 | 100 | 600 | 2,331 | NA | 400 | 3,431 |
| 2003 | 710 | 299 | 7585 | NA | 130 | 8724 |
| 2004 | 630 | 220 | 12,837 | NA | 28 | 13,715 |
| 2005 | 2 | 291 | 12,009 | NA | NA | 12,302 |

Troll: Area 12

Net: Areas 11-13

Sport: Based on July - August creel census in Area 12 and northern half of Area 13

¹ Since 1998, the catch accounting year for troll fisheries was set from October 1-September 30.

To make comparisons to previous years more meaningful, the same catch accounting period was applied for years prior to 1998.

² Preliminary estimate

NA=not available

Appendix A.6. Strait of Georgia/Fraser Chinook catches, 1975-2005.

| Year | Strait of Georgia/Fraser | | | | | |
|------|--------------------------|--------|-------------|-------------------------------|----------------------------|---------|
| | Troll ¹ | Net | Tidal Sport | Freshwater Sport ² | First Nations ³ | Total |
| 1975 | 174,001 | 66,119 | 398,000 | NA | 20,170 | 658,290 |
| 1976 | 200,229 | 73,018 | 490,000 | NA | 19,189 | 782,436 |
| 1977 | 248,082 | 85,222 | 372,000 | NA | 23,310 | 728,614 |
| 1978 | 217,955 | 50,247 | 500,000 | NA | 19,541 | 787,743 |
| 1979 | 255,057 | 49,038 | 350,000 | NA | 14,931 | 669,026 |
| 1980 | 273,077 | 31,161 | 204,100 | NA | 15,252 | 523,590 |
| 1981 | 239,266 | 19,985 | 197,239 | NA | 11,987 | 468,477 |
| 1982 | 179,040 | 22,971 | 124,390 | 96 | 35,687 | 362,184 |
| 1983 | 105,133 | 17,520 | 198,433 | NA | 15,756 | 336,842 |
| 1984 | 90,280 | 19,851 | 369,445 | 7,880 | 22,784 | 510,240 |
| 1985 | 55,888 | 31,006 | 234,838 | 1,874 | 10,895 | 334,501 |
| 1986 | 44,043 | 32,359 | 181,896 | 1,573 | 15,646 | 275,517 |
| 1987 | 38,084 | 13,016 | 121,081 | 4,876 | 14,525 | 191,582 |
| 1988 | 20,224 | 8,373 | 119,117 | 7,546 | 15,589 | 170,849 |
| 1989 | 28,444 | 23,833 | 132,846 | 918 | 5,983 | 192,024 |
| 1990 | 34,304 | 15,298 | 111,914 | 2,341 | 17,948 | 181,805 |
| 1991 | 32,412 | 15,407 | 115,523 | 1,616 | 22,185 | 187,143 |
| 1992 | 37,250 | 9,159 | 116,581 | 1,677 | 20,038 | 184,705 |
| 1993 | 33,293 | 16,153 | 127,576 | 1,930 | 20,597 | 199,549 |
| 1994 | 12,916 | 14,078 | 70,839 | 2,475 | 22,476 | 122,784 |
| 1995 | 138 | 6,263 | 62,173 | 9,158 | 20,790 | 98,522 |
| 1996 | 2 | 9,591 | 89,589 | 6,749 | 17,781 | 123,712 |
| 1997 | 908 | 28,342 | 56,332 | 4,180 | 29,497 | 119,259 |
| 1998 | 105 | 6,779 | 20,923 | 22,709 | 18,926 | 69,442 |
| 1999 | 80 | 3,906 | 43,588 | 10,071 | 28,226 | 85,871 |
| 2000 | 270 | 5,584 | 32,750 | 2,078 | 26,213 | 66,895 |
| 2001 | 0 | 4,301 | 31,259 | 23,729 | 28,460 | 87,749 |
| 2002 | 506 | 8,980 | 52,979 | 21,400 | 27,774 | 111,639 |
| 2003 | 17 | 12,277 | 19,981 | 20,363 | 29,634 | 82,272 |
| 2004 | 17 | 12,318 | 13,475 | 16,885 ⁴ | 41,141 | 89,246 |
| 2005 | 0 | 5,296 | 11,972 | 21,831 | 26,919 | 66,018 |

Troll: Areas 13-18 and 29; Net: Areas 14-19, 28 and 29; Sport: Areas 13-18, 19a, 28 and 29

¹ Since 1998, the catch accounting year for troll fisheries was set from October 1-September 30.

To make comparisons to previous years more meaningful, the same catch accounting period was applied for years prior to 1998.

² Prior to 1990, catch includes catch from Fraser systems only; catch records not available those years from non-Fraser systems.

³ No catch records are available for non-Fraser catch prior to 1990.

⁴ Underestimate. NA=not available

Appendix A.7. Canada - Strait of Juan de Fuca Chinook catches, 1975-2005.

| Year | Canada - Strait of Juan de Fuca | | | | |
|------|---------------------------------|-------------|-------------------------------|---------------|--------|
| | Net | Tidal Sport | Freshwater Sport ¹ | First Nations | Total |
| 1975 | 9,799 | NA | NA | NA | 9,799 |
| 1976 | 13,004 | NA | NA | NA | 13,004 |
| 1977 | 25,344 | NA | NA | NA | 25,344 |
| 1978 | 9,725 | NA | NA | NA | 9,725 |
| 1979 | 8,665 | NA | NA | NA | 8,665 |
| 1980 | 3,438 | 37,900 | NA | NA | 41,338 |
| 1981 | 9,982 | 29,832 | NA | NA | 39,814 |
| 1982 | 7,072 | 30,646 | NA | NA | 37,718 |
| 1983 | 328 | 30,228 | NA | NA | 30,556 |
| 1984 | 6,237 | 24,353 | NA | NA | 30,590 |
| 1985 | 17,164 | 27,843 | NA | NA | 45,007 |
| 1986 | 17,727 | 34,387 | NA | NA | 52,114 |
| 1987 | 6,782 | 24,878 | NA | NA | 31,660 |
| 1988 | 4,473 | 31,233 | NA | NA | 35,706 |
| 1989 | 21,238 | 32,539 | NA | NA | 53,777 |
| 1990 | 7,405 | 30,127 | NA | 42 | 37,574 |
| 1991 | 8,893 | 19,017 | NA | 250 | 28,160 |
| 1992 | 10,023 | 21,090 | NA | 302 | 31,415 |
| 1993 | 2,287 | 13,967 | NA | 317 | 16,571 |
| 1994 | 8,931 | 14,372 | NA | 600 | 23,903 |
| 1995 | 631 | 14,405 | NA | 751 | 15,787 |
| 1996 | 362 | 19,012 | NA | 20 | 19,394 |
| 1997 | 307 | 17,080 | NA | 42 | 17,429 |
| 1998 | 115 | 9,709 | NA | 1,500 | 11,324 |
| 1999 | 128 | 14,808 | NA | 52 | 14,988 |
| 2000 | 100 | 10,973 | NA | 272 | 11,345 |
| 2001 | 0 | 23,463 | NA | 135 | 23,598 |
| 2002 | 0 | 24,084 | NA | NA | 24,084 |
| 2003 | 292 | 26,630 | NA | NA | 26,922 |
| 2004 | 0 | 40,877 | NA | NA | 40,877 |
| 2005 | 153 | 30,480 | NA | NA | 30,633 |

Net: Area 20

Sport: Areas 19b and 20

¹ While catch records are poor, in-river sport catch is believed to be small

NA=not available

Appendix A.8. Washington - Strait of Juan de Fuca Chinook catches, 1975-2005.

| Year | Washington - Strait of Juan de Fuca | | | |
|------|-------------------------------------|--------|--------|---------|
| | Troll | Net | Sport | Total |
| 1975 | 5,752 | 8,048 | 81,681 | 95,481 |
| 1976 | 10,488 | 6,072 | 75,308 | 91,868 |
| 1977 | 8,915 | 14,930 | 53,238 | 77,083 |
| 1978 | 10,006 | 11,224 | 62,299 | 83,529 |
| 1979 | 7,804 | 10,939 | 67,094 | 85,837 |
| 1980 | 10,682 | 11,320 | 56,415 | 78,417 |
| 1981 | 15,638 | 18,541 | 51,352 | 85,531 |
| 1982 | 19,024 | 22,547 | 29,842 | 71,413 |
| 1983 | 18,489 | 16,141 | 58,060 | 92,690 |
| 1984 | 15,650 | 12,120 | 48,003 | 75,773 |
| 1985 | 11,808 | 12,784 | 44,267 | 68,859 |
| 1986 | 30,000 | 17,000 | 69,000 | 116,000 |
| 1987 | 45,000 | 11,000 | 53,000 | 109,000 |
| 1988 | 49,000 | 10,000 | 39,000 | 98,000 |
| 1989 | 65,000 | 10,000 | 52,000 | 127,000 |
| 1990 | 47,162 | 5,294 | 50,903 | 103,359 |
| 1991 | 37,127 | 3,390 | 39,667 | 80,184 |
| 1992 | 31,452 | 927 | 38,438 | 70,817 |
| 1993 | 9,794 | 1,482 | 32,434 | 43,710 |
| 1994 | 3,346 | 5,864 | 1,661 | 10,871 |
| 1995 | 6,397 | 4,769 | 6,349 | 17,515 |
| 1996 | 9,757 | 604 | 4,825 | 15,186 |
| 1997 | 829 | 492 | 12,238 | 13,559 |
| 1998 | 338 | 265 | 2,159 | 2,762 |
| 1999 | 544 | 589 | 1,990 | 3,123 |
| 2000 | 332 | 640 | 1,670 | 2,642 |
| 2001 | 1,974 | 931 | 4,819 | 7,724 |
| 2002 | 1,783 | 1,076 | 2,028 | 4,887 |
| 2003 | 436 | 908 | 5,290 | 6,634 |
| 2004 | 20,627 | 592 | 4,519 | 25,738 |
| 2005 | 5,344 | 175 | NA | NA |

Troll: Areas 5 and 6C; Area 4B from Jan. 1 - April 30 and Oct. 1 - Dec. 31

Net: Areas 4B, 5, and 6C

Sport: Areas 5 and 6, 4B Neah Bay "add-on" fishery

Appendix A.9. Washington - San Juan Chinook catches, 1975-2005.

| Year | Washington - San Juans | | | |
|------|------------------------|--------|--------|---------|
| | Troll | Net | Sport | Total |
| 1975 | 3 | 90,100 | 31,988 | 122,091 |
| 1976 | 0 | 66,832 | 34,505 | 101,337 |
| 1977 | 62 | 84,316 | 14,049 | 98,427 |
| 1978 | 3 | 87,565 | 15,083 | 102,651 |
| 1979 | 5 | 53,750 | 17,367 | 71,122 |
| 1980 | 0 | 64,338 | 12,231 | 76,569 |
| 1981 | 4 | 50,695 | 9,727 | 60,426 |
| 1982 | 0 | 38,763 | 6,953 | 45,716 |
| 1983 | 2 | 28,497 | 15,166 | 43,665 |
| 1984 | 83 | 33,432 | 25,759 | 59,274 |
| 1985 | 872 | 33,579 | 12,610 | 47,061 |
| 1986 | 0 | 21,000 | 15,000 | 36,000 |
| 1987 | 0 | 29,000 | 14,000 | 43,000 |
| 1988 | 0 | 32,000 | 9,000 | 41,000 |
| 1989 | 1,000 | 16,000 | 9,000 | 26,000 |
| 1990 | 666 | 8,608 | 7,370 | 16,644 |
| 1991 | 135 | 11,753 | 5,115 | 17,003 |
| 1992 | 172 | 14,011 | 6,788 | 20,971 |
| 1993 | 243 | 14,002 | 6,916 | 21,161 |
| 1994 | 73 | 13,908 | 5,795 | 19,776 |
| 1995 | 9 | 5,333 | 7,863 | 13,205 |
| 1996 | 153 | 3,934 | 12,674 | 16,761 |
| 1997 | 29 | 29,593 | 9,155 | 38,777 |
| 1998 | 376 | 3,804 | 3,069 | 7,249 |
| 1999 | 114 | 3 | 3,421 | 3,538 |
| 2000 | 22 | 1,091 | 4,447 | 5,560 |
| 2001 | 0 | 970 | 6,522 | 7,492 |
| 2002 | 0 | 2,231 | 4,823 | 7,054 |
| 2003 | 0 | 4,827 | 3,036 | 7,863 |
| 2004 | 123 | 5,183 | 1,998 | 7,304 |
| 2005 | 0 | 4,306 | NA | NA |

Troll: Areas 6, 6A, 7, and 7A

Net: Areas 6, 6A, 7 and 7A

Sport: Area 7

NA=not available

Appendix A.10. Washington – Other Puget Sound Chinook catches, 1975-2005.

| Year | Washington – Other Puget Sound | | |
|------|--------------------------------|---------|---------|
| | Net | Sport | Total |
| 1975 | 131,982 | 173,086 | 305,068 |
| 1976 | 141,281 | 151,246 | 292,527 |
| 1977 | 145,470 | 97,761 | 243,231 |
| 1978 | 150,298 | 116,979 | 267,277 |
| 1979 | 128,073 | 156,402 | 284,475 |
| 1980 | 171,516 | 142,799 | 314,315 |
| 1981 | 145,152 | 106,048 | 251,200 |
| 1982 | 149,274 | 85,703 | 234,977 |
| 1983 | 134,492 | 123,752 | 258,244 |
| 1984 | 180,248 | 102,740 | 282,988 |
| 1985 | 184,907 | 92,603 | 277,510 |
| 1986 | 153,000 | 88,000 | 241,000 |
| 1987 | 127,000 | 59,000 | 186,000 |
| 1988 | 133,000 | 63,000 | 196,000 |
| 1989 | 156,000 | 75,000 | 231,000 |
| 1990 | 179,593 | 71,000 | 250,593 |
| 1991 | 89,495 | 48,859 | 138,354 |
| 1992 | 63,460 | 51,656 | 115,116 |
| 1993 | 54,968 | 41,034 | 96,002 |
| 1994 | 63,577 | 44,181 | 107,758 |
| 1995 | 63,593 | 61,509 | 125,102 |
| 1996 | 61,658 | 58,538 | 120,196 |
| 1997 | 47,522 | 43,961 | 91,483 |
| 1998 | 50,915 | 30,016 | 80,931 |
| 1999 | 91,947 | 34,116 | 126,063 |
| 2000 | 79,494 | 29,328 | 108,822 |
| 2001 | 123,266 | 40,170 | 163,436 |
| 2002 | 106,409 | 35,836 | 142,245 |
| 2003 | 86,562 | 32,650 | 119,212 |
| 2004 | 69,283 | 25,316 | 94,599 |
| 2005 | 77,569 | NA | NA |

Net: Areas 6B, 6D, 7B, 7C, and 7E; Areas 8-13 (including all sub-areas); Areas 74C – 83F

Sport: Areas 8-13 and all Puget Sound Rivers

NA=not available

Appendix A.11. Washington – Inside Coastal Chinook catches, 1975-2005.

| Year | Washington – Inside Coastal | | |
|------|-----------------------------|--------|--------|
| | Net | Sport | Total |
| 1975 | 34,859 | 1,716 | 36,575 |
| 1976 | 51,995 | 2,219 | 54,214 |
| 1977 | 72,467 | 2,043 | 74,510 |
| 1978 | 32,662 | 3,399 | 36,061 |
| 1979 | 36,501 | 2,199 | 38,700 |
| 1980 | 47,681 | 1,476 | 49,157 |
| 1981 | 36,880 | 786 | 37,666 |
| 1982 | 33,271 | 1,114 | 34,385 |
| 1983 | 16,210 | 1,452 | 17,662 |
| 1984 | 16,239 | 1,319 | 17,558 |
| 1985 | 25,162 | 1,955 | 27,117 |
| 1986 | 29,000 | 3,000 | 32,000 |
| 1987 | 51,000 | 3,000 | 54,000 |
| 1988 | 74,000 | 7,000 | 81,000 |
| 1989 | 85,000 | 6,000 | 91,000 |
| 1990 | 57,770 | 5,000 | 62,770 |
| 1991 | 54,397 | 6,070 | 60,467 |
| 1992 | 64,223 | 6,577 | 70,800 |
| 1993 | 59,285 | 9,180 | 68,465 |
| 1994 | 46,059 | 7,454 | 53,513 |
| 1995 | 46,490 | 9,881 | 56,371 |
| 1996 | 55,408 | 12,059 | 67,467 |
| 1997 | 28,269 | 6,619 | 34,888 |
| 1998 | 20,266 | 6,569 | 26,835 |
| 1999 | 11,400 | 3,165 | 13,565 |
| 2000 | 15,660 | 3,179 | 18,839 |
| 2001 | 19,480 | 8,645 | 28,125 |
| 2002 | 25,260 | 3,524 | 28,784 |
| 2003 | 19,979 | 6,044 | 26,023 |
| 2004 | 28,363 | 12,006 | 40,369 |
| 2005 | 24,889 | NA | NA |

Net: Areas 2A - 2M; Areas 72B - 73H

Sport: All coastal rivers, Area 2.1, and Area 2.2 (when Area 2 is open)

NA=not available

Appendix A.12. Washington/Oregon North of Cape Falcon Chinook catches, 1975-2005.

| Year | Washington/Oregon North of Cape Falcon | | | |
|------|--|-------|---------|---------|
| | Troll | Net | Sport | Total |
| 1975 | 268,971 | 1,212 | 265,785 | 535,968 |
| 1976 | 371,239 | 203 | 215,319 | 586,761 |
| 1977 | 244,491 | 4 | 197,563 | 442,058 |
| 1978 | 150,673 | 4 | 104,306 | 254,983 |
| 1979 | 133,035 | 3 | 84,977 | 218,015 |
| 1980 | 125,709 | 1,215 | 59,099 | 186,023 |
| 1981 | 109,519 | 209 | 96,151 | 205,879 |
| 1982 | 154,720 | 267 | 114,952 | 269,939 |
| 1983 | 63,584 | 62 | 51,789 | 115,435 |
| 1984 | 15,392 | 0 | 6,980 | 22,372 |
| 1985 | 55,408 | 493 | 30,189 | 86,090 |
| 1986 | 52,000 | 0 | 23,000 | 75,000 |
| 1987 | 81,000 | 4,000 | 44,000 | 129,000 |
| 1988 | 108,000 | 3,000 | 19,000 | 130,000 |
| 1989 | 74,600 | 1,000 | 20,900 | 96,500 |
| 1990 | 65,800 | 0 | 32,900 | 98,700 |
| 1991 | 51,600 | 0 | 13,300 | 64,900 |
| 1992 | 69,000 | 0 | 18,900 | 87,900 |
| 1993 | 55,900 | 0 | 13,600 | 69,500 |
| 1994 | 4,500 | 0 | 0 | 4,500 |
| 1995 | 9,500 | 0 | 600 | 10,100 |
| 1996 | 12,300 | 0 | 200 | 12,500 |
| 1997 | 20,500 | 0 | 4,100 | 24,600 |
| 1998 | 20,615 | 0 | 2,292 | 22,907 |
| 1999 | 44,923 | 0 | 10,821 | 55,744 |
| 2000 | 20,152 | 0 | 9,242 | 29,394 |
| 2001 | 54,163 | 0 | 25,592 | 79,755 |
| 2002 | 106,412 | 0 | 60,575 | 166,987 |
| 2003 | 101,683 | 0 | 36,513 | 138,196 |
| 2004 | 88,175 | 0 | 27,090 | 115,265 |
| 2005 | 87,126 | 0 | 40,011 | 127,137 |

Troll: OR Area 2; WA Areas 1, 2, 3 and 4: Area 4B from May 1 through Sept. 30 (during PFMC management)

Net: WA Areas 1, 2, 3, 4, 4A

Sport: OR Area 2; WA Areas 1, 1.1, 1.2, 2, 3, 4 and 2.2 (when Area 2 is open)

Appendix A.13. Columbia River Chinook catches, 1975-2005.

| Year | Columbia River ¹ | | | |
|------|-----------------------------|--------------------------|---------|---------|
| | Net | Ceremonial & Subsistence | Sport | Total |
| 1975 | 323,000 | | 34,870 | 357,870 |
| 1976 | 288,400 | | 42,527 | 330,927 |
| 1977 | 255,600 | | 58,838 | 314,438 |
| 1978 | 189,100 | | 56,582 | 245,682 |
| 1979 | 169,691 | 7,865 | 38,700 | 216,256 |
| 1980 | 146,356 | 2,938 | 14,860 | 164,154 |
| 1981 | 86,554 | 3,647 | 20,882 | 111,083 |
| 1982 | 151,479 | 8,548 | 30,984 | 191,011 |
| 1983 | 55,946 | 9,669 | 22,709 | 88,324 |
| 1984 | 124,683 | 10,029 | 43,498 | 178,210 |
| 1985 | 143,371 | 9,227 | 45,104 | 197,702 |
| 1986 | 271,813 | 13,386 | 56,168 | 341,367 |
| 1987 | 480,361 | 11,846 | 105,103 | 597,310 |
| 1988 | 486,784 | 15,947 | 97,622 | 600,353 |
| 1989 | 273,574 | 14,525 | 87,636 | 375,735 |
| 1990 | 142,506 | 12,343 | 77,412 | 232,260 |
| 1991 | 104,884 | 7,744 | 76,807 | 189,435 |
| 1992 | 53,385 | 6,898 | 53,629 | 113,912 |
| 1993 | 50,258 | 9,898 | 59,489 | 119,644 |
| 1994 | 31,258 | 8,238 | 27,852 | 67,348 |
| 1995 | 30,370 | 13,294 | 35,321 | 78,985 |
| 1996 | 59,577 | 25,322 | 30,598 | 115,497 |
| 1997 | 52,163 | 32,677 | 44,720 | 129,560 |
| 1998 | 35,033 | 19,482 | 33,402 | 87,917 |
| 1999 | 53,106 | 36,587 | 43,803 | 133,496 |
| 2000 | 59,162 | 24,086 | 49,070 | 132,318 |
| 2001 | 132,284 | 64,985 | 129,219 | 326,488 |
| 2002 | 185,072 | 51,926 | 126,006 | 363,004 |
| 2003 | 178,736 | 44,362 | 127,589 | 350,687 |
| 2004 | 201,432 | 27,890 | 125,950 | 355,272 |
| 2005 | 151,846 | 30,044 | 81,725 | 263,615 |

¹ The historical time series of catches in this year's report has changed from last year's report.

Appendix A.14. Oregon Chinook catches, 1975-2005.

| Year | Oregon | | |
|------|--------|--------|--------|
| | Troll | Sport | Total |
| 1975 | 300 | 19,000 | 19,300 |
| 1976 | 1,000 | 21,000 | 22,000 |
| 1977 | 3,000 | 34,000 | 37,000 |
| 1978 | 1,000 | 37,000 | 38,000 |
| 1979 | 800 | 31,000 | 31,800 |
| 1980 | 300 | 22,000 | 22,300 |
| 1981 | 300 | 28,000 | 28,300 |
| 1982 | 500 | 23,000 | 23,500 |
| 1983 | 700 | 19,000 | 19,700 |
| 1984 | 1,088 | 27,000 | 28,088 |
| 1985 | 1,700 | 25,000 | 26,700 |
| 1986 | 1,900 | 33,000 | 34,900 |
| 1987 | 3,600 | 46,000 | 49,600 |
| 1988 | 4,800 | 49,000 | 53,800 |
| 1989 | 4,500 | 45,000 | 49,500 |
| 1990 | 0 | 38,000 | 38,000 |
| 1991 | 0 | 44,500 | 44,500 |
| 1992 | 384 | 39,000 | 39,384 |
| 1993 | 649 | 52,000 | 52,649 |
| 1994 | 371 | 33,590 | 33,961 |
| 1995 | 206 | 48,366 | 48,572 |
| 1996 | 989 | 56,202 | 57,191 |
| 1997 | 513 | 37,659 | 38,172 |
| 1998 | 858 | 37,990 | 38,848 |
| 1999 | 1,233 | 30,735 | 31,968 |
| 2000 | 1,860 | 33,262 | 35,122 |
| 2001 | 1,184 | 54,988 | 56,172 |
| 2002 | 1,633 | 61,085 | 62,718 |
| 2003 | 1,459 | 67,939 | 69,398 |
| 2004 | 2,258 | 71,726 | 73,984 |
| 2005 | 1,956 | NA | NA |

Troll: Late season off Elk River mouth.

Sport: Estuary and inland.

NA = not available.

Appendix B. Escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005.

| | <u>Page</u> |
|---|--------------------|
| Appendix B.1. Southeast Alaska and Transboundary river escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005. | 102 |
| Appendix B.2. Canadian escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005. | 104 |
| Appendix B.3. Puget Sound escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005. | 106 |
| Appendix B.4. Washington Coast escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005. | 107 |
| Appendix B.5. Columbia River escapements and terminal runs of PSC CTC wild Chinook escapement indicator stocks, 1975-2005. | 108 |
| Appendix B.6. Oregon Coastal escapements and terminal runs of PSC Chinook Technical Committee wild Chinook salmon escapement indicator stocks, 1975-2005. | 109 |

Appendix B.1. Southeast Alaska and Transboundary river escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005.

| Southeast Alaska | | | | | | |
|------------------|-------|--------|--------|--------|---------|-------|
| Year | Situk | | King | | Blossom | Keta |
| | esc. | t. run | Salmon | Andrew | Index | Index |
| | | | esc. | esc. | esc. | esc. |
| 1975 | | | 62 | 520 | 146 | 203 |
| 1976 | 1,421 | 3,184 | 96 | 404 | 68 | 84 |
| 1977 | 1,732 | 2,981 | 199 | 456 | 112 | 230 |
| 1978 | 808 | 1,745 | 84 | 388 | 143 | 392 |
| 1979 | 1,284 | 3,089 | 113 | 327 | 54 | 426 |
| 1980 | 905 | 2,504 | 104 | 282 | 89 | 192 |
| 1981 | 702 | 1,857 | 139 | 536 | 159 | 329 |
| 1982 | 434 | 949 | 354 | 672 | 345 | 754 |
| 1983 | 592 | 1,290 | 245 | 366 | 589 | 822 |
| 1984 | 1,726 | 2,948 | 265 | 389 | 508 | 610 |
| 1985 | 1,521 | 2,916 | 175 | 640 | 709 | 624 |
| 1986 | 2,067 | 2,873 | 255 | 1,416 | 1,278 | 690 |
| 1987 | 1,379 | 2,874 | 196 | 1,576 | 1,349 | 768 |
| 1988 | 868 | 1,596 | 208 | 1,128 | 384 | 575 |
| 1989 | 637 | 1,377 | 240 | 1,060 | 344 | 1,155 |
| 1990 | 628 | 1,643 | 179 | 1,328 | 257 | 606 |
| 1991 | 889 | 2,095 | 134 | 800 | 239 | 272 |
| 1992 | 1,595 | 3,819 | 99 | 1,556 | 150 | 217 |
| 1993 | 952 | 2,558 | 259 | 2,120 | 303 | 362 |
| 1994 | 1,271 | 6,085 | 207 | 1,144 | 161 | 306 |
| 1995 | 4,330 | 14,987 | 144 | 686 | 217 | 175 |
| 1996 | 1,800 | 8,100 | 284 | 670 | 220 | 297 |
| 1997 | 1,878 | 6,601 | 357 | 586 | 132 | 246 |
| 1998 | 924 | 5,420 | 132 | 974 | 91 | 180 |
| 1999 | 1,461 | 7,208 | 300 | 1,210 | 212 | 276 |
| 2000 | 1,785 | 4,941 | 137 | 1,380 | 231 | 300 |
| 2001 | 656 | 2,317 | 147 | 2,108 | 204 | 343 |
| 2002 | 1,000 | 3,017 | 153 | 1,752 | 224 | 411 |
| 2003 | 2,117 | 6,280 | 117 | 1,190 | 203 | 322 |
| 2004 | 748 | 3,275 | 134 | 3,068 | 333 | 376 |
| 2005 | 613 | 1,171 | 141 | 2,030 | 445 | 497 |
| Goal Lower | 500 | | 120 | 650 | 250 | 250 |
| Goal Upper | 1,000 | | 240 | 1,500 | 500 | 500 |

(continued)

Appendix B.1. (Page 2 of 2).

| Transboundary Rivers | | | | | | |
|----------------------|----------------------------------|--------------|-----------------|-----------------------|----------------------------|-----------------|
| Year | Alsek (Klukshu) Index esc. | Taku esc. | Stikine esc. | Unuk Index esc. | Chickamin Index esc. | Chilkat esc. |
| 1975 | | 12,920 | 7,571 | | 370 | |
| 1976 | 1,064 | 24,582 | 5,723 | | 157 | |
| 1977 | 2,698 | 29,496 | 11,445 | 974 | 363 | |
| 1978 | 2,530 | 17,124 | 6,835 | 1,106 | 308 | |
| 1979 | 3,104 | 21,617 | 12,610 | 576 | 239 | |
| 1980 | 2,487 | 39,239 | 30,573 | 1,016 | 445 | |
| 1981 | 1,963 | 49,559 | 36,057 | 731 | 384 | |
| 1982 | 1,969 | 23,847 | 40,488 | 1,351 | 571 | |
| 1983 | 2,237 | 9,795 | 6,424 | 1,125 | 599 | |
| 1984 | 1,572 | 20,778 | 13,995 | 1,837 | 1,102 | |
| 1985 | 1,283 | 35,916 | 16,037 | 1,184 | 956 | |
| 1986 | 2,607 | 38,110 | 14,889 | 2,126 | 1,745 | |
| 1987 | 2,491 | 28,935 | 24,632 | 1,973 | 975 | |
| 1988 | 1,994 | 44,524 | 37,554 | 1,746 | 786 | |
| 1989 | 2,202 | 40,329 | 24,282 | 1,149 | 934 | |
| 1990 | 1,698 | 52,143 | 22,619 | 591 | 564 | |
| 1991 | 2,223 | 51,645 | 23,206 | 655 | 487 | 5,897 |
| 1992 | 1,243 | 55,889 | 34,129 | 874 | 346 | 5,284 |
| 1993 | 3,221 | 66,125 | 58,962 | 1,068 | 389 | 4,472 |
| 1994 | 3,620 | 48,368 | 33,094 | 711 | 388 | 6,795 |
| 1995 | 5,397 | 33,805 | 16,784 | 722 | 356 | 3,790 |
| 1996 | 3,382 | 79,019 | 28,949 | 1,167 | 422 | 4,920 |
| 1997 | 2,829 | 114,938 | 26,996 | 636 | 272 | 8,100 |
| 1998 | 1,347 | 31,039 | 25,968 | 840 | 391 | 3,675 |
| 1999 | 2,166 | 19,734 | 19,947 | 680 | 492 | 2,271 |
| 2000 | 1,321 | 30,529 | 27,531 | 1,341 | 801 | 2,035 |
| 2001 | 1,738 | 42,980 | 63,523 | 2,019 | 1,010 | 4,517 |
| 2002 | 2,141 | 52,409 | 50,875 | 897 | 1,013 | 4,051 |
| 2003 | 1,661 | 36,435 | 46,824 | 1,121 | 964 | 5,657 |
| 2004 | 2,455 | 68,199 | 48,900 | 1,008 | 798 | 3,422 |
| 2005 | 963 | 39,007 | 44,033 | 929 | 924 | 3,366 |
| Goal Lower | 1,100 | 30,000 | 14,000 | 650 | 450 | 1,750 |
| Goal Upper | 2,300 | 55,000 | 28,000 | 1,400 | 900 | 3,500 |

Appendix B.2. Canadian escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005.

| Year | Northern B.C. | | | | | | | | |
|------|--------------------------|-----------------------------|------------|--------|------------------|---------|-------------------------|---------------------------|---------------------------|
| | Area 1 Yakoun esc. | Area 3 ¹ Nass | | | Area 4 Skeena | | Area 8 Dean Index | Area 9 Rivers Inlet | Area 10 Smith Inlet |
| | | Above GW ¹ | Total esc. | t. run | esc. | t. run | | | |
| 1975 | 1,500 | | 14,895 | 17,874 | 20,319 | | | 3,280 | 960 |
| 1976 | 700 | | 13,819 | 16,583 | 13,078 | | | 1,640 | 1,000 |
| 1977 | 800 | 13,688 | 14,288 | 18,410 | 29,018 | 39,606 | | 2,225 | 1,050 |
| 1978 | 600 | 15,485 | 16,885 | 21,807 | 22,661 | 35,055 | 3,500 | 2,800 | 2,100 |
| 1979 | 400 | 11,253 | 12,783 | 16,229 | 18,488 | 28,166 | 4,000 | 2,150 | 500 |
| 1980 | 600 | 13,476 | 14,855 | 18,744 | 23,429 | 38,626 | 2,000 | 2,325 | 1,200 |
| 1981 | 750 | 12,625 | 13,925 | 17,606 | 24,523 | 42,018 | 3,500 | 3,175 | 1,020 |
| 1982 | 1,400 | 7,959 | 10,359 | 13,287 | 17,092 | 35,185 | | 2,250 | 1,500 |
| 1983 | 600 | 13,252 | 16,301 | 20,516 | 23,562 | 39,510 | 500 | 3,320 | 1,050 |
| 1984 | 300 | 20,967 | 24,967 | 31,408 | 37,598 | 53,516 | 4,500 | 1,400 | 770 |
| 1985 | 1,500 | 17,782 | 19,694 | 24,768 | 53,599 | 76,544 | 4,000 | 3,371 | 230 |
| 1986 | 500 | 36,523 | 38,123 | 47,967 | 59,968 | 87,566 | 3,300 | 7,623 | 532 |
| 1987 | 2,000 | 19,540 | 20,986 | 26,568 | 59,120 | 76,349 | 1,144 | 5,239 | 1,050 |
| 1988 | 2,000 | 15,345 | 16,715 | 21,094 | 68,705 | 102,563 | 1,300 | 4,429 | 1,050 |
| 1989 | 2,800 | 28,133 | 29,175 | 36,594 | 57,202 | 83,439 | 2,300 | 3,265 | 225 |
| 1990 | 2,000 | 24,051 | 26,551 | 33,384 | 55,976 | 89,447 | 2,000 | 4,039 | 510 |
| 1991 | 1,900 | 6,907 | 8,259 | 13,136 | 52,753 | 79,343 | 2,400 | 6,635 | 500 |
| 1992 | 2,000 | 16,808 | 17,408 | 25,405 | 63,392 | 92,184 | 3,000 | 7,500 | 500 |
| 1993 | 1,000 | 24,814 | 26,508 | 36,678 | 66,977 | 96,018 | 700 | 10,000 | 500 |
| 1994 | 2,000 | 21,169 | 25,689 | 32,864 | 48,712 | 68,127 | 1,300 | 3,500 | 700 |
| 1995 | 1,500 | 7,844 | 8,776 | 16,187 | 34,390 | 48,351 | 1,100 | 3,196 | 400 |
| 1996 | 3,000 | 21,842 | 22,712 | 30,889 | 73,684 | 96,453 | 2,000 | 3,000 | 250 |
| 1997 | 2,500 | 18,702 | 20,584 | 27,658 | 42,539 | 65,350 | 1,400 | 4,980 | 100 |
| 1998 | 3,000 | 23,213 | 25,361 | 34,922 | 46,744 | 65,167 | 3,000 | 5,367 | 1,100 |
| 1999 | 3,200 | 11,544 | 13,118 | 22,310 | 43,775 | 70,993 | 1,800 | 2,739 | 500 |
| 2000 | 3,600 | 18,912 | 20,565 | 31,159 | 51,804 | 77,320 | 1,200 | 6,700 | 500 |
| 2001 | 3,500 | 29,687 | 31,915 | 44,595 | 81,504 | 112,346 | 3,795 | 5,062 | 300 |
| 2002 | 3,000 | 13,773 | 15,382 | 21,528 | 44,771 | 63,069 | 3,731 | 5,031 | - ² |
| 2003 | 4,000 | 26,940 | 28,330 | 36,503 | 56,758 | 82,410 | 3,700 | 1,900 | - ² |
| 2004 | 4,500 | 15,912 | 18,185 | 25,137 | 44,243 | 61,065 | 3,500 | 3,950 | - ² |
| 2005 | 5,000 | 14,901 | 16,595 | 24,067 | 29,067 | 39,278 | 2,200 | 5,585 | - ² |

¹ GW refers to Gitwinksihlkw, the location of the lower fish wheels on the Nass River used to capture Chinook for the mark-recapture estimate.

² The Docee River was dropped as an escapement indicator due to an inability to obtain reliable escapement estimates.

Appendix B.2. (Page 2 of 2).

| Year | Southern B.C. | | | | Fraser River | | | | | |
|---------|---|------------------------------------|--------|------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-----------------------------|-------------------------|
| | W. Coast Vancouver Island esc. | Lower Georgia Strait esc. | t. run | Upper Georgia Strait esc. | Fraser Spring Age 1.2 esc. | Fraser Spring Age 1.3 esc. | Fraser Summer Age 0.3 esc. | Fraser Summer Age 1.3 esc. | Fraser Spr/sum t. run | Harrison esc. t. run |
| 1975 | 800 | 5,475 | 6,390 | | 7,179 | 8,184 | 26,875 | 16,875 | 119,081 | |
| 1976 | 1,075 | 4,340 | 5,390 | | 4,600 | 10,307 | 4,925 | 13,630 | 98,691 | |
| 1977 | 1,835 | 6,530 | 7,590 | 3,880 | 3,675 | 13,261 | 19,600 | 17,240 | 132,553 | |
| 1978 | 2,750 | 6,495 | 7,035 | 6,150 | 4,305 | 15,725 | 16,700 | 19,200 | 109,119 | |
| 1979 | 2,048 | 10,686 | 11,209 | 4,127 | 2,770 | 14,985 | 18,275 | 10,205 | 101,252 | |
| 1980 | 5,974 | 8,819 | 10,519 | 1,367 | 6,255 | 16,521 | 8,350 | 13,625 | 71,504 | |
| 1981 | 5,050 | 6,007 | 7,607 | 1,945 | 2,975 | 12,274 | 13,120 | 12,202 | 62,668 | |
| 1982 | 6,812 | 6,186 | 6,657 | 3,260 | 5,510 | 15,010 | 6,850 | 15,088 | 85,140 | |
| 1983 | 2,700 | 6,582 | 6,862 | 3,770 | 2,641 | 24,225 | 9,500 | 16,604 | 72,526 | |
| 1984 | 3,862 | 8,456 | 8,861 | 4,600 | 6,380 | 30,370 | 15,522 | 13,595 | 95,681 | 120,837 131,740 |
| 1985 | 3,700 | 4,589 | 5,242 | 4,600 | 9,477 | 43,168 | 20,375 | 19,099 | 121,941 | 174,778 181,367 |
| 1986 | 2,760 | 3,105 | 3,776 | 1,630 | 10,275 | 48,446 | 22,460 | 32,505 | 144,617 | 162,596 177,662 |
| 1987 | 2,570 | 3,276 | 3,781 | 6,450 | 5,049 | 48,271 | 22,404 | 27,646 | 128,699 | 79,038 81,799 |
| 1988 | 4,560 | 7,957 | 8,638 | 3,300 | 4,003 | 41,783 | 29,567 | 32,066 | 129,587 | 35,116 38,285 |
| 1989 | 6,220 | 7,087 | 8,142 | 5,550 | 6,126 | 31,994 | 24,200 | 16,200 | 106,843 | 74,685 76,294 |
| 1990 | 3,660 | 7,023 | 7,627 | 2,320 | 3,225 | 41,560 | 25,425 | 33,747 | 135,124 | 177,375 180,837 |
| 1991 | 5,060 | 8,343 | 8,613 | 3,340 | 3,495 | 27,296 | 26,250 | 28,097 | 116,555 | 90,638 93,363 |
| 1992 | 4,830 | 11,377 | 11,637 | 5,268 | 5,937 | 33,038 | 32,200 | 38,011 | 130,249 | 130,411 132,042 |
| 1993 | 4,530 | 8,418 | 8,713 | 1,574 | 7,870 | 32,796 | 13,300 | 21,385 | 110,237 | 118,998 120,600 |
| 1994 | 4,080 | 7,463 | 7,808 | 1,237 | 10,696 | 51,655 | 25,350 | 23,657 | 145,303 | 98,334 100,839 |
| 1995 | 3,710 | 18,732 | 19,265 | 4,227 | 9,670 | 45,237 | 20,550 | 26,371 | 134,478 | 28,616 29,840 |
| 1996 | 6,026 | 16,465 | 17,275 | 3,600 | 20,726 | 38,398 | 50,900 | 43,142 | 185,559 | 37,394 38,568 |
| 1997 | 7,197 | 11,742 | 11,933 | 5,266 | 9,878 | 44,373 | 49,250 | 40,882 | 202,795 | 70,514 72,061 |
| 1998 | 11,643 | 8,246 | 9,319 | 10,350 | 3,003 | 37,862 | 68,033 | 36,750 | 169,333 | 188,425 189,103 |
| 1999 | 10,186 | 8,481 | 9,181 | 9,500 | 8,751 | 20,740 | 53,204 | 25,138 | 140,939 | 107,016 107,884 |
| 2000 | 4,675 | 7,933 | 8,500 | 12,850 | 11,731 | 26,773 | 45,161 | 25,869 | 155,209 | 77,035 78,098 |
| 2001 | 2,737 | 5,315 | 8,280 | 9,885 | 10,607 | 31,512 | 74,132 | 33,980 | 177,008 | 73,134 74,419 |
| 2002 | 4,036 | 3,840 | 6,022 | 12,865 | 16,423 | 42,408 | 85,132 | 34,886 | 221,020 | 89,968 91,122 |
| 2003 | 4,456 | 3,310 | 5,970 | 13,978 | 17,137 | 45,441 | 70,164 | 44,451 | 231,689 | 247,121 251,453 |
| 2004 | 8,491 | 2,602 | 4,140 | 13,365 | 12,156 | 31,614 | 53,764 | 30,980 | 194,440 | 135,895 138,890 |
| 2005 | 3,969 | 2,527 | 5,179 | 13,365 | 3,898 | 21,458 | 88,329 | 18,586 | 172,281 | 86,730 92,993 |
| Goal LL | | | | | | | | | | 75,100 |
| Goal UL | | | | | | | | | | 98,500 |

Appendix B.3. Puget Sound escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005.

| Year | Puget Sound | | | | | | | | | | | | | |
|------|---------------|--------|-----------------|--------|---------------|--------|-----------|--------|--------|--------|----------------------|---------|----------------------|--------|
| | Skagit Spring | | Skagit Sum/fall | | Stillaguamish | | Snohomish | | Green | | Nooksack Spring esc. | | Lake Washington Fall | |
| | esc. | t. run | esc. | t. run | esc. | t. run | esc. | t. run | esc. | t. run | N. Fork | S. Fork | esc. | t. run |
| 1975 | 627 | 627 | 11,320 | 24,625 | 1,198 | 1,635 | 4,485 | 6,123 | 3,394 | 6,238 | | | 656 | 881 |
| 1976 | 633 | 633 | 14,120 | 23,306 | 2,140 | 4,002 | 5,315 | 9,889 | 3,140 | 7,732 | | | 719 | 759 |
| 1977 | 520 | 520 | 9,218 | 17,994 | 1,475 | 2,549 | 5,565 | 9,618 | 3,804 | 5,366 | | | 675 | 728 |
| 1978 | 932 | 932 | 13,075 | 20,030 | 1,232 | 1,959 | 7,931 | 12,591 | 3,304 | 4,349 | | | 890 | 1,202 |
| 1979 | 818 | 818 | 13,306 | 21,443 | 1,042 | 2,366 | 5,903 | 12,706 | 9,704 | 10,730 | | | 1,289 | 1,430 |
| 1980 | 1,408 | 1,408 | 20,058 | 28,938 | 821 | 2,647 | 6460 | 16,688 | 7743 | 10,608 | | | 1360 | 1,431 |
| 1981 | 1,045 | 1,045 | 8,283 | 19,675 | 630 | 2,783 | 3368 | 8,968 | 3606 | 4,912 | | | 721 | 792 |
| 1982 | 753 | 753 | 9,910 | 20,722 | 773 | 3,058 | 4379 | 8,470 | 1840 | 3,850 | | | 885 | 1,148 |
| 1983 | 554 | 554 | 8,723 | 14,671 | 387 | 925 | 4549 | 10,386 | 3679 | 13,290 | | | 1332 | 2,124 |
| 1984 | 696 | 696 | 12,628 | 15,005 | 374 | 883 | 3762 | 8,480 | 3353 | 5,381 | 45 | 188 | 1252 | 3,436 |
| 1985 | 2,634 | 2,634 | 16,002 | 25,075 | 1,223 | 2,455 | 4,873 | 9,005 | 2,908 | 7,444 | 258 | 445 | 949 | 2,305 |
| 1986 | 1,922 | 1,922 | 17,908 | 21,585 | 1,277 | 2,416 | 4,534 | 8,267 | 4,792 | 5,784 | 226 | 170 | 1,470 | 2,419 |
| 1987 | 1,745 | 1,745 | 9,409 | 13,037 | 1,321 | 1,906 | 4,689 | 6,670 | 10,338 | 11,724 | 181 | 248 | 2,038 | 4,124 |
| 1988 | 1,743 | 1,743 | 11,468 | 14,647 | 726 | 1,185 | 4,513 | 7,389 | 7,994 | 9,207 | 456 | 233 | 792 | 2,373 |
| 1989 | 1,400 | 1,809 | 6,684 | 12,787 | 811 | 1,642 | 3,138 | 6,142 | 11,512 | 15,000 | 303 | 606 | 1,011 | 1,688 |
| 1990 | 1,511 | 1,546 | 16,792 | 19,172 | 842 | 1,739 | 4,209 | 8,345 | 7,035 | 15,200 | 10 | 142 | 787 | 1,128 |
| 1991 | 1,236 | 1,273 | 5,824 | 8,423 | 1,632 | 2,913 | 2,783 | 4,964 | 10,548 | 14,967 | 108 | 365 | 661 | 1,415 |
| 1992 | 986 | 1,010 | 7,348 | 9,201 | 780 | 1,247 | 2,708 | 4,319 | 5,267 | 9,941 | 498 | 103 | 790 | 1,349 |
| 1993 | 782 | 812 | 5,801 | 6,879 | 928 | 1,299 | 3,866 | 5,602 | 2,476 | 5,202 | 449 | 235 | 245 | 304 |
| 1994 | 470 | 496 | 5,656 | 6,586 | 954 | 1,285 | 3,626 | 4,885 | 4,078 | 7,963 | 45 | 118 | 888 | 891 |
| 1995 | 855 | 887 | 6,985 | 9,209 | 822 | 920 | 3,176 | 5,000 | 7,939 | 9,743 | 230 | 290 | 930 | 944 |
| 1996 | 1,051 | 1,078 | 10,706 | 12,286 | 1,244 | 1,244 | 4,851 | 7,921 | 6,026 | 8,668 | 534 | 203 | 336 | 341 |
| 1997 | 1,041 | 1,064 | 4,951 | 6,134 | 1,156 | 1,167 | 4,292 | 4,334 | 11,800 | 12,097 | 570 | 180 | 294 | 296 |
| 1998 | 1,086 | 1,091 | 14,700 | 14,976 | 1,540 | 1,558 | 6,304 | 6,344 | 9,115 | 10,627 | 368 | 157 | 697 | 697 |
| 1999 | 471 | 476 | 5,002 | 5,249 | 1,098 | 1,101 | 4,799 | 4,817 | 13,173 | 14,595 | 823 | 166 | 778 | 778 |
| 2000 | 1,021 | 1,025 | 17,024 | 17,206 | 1,647 | 1,647 | 6,092 | 8,400 | 10,526 | 16,222 | 1,245 | 284 | 347 | 347 |
| 2001 | 1,856 | 1,866 | 13,868 | 14,081 | 1,312 | 1,351 | 8,164 | 8,395 | 21,402 | 24,594 | 2,209 | 267 | 1,269 | 1,516 |
| 2002 | 1,076 | 1,092 | 19,671 | 19,887 | 1,636 | 1,641 | 7,220 | 7,245 | 14,857 | 16,460 | 3,741 | 289 | 637 | 647 |
| 2003 | 909 | 987 | 9,964 | 10,946 | 1,067 | 1,095 | 6,211 | 6,364 | 10,405 | 12,765 | 2,857 | 204 | 771 | 800 |
| 2004 | 1,622 | 1,622 | 23,750 | 24,241 | 1,506 | 1,531 | 10,606 | 10,780 | 13,991 | 20,631 | 2,064 | 130 | 730 | 773 |
| 2005 | 1,305 | NA | 20,803 | 23,396 | 963 | 991 | 4,484 | 4,611 | 4,089 | 4,708 | 2,047 | 120 | 726 | 786 |

Appendix B.4. Washington Coast escapements and terminal runs of PSC Chinook Technical Committee wild Chinook escapement indicator stocks, 1975-2005.

| Year | Washington Coast | | | | | | | | | | | | | |
|------|-------------------|--------|-----------------|--------|-------------|--------|----------|--------|-----------|--------|----------------|--------|-------------|--------|
| | Quillayute Summer | | Quillayute Fall | | Hoh Spr/Sum | | Hoh Fall | | Hoko Fall | | Queets Spr/Sum | | Queets Fall | |
| | esc. | t. run | esc. | t. run | esc. | t. run | esc. | t. run | esc. | t. run | esc. | t. run | esc. | t. run |
| 1975 | | | | | | | | | | | | | | |
| 1976 | 1,300 | 1,700 | | | 600 | 1,300 | 2,500 | 3,100 | | | 505 | 737 | | |
| 1977 | 3,800 | 5,300 | | | 1,000 | 2,000 | 2,100 | 3,800 | | | 732 | 1,155 | 600 | 1,000 |
| 1978 | 2,300 | 2,700 | | | 1,400 | 2,472 | 1,900 | 2,900 | | | 1,110 | 1,406 | 800 | 1,700 |
| 1979 | 2,100 | 3,900 | | | 1,400 | 2,326 | 1,700 | 2,200 | | | 870 | 1,369 | 1,000 | 1,600 |
| 1980 | 964 | 1,500 | 6,700 | 7,600 | 800 | 1,079 | 2,200 | 2,800 | | | 1,038 | 1,213 | 400 | 1,100 |
| 1981 | 815 | 1,700 | 5,963 | 7,102 | 1,498 | 2,005 | 3,100 | 4,000 | | | 988 | 1,329 | 200 | 600 |
| 1982 | 1,126 | 2,700 | 7,107 | 9,651 | 1,553 | 2,125 | 4,500 | 5,800 | | | 781 | 1,244 | 600 | 900 |
| 1983 | 548 | 1,800 | 3,069 | 5,530 | 1,696 | 2,233 | 2,500 | 3,300 | | | 1,044 | 1,173 | 610 | 669 |
| 1984 | 618 | 1,000 | 9,128 | 10,447 | 1,430 | 2,005 | 1,900 | 2,600 | | | 958 | 1,189 | 5,606 | 13,296 |
| 1985 | 550 | 700 | 6,145 | 8,367 | 978 | 1,353 | 1,725 | 2,720 | | | 677 | 886 | 800 | 850 |
| 1986 | 853 | 1,000 | 10,006 | 13,380 | 1,248 | 1,912 | 4,981 | 6,000 | 801 | 839 | 925 | 1,193 | 2,600 | 3,800 |
| 1987 | 666 | 1,600 | 12,352 | 20,349 | 1,710 | 2,480 | 4,006 | 6,147 | 581 | 606 | 598 | 1,543 | 200 | 600 |
| 1988 | 2,599 | 3,943 | 15,168 | 22,115 | 2,605 | 3,708 | 4,128 | 6,873 | 784 | 821 | 1,765 | 2,267 | 11,656 | 21,639 |
| 1989 | 2,407 | 3,472 | 9,951 | 17,260 | 4,697 | 6,820 | 5,148 | 8,682 | 845 | 862 | 2,568 | 3,954 | 600 | 900 |
| 1990 | 1,483 | 1,840 | 13,711 | 16,914 | 3,886 | 5,294 | 4,236 | 6,327 | 493 | 498 | 1,780 | 2,480 | 4,300 | 8,000 |
| 1991 | 1,188 | 1,500 | 6,292 | 7,631 | 1,078 | 1,693 | 1,420 | 2,628 | 1,008 | 1,024 | 630 | 761 | 610 | 669 |
| 1992 | 1,009 | 1,271 | 6,342 | 7,750 | 1,018 | 1,443 | 4,003 | 5,139 | 741 | 750 | 375 | 505 | 800 | 850 |
| 1993 | 1,292 | 1,531 | 5,254 | 5,735 | 1,411 | 2,065 | 2,280 | 3,951 | 894 | 908 | 713 | 788 | 2,600 | 3,800 |
| 1994 | 974 | 1,187 | 4,932 | 5,692 | 1,699 | 2,372 | 3,967 | 4,322 | 429 | 440 | 705 | 727 | 600 | 900 |
| 1995 | 1,333 | 1,731 | 5,532 | 6,716 | 1,132 | 1,686 | 2,202 | 2,912 | 929 | 949 | 625 | 662 | 610 | 669 |
| 1996 | 1,170 | 1,388 | 7,316 | 9,293 | 1,371 | 2,083 | 3,022 | 4,061 | 1,256 | 1,258 | 776 | 891 | 5,482 | 8,997 |
| 1997 | 890 | 1,177 | 5,405 | 6,047 | 1,826 | 2,582 | 1,773 | 3,034 | 868 | 888 | 540 | 693 | 1,128 | 1,130 |
| 1998 | 1,599 | 1,829 | 6,752 | 7,940 | 1,287 | 1,880 | 4,257 | 5,388 | 1,702 | 1,702 | 492 | 537 | 1,128 | 1,130 |
| 1999 | 713 | 818 | 3,334 | 4,758 | 928 | 1,081 | 1,924 | 2,941 | 1,550 | 1,550 | 373 | 426 | 1,128 | 1,130 |
| 2000 | 989 | 1,149 | 3,730 | 4,794 | 492 | 529 | 1,749 | 2,632 | 730 | 730 | 248 | 250 | 1,128 | 1,130 |
| 2001 | 1,225 | 1,399 | 5,136 | 7,545 | 1,159 | 1,231 | 2,560 | 4,116 | 838 | 838 | 548 | 565 | 1,128 | 1,130 |
| 2002 | 1,002 | 1,100 | 6,067 | 9,512 | 2,464 | 3,375 | 4,415 | 5,716 | 680 | 680 | 738 | 755 | 1,128 | 1,130 |
| 2003 | 1,219 | 1,308 | 7,398 | 9,469 | 1,228 | 1,646 | 1,649 | 2,319 | 1,100 | 1,100 | 189 | 195 | 1,128 | 1,130 |
| 2004 | 1,093 | 1,153 | 3,912 | 6,133 | 1,786 | 2,239 | 3,211 | 4,410 | 1,088 | 1,088 | 604 | 619 | 1,128 | 1,130 |
| 2005 | 876 | 958 | 6,406 | 8,420 | 1,193 | 1,389 | 4,180 | 5,267 | 283 | 283 | 294 | 302 | 1,128 | 1,130 |
| Goal | | | 3,000 | | 900 | | 1,200 | | | | 700 | | 2,500 | |

Appendix B.5. Columbia River escapements and terminal runs of PSC CTC wild Chinook escapement indicator stocks, 1975-2005.

| Year | Columbia Upriver | | Columbia Upriver Summers /1 | | | | | | Columbia Upriver Fall Chinook | | | | | | | | |
|------|------------------|---------|-----------------------------|--------|-------------|--------|--------|--------|-------------------------------|--------|--------------------|-------------------------|--------|------------|---------|---------|--------|
| | Spring | | Mid-Columbia | | Snake River | | Total | | Lewis River /2 | | Deschutes River /3 | | | Brights /4 | | | |
| | esc. | t. run | esc. | t. run | esc. | t. run | esc. | t. run | esc. | t. run | esc. | esc. | t. run | esc. | t. run | | |
| 1975 | | | | | | | | | 13,859 | 13,859 | Mark Recapture | Above Falls Expanded | | 29,600 | 164,509 | | |
| 1976 | | | | | | | | | 3,371 | 3,371 | | | | | 27,700 | 109,726 | |
| 1977 | | | | | | | | | 6,930 | 6,930 | | | | 7,484 | 9,345 | 35,600 | 85,755 |
| 1978 | | | | | | | | | 5,363 | 5,363 | | | | 5,049 | 7,020 | 25,800 | 78,280 |
| 1979 | 31,314 | 32,566 | 17,108 | 18,031 | 2,714 | 1,709 | 19,822 | 19,741 | 8,023 | 8,023 | | 4,091 | 5,683 | 28,700 | 83,517 | | |
| 1980 | 32,775 | 33,876 | 16,583 | 17,494 | 2,688 | 2,919 | 19,271 | 20,413 | 16,394 | 16,856 | | 3,159 | 5,110 | 27,700 | 71,690 | | |
| 1981 | 34,235 | 36,091 | 11,826 | 12,741 | 3,306 | 4,474 | 15,132 | 17,215 | 19,297 | 20,298 | | 4,085 | 5,922 | 18,114 | 60,678 | | |
| 1982 | 39,598 | 42,589 | 8,271 | 9,151 | 4,210 | 4,745 | 12,481 | 13,896 | 8,370 | 10,126 | | 7,406 | 9,422 | 27,226 | 69,578 | | |
| 1983 | 31,559 | 32,962 | 7,705 | 7,932 | 3,895 | 4,576 | 11,600 | 12,508 | 13,540 | 14,489 | | 4,681 | 6,177 | 42,681 | 79,923 | | |
| 1984 | 25,171 | 27,039 | 12,369 | 12,689 | 5,429 | 5,079 | 17,798 | 17,768 | 7,132 | 8,128 | | 4,404 | 5,374 | 45,452 | 126,026 | | |
| 1985 | 32,292 | 33,480 | 12,276 | 13,257 | 5,062 | 3,885 | 17,338 | 17,142 | 7,491 | 8,241 | | 3,785 | 4,592 | 72,758 | 191,808 | | |
| 1986 | 40,550 | 43,113 | 10,640 | 11,361 | 6,154 | 5,824 | 16,794 | 17,185 | 11,983 | 13,504 | | 5,355 | 6,508 | 90,961 | 275,061 | | |
| 1987 | 34,980 | 37,286 | 13,769 | 14,931 | 5,891 | 7,519 | 19,660 | 22,450 | 12,935 | 14,173 | | 6,776 | 8,833 | 121,171 | 411,823 | | |
| 1988 | 32,405 | 34,885 | 12,527 | 13,442 | 6,145 | 8,304 | 18,672 | 21,747 | 12,059 | 13,636 | | 5,982 | 8,373 | 97,781 | 331,542 | | |
| 1989 | 32,346 | 35,045 | 17,071 | 17,179 | 3,169 | 3,397 | 20,240 | 20,577 | 21,199 | 22,813 | | 4,777 | 6,507 | 83,100 | 254,795 | | |
| 1990 | 30,189 | 32,439 | 12,883 | 12,976 | 5,093 | 5,123 | 17,976 | 18,099 | 17,506 | 18,784 | | 2,224 | 3,194 | 48,891 | 150,399 | | |
| 1991 | 19,969 | 21,308 | 9,383 | 9,504 | 3,809 | 3,510 | 13,192 | 13,015 | 9,066 | 10,354 | | 3,678 | 3,832 | 39,625 | 99,454 | | |
| 1992 | 33,479 | 35,670 | 6,133 | 6,200 | 3,014 | 3,125 | 9,147 | 9,325 | 6,307 | 7,129 | | 2,777 | 2,814 | 38,879 | 78,202 | | |
| 1993 | 29,349 | 31,280 | 8,962 | 9,235 | 7,889 | 4,520 | 16,851 | 13,755 | 7,025 | 8,106 | | 8,235 | 8,246 | 41,853 | 94,662 | | |
| 1994 | 9,047 | 9,530 | 11,768 | 11,967 | 795 | 907 | 12,563 | 12,874 | 9,939 | 10,541 | | 5,455 | 5,524 | 66,470 | 127,315 | | |
| 1995 | 4,681 | 4,928 | 9,081 | 9,419 | 692 | 841 | 9,773 | 10,260 | 9,718 | 12,155 | | 7,581 | 7,617 | 53,470 | 98,842 | | |
| 1996 | 18,355 | 19,376 | 7,589 | 7,873 | 2,607 | 2,832 | 10,196 | 10,704 | 13,971 | 13,971 | | 8,759 | 8,837 | 51,973 | 134,356 | | |
| 1997 | 17,080 | 18,312 | 8,362 | 8,508 | 10,709 | 7,536 | 19,071 | 16,043 | 8,670 | 8,670 | | 20,678 | 20,811 | 49,074 | 140,916 | | |
| 1998 | 17,226 | 18,156 | 9,525 | 9,757 | 4,355 | 4,739 | 13,880 | 14,496 | 5,929 | 5,929 | | 10,923 | 11,428 | 40,012 | 130,874 | | |
| 1999 | 11,490 | 12,053 | 16,634 | 17,010 | 3,260 | 3,514 | 19,894 | 20,524 | 3,184 | 3,184 | | 3,997 | 4,370 | 44,867 | 161,436 | | |
| 2000 | 49,408 | 52,616 | 16,901 | 17,092 | 3,933 | 4,017 | 20,834 | 21,109 | 9,820 | 9,820 | | 3,230 | 3,637 | 62,675 | 152,107 | | |
| 2001 | 93,011 | 107,225 | 38,708 | 39,295 | 13,735 | 14,623 | 52,443 | 53,918 | 13,886 | 14,186 | 12,595 | 11,161 | 12,929 | 86,908 | 219,562 | | |
| 2002 | 76,976 | 86,463 | 67,676 | 71,607 | 22,159 | 20,104 | 89,835 | 91,711 | 16,380 | 18,230 | 15,505 | 12,252 | 16,475 | 116,237 | 260,794 | | |
| 2003 | 63,970 | 69,511 | 58,613 | 65,367 | 16,422 | 16,672 | 75,035 | 82,039 | 18,505 | 20,505 | 18,568 | 12,590 | 19,646 | 160,677 | 353,545 | | |
| 2004 | 57,233 | 62,767 | 44,536 | 53,674 | 8,813 | 10,206 | 53,349 | 63,879 | 15,342 | 17,133 | 13,369 | 11,879 | 14,593 | 150,440 | 353,265 | | |
| 2005 | 35,833 | 38,266 | 39,138 | 50,505 | 6,736 | 7,585 | 45,874 | 58,090 | 11,348 | 13,348 | 8,924 | 13,550 | 9,759 | 112,679 | 258,985 | | |
| Goal | | | 17,857 | | | | | | 5,700 | | | | | 40,000 | | | |

1/ Columbia Upriver Summers are a single escapement indicator stock with an agency management goal of 85,000. Mid-Columbia summers and Snake River summers exhibit different life history types. Only Mid-Columbia is included in the model stock. Based on a S-R analysis of model data, the interim goal for Mid-Columbia Summers is 17,857 until better data can be compiled.

2/ This is the number of naturally spawning adult fish in the Lewis River. The terminal run given is the escapement plus the Lewis River sport catch of wild adults.

3/ The first column is based on a mark-recapture project for the entire river. The second column is based on using the ratio of redds above and below Sherar's Falls. The agencies' management goal is 4000.

4/ The CRFMP stated an interim escapement goal of 40,000 natural spawning URBs at McNary Dam, including 38,700 for Hanford Reach and 1,100 Snake River. In 1990, the escapement goal was increased to 45,000 for increased hatchery programs. In 1994, a management goal of 46,000 was established, and in 1995, the management goal was retained while the escapement goal was reduced to 43,500. In 2002, the CRFMP escapement goal of 40,000 was agreed to by the CTC. Escapement numbers given are McNary adult dam count minus adult sport and broodstock above the dam. The terminal run is the Columbia River mouth terminal run of Upriver Brights minus the Deschutes River fall Chinook terminal run.

Appendix B.6. Oregon Coastal escapements and terminal runs of PSC Chinook Technical Committee wild Chinook salmon escapement indicator stocks, 1975-2005.

| Year | Oregon | | | | | | |
|------|---------|--------|--------|--------|---------|--------|------------------|
| | Nehalem | | Siletz | | Siuslaw | | Umpqua River |
| | esc. | t. run | esc. | t. run | esc. | t. run | Redd Count Index |
| 1975 | 5,197 | 5,303 | 2,062 | 2,689 | 4,427 | 4,548 | na |
| 1976 | 9,807 | 9,908 | 1,326 | 2,036 | 7,999 | 8,153 | na |
| 1977 | 11,478 | 12,093 | 3,314 | 3,919 | 9,492 | 10,362 | na |
| 1978 | 12,059 | 12,960 | 2,062 | 3,703 | 5,872 | 6,879 | 400 |
| 1979 | 12,205 | 12,841 | 7,217 | 8,907 | 8,040 | 8,799 | na |
| 1980 | 5,555 | 6,379 | 3,680 | 4,823 | 10,630 | 11,183 | 697 |
| 1981 | 10,752 | 11,272 | 4,435 | 6,755 | 8,724 | 9,342 | 890 |
| 1982 | 5,085 | 5,675 | 3,415 | 4,514 | 10,870 | 11,774 | 1,011 |
| 1983 | 4,431 | 4,892 | 2,136 | 3,152 | 4,186 | 4,885 | 1,628 |
| 1984 | 20,341 | 21,623 | 3,461 | 4,571 | 11,168 | 12,437 | 2,594 |
| 1985 | 18,670 | 19,432 | 6,628 | 7,531 | 14,822 | 15,553 | 2,246 |
| 1986 | 10,389 | 11,873 | 6,748 | 7,639 | 14,844 | 15,775 | 1,573 |
| 1987 | 13,560 | 15,654 | 4,577 | 5,906 | 17,603 | 19,031 | 2,795 |
| 1988 | 14,889 | 17,138 | 7,805 | 8,992 | 41,746 | 43,975 | 3,778 |
| 1989 | 10,389 | 11,903 | 4,401 | 5,644 | 28,279 | 31,065 | 6,162 |
| 1990 | 5,104 | 6,726 | 4,313 | 5,148 | 26,799 | 28,893 | 3,761 |
| 1991 | 5,557 | 7,649 | 5,633 | 6,597 | 26,100 | 29,011 | 6,717 |
| 1992 | 9,060 | 11,780 | 6,044 | 7,217 | 26,090 | 27,958 | 8,149 |
| 1993 | 5,345 | 9,309 | 4,342 | 6,244 | 10,446 | 13,567 | 3,364 |
| 1994 | 6,486 | 9,400 | 10,475 | 11,990 | 23,570 | 25,584 | 7,128 |
| 1995 | 5,194 | 8,797 | 5,164 | 7,626 | 26,715 | 30,216 | 11,388 |
| 1996 | 9,211 | 13,241 | 7,394 | 9,917 | 33,051 | 38,485 | 10,019 |
| 1997 | 10,026 | 13,053 | 3,726 | 5,814 | 22,305 | 26,195 | 7,286 |
| 1998 | 8,245 | 11,134 | 5,516 | 7,247 | 24,708 | 28,907 | 1,104 |
| 1999 | 8,063 | 10,008 | 4,166 | 6,002 | 29,610 | 32,556 | 1,804 |
| 2000 | 5,257 | 7,491 | 4,982 | 7,626 | 12,999 | 16,830 | 3,140 |
| 2001 | 9,459 | 13,412 | 10,582 | 14,159 | 29,748 | 34,400 | 6,510 |
| 2002 | 18,089 | 22,425 | 14,054 | 18,195 | 41,058 | 46,177 | 3,831 |
| 2003 | 10,906 | 15,005 | 11,149 | 15,345 | 56,546 | 63,754 | 8,918 |
| 2004 | 9,975 | NA | 3,902 | NA | 34,427 | NA | 7,487 |
| 2005 | 7,038 | NA | 6,426 | NA | 16,619 | NA | 3,084 |
| Goal | 6,989 | | 2,944 | | 12,925 | | |

Appendix C. Relationship between exploitation rate indicator stocks, escapement indicator stocks, model stocks, and additional management action stocks identified in the PST annex.

| | <u>Page</u> |
|---|--------------------|
| Appendix C.1. Indicator stocks for Southeast Alaska and Transboundary Rivers..... | 112 |
| Appendix C.2. Indicator stocks for Canada. | 113 |
| Appendix C.3. Indicator stocks for Puget Sound..... | 114 |
| Appendix C.4. Indicator stocks for the Washington Coast..... | 115 |
| Appendix C.5. Indicator stocks for Columbia River and Oregon Coast. | 116 |

Appendix C.1. Indicator stocks for Southeast Alaska and Transboundary Rivers.

| Area | Annex Stock Group ¹ | Annex Indicator Stocks | Run Type | Escapement Indicator Stock | Escapement Objective | Model Stock | Escapement Goal in Model | Exploitation Rate Indicator Stock | CWT Acronym |
|----------------------|--------------------------------|------------------------|----------|----------------------------|----------------------|-----------------|--------------------------|---|-------------|
| SEAK/TBR | | | Spring | Taku | 30,000–55,000 | Alaska South SE | 9,110 | NA | AKS |
| | | | | Stikine | 14,000–28,000 | | | NA | |
| Yakutat | | | | Situk | 500–1,000 | | | NA | |
| | | | | Alsek | 1,100–2,300 | | | NA | |
| SEAK Northern Inside | | | | Chilkat | 1,750-3,500 | | | NA | |
| | | | | King Salmon | 120–240 | | | Alaska Spring (Little Port Walter, Neets Bay Hatchery, Whitman Lake Hatchery, Carroll Inlet Releases, Deer Mountain Hatchery, Crystal Lake Hatchery) | |
| SEAK Central Inside | | | | Andrew Creek | 650–1,500 | | | | |
| SEAK Southern Inside | | | | Unuk | 650–1,400 | | | | |
| | | | | Chickamin | 450–900 | | | | |
| | | | | Blossom | 250–500 | | | | |
| | | | | Keta | 250–500 | | | | |

¹ SEAK fisheries will be managed to achieve escapement objectives for Southeast Alaska and Transboundary River Chinook stocks.
NA = not available

Appendix C.2. Indicator stocks for Canada.

| Area | Annex Stock Group | Annex Indicator Stocks | Run Type | Escapement Indicator Stock | Escapement Objective | Model Stock | Escapement Goal in Model | Exploitation Rate Indicator Stock | CWT Acronym |
|-------------------------|-----------------------------------|---|---------------|--|-------------------------------------|----------------------------------|--------------------------|-----------------------------------|-------------|
| NBC-Area 1 | North / Central British Columbia | Yakoun | Summer | Yakoun | Escapement goal range by stock | North / Central BC | 117,500 | Kitsumkalum | KLM |
| NBC-Area 3 | | Nass | Spring/Summer | Nass | | | | | |
| NBC-Area 4 | | Skeena | | Skeena | | | | | |
| CBC-Area 8 | | | Spring | Dean | | | | | |
| CBC-Area 9 | | | Spring/Fall | Rivers Inlet | | | | | |
| WCVI | West Coast Vancouver Island Falls | Artlish, Burman, Gold, Kauok, Tahsis, Tashish, Marble | Fall | WCVI Aggregate (Artlish, Burman, Kauok, Tahsis, Tashish, Marble) | Escapement goal range for aggregate | WCVI Natural | 42,734 | Robertson Creek | RBT |
| | | | | | | WCVI Hatchery | 6,472 | | |
| Upper Strait of Georgia | Upper Strait of Georgia | Klinaklini, Kakweikan, Wakeman, Kingcome, Nimpkish | Summer/ Fall | Upper Strait of Georgia (Klinaklini, Kakweikan, Wakeman, Kingcome, Nimpkish) | Escapement goal range for aggregate | Upper Strait of Georgia | 23,300 | Quinsam | QUI |
| Lower Strait of Georgia | Lower Strait of Georgia | | Summer/ Fall | | | Lower Strait of Georgia Hatchery | 5,318 | Puntledge | PPS |
| | | | | | | | | Big Qualicum | BQR |
| | | Cowichan, Nanaimo | Fall | Lower Strait of Georgia (Cowichan / Nanaimo) | Escapement goal range for aggregate | Lower Strait of Georgia Natural | 21,935 | Cowichan | COW |
| Fraser River | Fraser Early | Upper Fraser Mid Fraser Thompson | Spring | Fraser Spring-run Age 1.2 | Escapement goal range by stock | Fraser Early | 93,700 | NA | |
| | | | | Fraser Spring-run Age 1.3 | | | | | |
| | | | Summer | Fraser Summer-run Age 1.3 | | | | | |
| | | | | Fraser Summer-run Age 0.3 | | | | | |
| | Fraser Late | Harrison River | Fall | Harrison River | 75,100-98,500 | Fraser Late | 75,100 | Chilliwack | CHI |

Appendix C.3. Indicator stocks for Puget Sound.

| Area | Annex Stock Group | Annex Indicator Stocks | Run Type | Escapement Indicator Stock | Escapement Objective | Model Stock | Escapement Goal in Model | Exploitation Rate Indicator Stock | CWT Acronym |
|-------------------------------------|--|------------------------|-----------------|----------------------------|--------------------------------|---------------------------------|--------------------------|--|-------------|
| North/ Central Puget Sound | North Puget Sound Natural Springs | Nooksack | Spring | Nooksack | Escapement goal range by stock | Nooksack Spring | 4,000 | Nooksack Spring Fingerling Nooksack Spring Yearling | NKF NKS |
| | | Skagit | | Skagit spring | | | | Skagit Spring Fingerling Skagit Spring Yearling | SKF SKS |
| | North Puget Sound Natural Summer/Falls | Nooksack | Summer/ Fall | | Escapement goal range by stock | Nooksack Fall | 11,923 | Samish Fall Fingerling | SAM |
| | | Snohomish | | Snohomish | | Snohomish Wild | 5,250 | NA | |
| | | Skagit group | | Skagit sum/fall | | Skagit Wild | 9,778 | Skagit Summer Fingerling | SSF |
| | | Lake Washington | | Lake Washington Falls | | Puget Sound Natural Fingerling | 16,966 | NA | |
| | | Green River | | Green River | | | | | |
| | | Stillaguamish | | Stillaguamish | | Stillaguamish Wild | 2,000 | Stillaguamish Fall Fingerling | STL |
| | | | | | | | | Nisqually Fall Fingerling | NIS |
| | | | | | | | | Univ. of Washington Accelerated Fall | UWA |
| Hood Canal | Not an Annex stock | | Fall | | | | | George Adams Fall Fingerling | GAD |
| South Puget Sound | Not an annex stock | | Fall | | | Puget Sound Hatchery Fingerling | 24,769 | South Puget Sound Fall Fingerling | SPS |
| | | | | | | | | South Puget Sound Fall Yearling | SPY |
| | | | | | | | | Squaxin Pens Fall Yearling | SQP |
| | | | Spring | | | Puget Sound Hatchery Yearling | 9,136 | White River Spring Yearling | WRY |
| | | | | | | | | | |

NA = not available

Appendix C.4. Indicator stocks for the Washington Coast.

| Area | Annex Stock Group | Annex Indicator Stocks | Run Type | Escapement Indicator Stock | Escapement Objective | Model Stock | Escapement Goal in Model | Exploitation Rate Indicator Stock | CWT Acronym |
|---------------------------|----------------------------------|------------------------|-------------------|----------------------------|--------------------------------|-----------------------------|--------------------------|-----------------------------------|-------------|
| WA Coast/ Juan de Fuca | Washington Coastal Fall Naturals | Hoko | Fall | Hoko | | | | Elwha Fall Fingerling | ELW |
| | | | | | | | | Hoko Fall Fingerling | HOK |
| | | Grays Harbor | | Grays Harbor Fall | Escapement goal range by stock | Washington Coastal Wild | 21,500 | NA | |
| | | Queets | | Queets Fall | | | | Sooes Fall Fingerling | SOO |
| | | Hoh | | Hoh Fall | | | | NA | |
| | | Quillayute | | Quillayute Fall | | | | NA | |
| | | Queets | | Queets Fall | | | | Queets Fall Fingerling | QUE |
| | Not an annex stock | | Fall | | | Washington Coastal Hatchery | 6,703 | NA | |
| | Not an annex stock | | Spring | Grays Harbor Spring | | | | NA | |
| | Not an annex stock | | Spring/ Summer | Queets Spring/Summer | | | | NA | |
| | | | | Hoh Spring/Summer | | | | NA | |
| | Not an annex stock | | Summer | Quillayute Summer | | | | NA | |

NA = not available

Appendix C.5. Indicator stocks for Columbia River and Oregon Coast.

| Area | Annex Stock Group | Annex Indicator Stocks | Run Type | Escapement Indicator Stock | Escapement Objective | Model Stock | Escapement Goal in Model | Exploitation Rate Indicator Stock | CWT Acronym |
|--------------------|--|------------------------|----------|--------------------------------|----------------------|----------------------------|--------------------------|-----------------------------------|-------------|
| Columbia River | Not an Annex stock | | Spring | | | Cowlitz Spring Hatchery | 2,500 | NA | |
| | | | | | | Willamette River Hatchery | 13,500 | Willamette Spring | WSH |
| | Columbia River Summers | Mid-Columbia Summers | Summer | Mid Columbia Summer | 17,857 ¹ | Columbia River Summer | 17,857 | Columbia Summers | SUM |
| | Columbia River Falls | | Fall | | | Fall Cowlitz Hat. | 8,800 | Cowlitz Tule | CWF |
| | | | | | | Spring Creek Hatchery | 7,000 | Spring Creek Tule | SPR |
| | | | | | | Lower Bonneville Hatchery | 26,200 | Columbia Lower River Hatchery | LRH |
| | | Upriver Brights | | Columbia Upriver Bright | | Columbia Upriver Brights | 40,000 | Columbia Upriver Bright | URB |
| | | | | | | | | Hanford Wild | HAN |
| | | Deschutes | | Deschutes River Fall | | | | NA | |
| | | | | | | Lyons Ferry | 3,430 | Lyons Ferry | LYF |
| | | | | | | Mid Columbia River Brights | 12,500 | NA | |
| | | Lewis River | | Lewis | 5,700 | Lewis River Wild | 5,700 | Lewis River Wild | LRW |
| North Oregon Coast | Far North Migrating Oregon Coastal Falls | Nehalem | Fall | Nehalem | 6,989 | Oregon Coast | 62,382 | Salmon River | |
| | | Siuslaw | | Siuslaw | 12,925 | | | | |
| | | Siletz | | Siletz | 2,944 | | | | |
| Mid-Oregon Coast | Not an Annex stock | | Fall | Umpqua | | | | NA | |
| | | | | Mid South Oregon Coastal Falls | | | | NA | |

¹ Interim goal for modeling based on stock recruitment analysis of model data.

NA – not available

Appendix D. ISBM indices.

| | <u>Page</u> |
|---|--------------------|
| Appendix D.1. ISBM Indices for Canadian fisheries, from both the CWT-based exploitation rate analysis (1999-2003) and the Chinook model (1999-2005) used to establish the AI for each year. Order of the stock groups correspond to Annex 4, Chapter 3, Attachment IV and V of the PST 1999 Revised Annexes. | 118 |
| Appendix D.2. ISBM Indices for U.S. fisheries, from both the CWT-based exploitation rate analysis (1999-2003) and the Chinook model (1999-2005) used to establish the AI for each year. Order of the stock groups correspond to Annex 4, Chapter 3, Attachment IV and V of the PST 1999 Revised Annexes | 120 |

Appendix D.1. ISBM Indices for Canadian fisheries, from both the CWT-based exploitation rate analysis (2000-2004) and the Chinook model (2000-2006) used to establish the AI for each year. Order of the stock groups correspond to Annex 4, Chapter 3, Attachment IV and V of the PST 1999 Revised Annexes.

| Stock Group | Escapement Indicator Stocks | Canadian ISBM Indices | | | | | | | | | | | |
|--|---|--------------------------|-------|-------|---------------------|-------|-----------------|-----------------|-----------------|-----------------|-----------------|---------------------|-----------------|
| | | CWT Indices ¹ | | | | | Model Indices | | | | | | |
| | | 2000 | 2001 | 2002 | 2003 | 2004 | 2000 CLB0107 | 2001 CLB0107 | 2002 CLB0206 | 2003 CLB0308 | 2004 CLB0404 | 2005 CLB0506 | 2006 CLB0604 |
| Lower Strait of Georgia | Cowichan Nanaimo ⁵ | 0.196 | 0.260 | 0.247 | 0.363 ⁶ | 0.284 | 0.232 | 0.325 | 0.541 | 0.490 | 0.593 | 0.381 ⁸ | 0.590 |
| | | 0.154 | 0.260 | 0.247 | NA ⁷ | NA | 0.113 | 0.246 | 0.190 | 0.498 | 0.695 | | |
| Fraser Late | Harrison River ³ | 0.073 | 0.090 | 0.105 | 0.055 ⁹ | 0.032 | 0.198 | 0.336 | 0.302 | 0.352 | 0.719 | 0.332 | 0.294 |
| North Puget Sound Natural Springs | Nooksack Skagit | 1.176 | 0.040 | 0.023 | 0.046 | NA | 0.156 | 0.241 | 0.195 | 0.251 | 0.273 | 0.314 | 0.993 |
| | | NA | NA | NA | NA | NA | NA | NA | NA | 0.251 | 0.273 | 0.314 | 0.993 |
| Upper Strait of Georgia | Klinaklini, Kakweikan, Wakeman, Kingcome, Nimpkish | 0.123 | 0.040 | 0.063 | 0.006 | 0.018 | 0.118 | 0.314 | 0.272 | 0.649 | 0.971 | 0.649 | 0.584 |
| Fraser Early (spring and summers) | Upper Fraser, Mid Fraser, Thompson | NA | NA | NA | NA | NA | 0.124 | 0.210 | 0.145 | 0.661 | 0.718 | 0.654 | 0.610 |
| West Coast Vancouver Island Falls | WCVI (Artlish, Burman, Kauok, Tahsis, Tashish, Marble) | 0.083 | 0.060 | 0.248 | 0.496 ¹⁰ | 0.488 | 0.327 | 0.244 | 0.342 | 0.744 | 0.927 | 0.728 | 1.082 |
| Puget Sound Natural Summer / Falls | Skagit | NA | NA | NA | NA | NA | 0.119 | 0.217 | 0.172 | 0.436 | 0.438 | 0.465 | 1.092 |
| | Stillaguamish | 0.111 | 0.145 | NA | NA | 0.027 | 0.234 | 0.469 | 0.375 | 0.513 | 0.567 | 0.587 | 1.166 |
| | Snohomish | NA | NA | NA | NA | NA | 0.116 | 0.222 | 0.176 | 0.435 | 0.445 | 0.457 | 1.101 |
| | Lake Washington | NA | NA | NA | NA | NA | 0.202 | 0.355 | 0.275 | 0.508 | 0.446 | 0.497 ¹¹ | 0.898 |
| | Green River | 0.154 | 0.350 | 0.323 | 0.328 | 0.162 | 0.202 | 0.356 | 0.275 | 0.508 | 0.466 | 0.497 ¹¹ | 0.914 |
| North / Central B. C. | Yakoun, Nass, Skeena, Area 8 | NA | NA | NA | NA | NA | 0.254 | 0.613 | 0.584 | 0.689 | 0.804 | 0.680 | 0.626 |
| Washington Coastal Fall Naturals ⁴ | Hoko, Grays Harbor, Queets, Hoh, Quillayute | NA | NA | NA | NA | NA | 0.161 | 0.354 | 0.292 | 0.292 | 0.435 | 0.457 | 0.363 |
| Columbia River Falls ⁴ | Upriver Brights | NA | NA | NA | NA | NA | 0.104 | 0.377 | 0.429 | 0.686 | 0.663 | 0.640 | 0.523 |
| | Deschutes | NA | NA | NA | NA | NA | 0.104 | 0.377 | 0.429 | 0.686 | 0.663 | 0.640 | 0.523 |
| | Lewis ³ | NA | NA | NA | NA | NA | 0.180 | 0.180 | 0.171 | 0.515 | 0.480 | 0.546 | 0.315 |
| Columbia R Summers ⁴ | Mid-Columbia Summers ³ | NA | NA | NA | NA | NA | 0.085 | 0.144 | 0.198 | 0.352 | 0.333 | 0.406 | 0.335 |
| Far North Migrating OR Coastal Falls ⁴ | Nehalem ³ , Siletz ³ , Siuslaw ³ | NA | NA | NA | NA | NA | 0.110 | 0.505 | 0.514 | 0.689 | 0.672 | 0.674 | 0.515 |

- ¹The CWT-based estimates, not the model estimates, are to be used in postseason assessments.
- ²NA means not available because of insufficient data (lack of stock specific tag codes, base period CWT recoveries, etc).
- ³Stock or stock group with an agreed CTC escapement goal.
- ⁴Stock group not in Annex Attachment IV.
- ⁵Indices for this stock are calculated from CWT recoveries for Cowichan; differences between Nanaimo and Cowichan stock indices are due to differences in terminal harvest.
- ⁶An inconsistency was discovered between the approaches used to calculate the model-based and CWT-based indices. The former included harvest rates for terminal sport while the latter did not. Terminal sport harvest rates are now included in the calculation of both indices. Further review is yet required to determine whether the base period terminal sport harvest rates obtained from analyses of Big Qualicum CWT recoveries adequately represent impacts that would have occurred on Cowichan Chinook.
- ⁷Several problems have been identified in the approach previously used to calculate the CWT-based indices for Nanaimo Chinook; indices for this stock will not be reported as their utility is questionable.
- ⁸Although model-based indices were previously calculated separately for Cowichan and Nanaimo Chinook, these did not adequately represent impacts on either LGS stock. This is because the model-based data represent an aggregate of the two stocks and methods do not currently exist to correctly disaggregate these data for calculation of the ISBM values. Until such methods are developed, a single index value only will be reported representing the aggregate.
- ⁹The terminal sport harvest rates for Chilliwack Hatchery Chinook, the indicator stock, were removed from the calculation for the Harrison River naturals this year because sport harvest has been essentially zero on the natural population.
- ¹⁰An inconsistency was discovered between the calculation of the model-based and CWT-based indices. The former included harvest rates for terminal sport while the latter did not. Terminal sport harvest rates are now included in the calculation of both indices. A further review of the indices for WCVI Chinook will be done to determine whether they represent impacts on the WCVI wild aggregate.
- ¹¹For the Canadian ISBM fisheries, both Lake Washington and Green are assumed to have the same distribution and thus the same index value.

Appendix D.2. ISBM Indices for U.S. fisheries, from both the CWT-based exploitation rate analysis (2000-2004) and the Chinook model (1999-2006) used to establish the AI for each year. Order of the stock groups correspond to Annex 4, Chapter 3, Attachment IV and V of the PST 1999 Revised Annexes.

| Stock Group | Escapement Indicator Stocks | US ISBM Indices | | | | | | | | | | | |
|---|--|--------------------------|--------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|--------------------|-----------------|
| | | CWT Indices ¹ | | | | | Model Indices | | | | | | |
| | | 2000 | 2001 | 2002 | 2003 | 2004 | 2000 CLB0107 | 2001 CLB0107 | 2002 CLB0206 | 2003 CLB0308 | 2004 CLB0404 | 2005 CLB0506 | 2006 CLB0604 |
| Washington Coastal Fall Naturals | Hoko | NA | NA | NA ¹ | NA ¹ | NA ¹ | 0.34 | 0.56 | 0.48 | 0.682 | 0.966 | 0.444 | 0.442 |
| | Grays Harbor | 1.630 | 0.860 | 0.540 | 0.150 | 0.530 | 0.430 | 0.450 | 0.840 | 0.494 | 0.573 | 0.222 | 0.544 |
| | Queets | 0.850 | 1.440 | 0.840 | 0.850 | 0.840 | 0.420 | 0.440 | 1.050 | 1.063 | 0.932 | 1.023 | 1.022 |
| | Hoh | 2.750 | 1.660 | 0.950 | 1.340 | 1.220 | 0.730 | 0.760 | 1.260 | 1.208 | 1.214 | 1.499 | 1.493 |
| | Quillayute | 2.470 | 1.480 | 1.420 | 0.990 | 1.150 | 0.720 | 0.750 | 1.310 | 1.292 | 1.139 | 1.133 | 0.673 |
| Columbia River Falls | Upriver Brights | 2.530 | 1.350 | 1.850 | 1.430 | 1.740 | 1.090 | 0.990 | 0.910 | 1.022 | 0.906 | 0.734 | 0.814 |
| | Deschutes | 0.710 | 0.520 | 0.590 | 0.490 | 0.510 | 0.880 | 0.740 | 0.550 | 0.561 | 0.475 | 0.483 | 0.437 |
| | Lewis ⁵ | 0.360 | 0.580 | 0.560 | 1.030 | 0.170 | 0.160 | 1.700 | 0.930 | 0.851 | 1.008 | 1.058 | 1.861 |
| Puget Sound Natural Summer / Falls | Skagit | NA | NA | NA | NA | NA | 0.210 | 0.780 | 0.270 | 0.406 | 0.157 | 0.195 | 0.258 |
| | Stillaguamish | 0.040 | 0.890 | NA | NA | 0.010 | 0.140 | 0.400 | 0.200 | 0.184 | 0.224 | 0.185 | 0.493 |
| | Snohomish | NA | NA | NA | NA | NA | 0.050 | 0.600 | 0.150 | 0.072 | 0.110 | 0.891 | 0.199 |
| | Lake Washington | NA | NA | NA | NA | NA | 0.480 | 0.590 | 1.250 | 0.768 | 0.411 | 0.373 | 0.613 |
| | Green R | 0.700 | 1.180 | 1.070 | 1.030 | 1.010 | 0.480 | 0.600 | 0.350 | 0.263 | 0.260 | 0.202 | 0.361 |
| Fraser Late | Harrison River ⁵ | 0.130 | 0.310 | 0.410 | 0.640 | 0.320 | 0.390 | 0.620 | 0.720 | 0.981 | 1.058 | 0.670 | 0.787 |
| Columbia R Summers | Mid-Columbia Summers ⁵ | 4.820 | 5.320 | 7.250 | 10.040 | 2.690 | 0.090 | 0.140 | 0.820 | 0.794 | 0.715 | 0.545 | 0.696 |
| Far North Migrating OR Coastal Falls | Nehalem ⁵ | 1.970 | 1.940 | 2.170 | 3.110 | 1.800 | 2.660 | 2.750 | 2.610 | 2.346 | 2.230 | 2.090 | 1.912 |
| | Siletz ⁵ | 1.160 | 1.190 | 1.310 | 1.590 | 2.290 | 1.790 | 1.870 | 1.330 | 1.302 | 1.288 | 1.233 | 1.237 |
| | Siuslaw ⁵ | 2.450 | 2.180 | 2.560 | 3.820 | 1.030 | 0.930 | 0.950 | 3.340 | 2.856 | 2.816 | 2.643 | 1.095 |
| North Puget Sound Natural Springs | Nooksack | 0.000 | 0.040 | NA | NA | NA | 0.200 | 0.010 | 0.000 | 0.121 | 0.974 | 0.222 | 0.121 |
| | Skagit | NA | NA | 1.120 | NA | NA | ID | 0.070 | 0.060 | 0.119 | 0.663 | 0.213 | 0.161 |
| Lower Strait of Georgia ⁴ | Cowichan, | 0.690 | 11.350 | 5.780 | 4.990 | 7.250 | 0.210 | 0.480 | 0.220 | 0.452 | 0.915 | 0.407 ⁸ | 0.271 |
| | Nanaimo | 0.690 | 11.350 | 5.780 | 4.990 | 7.250 | 0.210 | 0.480 | 0.220 | 0.452 | 0.915 | | |
| Upper Strait of Georgia ⁴ | Klinaklini, Kakweikan, Wakeman, Kingcome, Nimpkish | NA | NA | NA | NA | NA | NC | NC | NC | NC | NC | NC | NC |
| Fraser Early (spring and summers) ⁴ | Upper Fraser, Mid Fraser, Thompson | NA | NA | NA | NA | NA | 0.150 | 0.700 | 0.150 | 0.277 | 0.839 | 0.257 | 0.224 |
| West Coast Vancouver Island Falls ⁴ | WCVI (Artlish, Burman, Kauok, Tahsis, Tashish, Marble) | NA | NA | NA | NA | NA | 0.380 | 0.730 | 0.270 | 0.658 | 0.540 | 0.290 | 0.128 |
| North / Central B. C. | Yakoun, Nass, Skeena, Area 8 | NA | NA | NA | NA | NA | NC | NC | NC | NC | NC | NC | NC |

⁸. See the footnote for the corresponding value in the table of indices for the Canadian ISBM fisheries.

Appendix E. Percent distribution of landed catch and total mortality among fisheries and escapement for exploitation rate indicator stocks by calendar year.

These data result from cohort analysis of CWT recoveries for the indicator stocks; data within a row for each calendar year sum to 100%. Some changes are present in these distribution tables compared to those presented in previous reports. There are various reasons for the changes including updates to escapement time series, in the case of some Columbia River stocks. Also, a computational rule used in producing the stock-specific distribution tables determines whether data are reported for any particular calendar year. The rule is that at least three year classes of CWT recoveries (out of four or five) must be available in any calendar year. Lack of CWT releases in recent years for some of the indicators has resulted in no distribution data for 2000-2003. Missing broods are noted in the appropriated tables.

LIST OF APPENDIX E TABLES

| | <u>Page</u> |
|---|--------------------|
| Appendix E.1. Percent distribution of Alaska Spring Chinook reported catch among fisheries and escapement..... | 125 |
| Appendix E.2. Percent distribution of Alaska Spring Chinook total fishing mortalities among fisheries and escapement..... | 126 |
| Appendix E.3. Percent distribution of Kitsumkalum River Summer Chinook reported catch among fisheries and escapement (NA=not available). | 127 |
| Appendix E.4. Percent distribution of Kitsumkalum River Summer Chinook total fishing mortalities among fisheries and escapement (NA=not available)..... | 128 |
| Appendix E.5. Percent distribution of Robertson Creek Fall Chinook reported catch among fisheries and escapement..... | 129 |
| Appendix E.6. Percent distribution of Robertson Creek Fall Chinook total fishing mortalities among fisheries and escapement..... | 130 |
| Appendix E.7. Percent distribution of Quinsam River Fall Chinook reported catch among fisheries and escapement..... | 131 |
| Appendix E.8. Percent distribution of Quinsam River Fall Chinook total fishing mortalities among fisheries and escapement..... | 132 |
| Appendix E.9. Percent distribution of Puntledge River Summer Chinook reported catch among fisheries and escapement. | 133 |
| Appendix E.10. Percent distribution of Puntledge River Summer Chinook total fishing mortalities among fisheries and escapement..... | 134 |
| Appendix E.11. Percent distribution of Big Qualicum River Fall Chinook reported catch among fisheries and escapement. | 135 |
| Appendix E.12. Percent distribution of Big Qualicum Fall Chinook total fishing mortalities among fisheries and escapement..... | 136 |
| Appendix E.13. Percent distribution of Cowichan River Fall Chinook reported catch among fisheries and escapement..... | 137 |
| Appendix E.14. Percent distribution of Cowichan River Fall Chinook total fishing mortalities among fisheries and escapement..... | 137 |
| Appendix E.15. Percent distribution of Chilliwack River Fall Chinook reported catch among fisheries and escapement..... | 138 |

| | | |
|----------------|--|-----|
| Appendix E.16. | Percent distribution of Chilliwack River Fall Chinook total fishing mortalities among fisheries and escapement..... | 139 |
| Appendix E.17. | Percent distribution of Nooksack Spring Fingerling Chinook reported catch among fisheries and escapement. | 140 |
| Appendix E.18. | Percent distribution of Nooksack Spring Fingerling Chinook total fishing mortalities among fisheries and escapement. | 140 |
| Appendix E.19. | Percent distribution of Nooksack Spring Yearling Chinook reported catch among fisheries and escapement. | 141 |
| Appendix E.20. | Percent distribution of Nooksack Spring Yearling Chinook total fishing mortalities among fisheries and escapement. | 141 |
| Appendix E.21. | Percent distribution of Skagit Spring Fingerling Chinook reported catch among fisheries and escapement. | 142 |
| Appendix E.22. | Percent distribution of Skagit Spring Fingerling Chinook total fishing mortalities among fisheries and escapement..... | 142 |
| Appendix E.23. | Percent distribution of Skagit Spring Yearling Chinook reported catch among fisheries and escapement..... | 143 |
| Appendix E.24. | Percent distribution of Skagit Spring Yearling Chinook total fishing mortalities among fisheries and escapement..... | 143 |
| Appendix E.25. | Percent distribution of Samish Fall Fingerling Chinook reported catch among fisheries and escapement..... | 144 |
| Appendix E.26. | Percent distribution of Samish Fall Fingerling Chinook total fishing mortalities among fisheries and escapement..... | 144 |
| Appendix E.27. | Percent distribution of Skagit Summer Fingerling Chinook reported catch among fisheries and escapement. | 145 |
| Appendix E.28. | Percent distribution of Skagit Summer Fingerling Chinook total fishing mortalities among fisheries and escapement. | 145 |
| Appendix E.29. | Percent distribution of Stillaguamish Fall Fingerling Chinook reported catch among fisheries and escapement (NA=not available). | 146 |
| Appendix E.30. | Percent distribution of Stillaguamish Fall Fingerling Chinook total fishing mortalities among fisheries and escapement. | 146 |
| Appendix E.31. | Percent distribution of Nisqually Fall Fingerling Chinook reported catch among fisheries and escapement. | 147 |
| Appendix E.32. | Percent distribution of Nisqually Fall Fingerling Chinook total fishing mortalities among fisheries and escapement..... | 148 |
| Appendix E.33. | Percent distribution of George Adams Fall Fingerling Chinook among fisheries reported catch and escapement..... | 149 |
| Appendix E.34. | Percent distribution of George Adams Fall Fingerling Chinook total fishing among fisheries and escapement..... | 150 |
| Appendix E.35. | Percent distribution of South Puget Sound Fall Fingerling Chinook reported catch among fisheries and escapement. | 151 |
| Appendix E.36. | Percent distribution of South Puget Sound Fall Fingerling Chinook total fishing mortalities among fisheries and escapement. | 152 |
| Appendix E.37. | Percent distribution of South Puget Sound Fall Yearling Chinook reported catch among fisheries and escapement. | 153 |
| Appendix E.38. | Percent distribution of South Puget Sound Fall Yearling Chinook for total fishing mortalities among fisheries and escapement. | 154 |

| | | |
|----------------|--|-----|
| Appendix E.39. | Percent distribution of Squaxin Pens Fall Yearling Chinook reported catch among fisheries and escapement. | 155 |
| Appendix E.40. | Percent distribution of Squaxin Pens Fall Yearling Chinook total fishing mortalities among fisheries and escapement. | 155 |
| Appendix E.41. | Percent distribution of White River Spring Yearling Chinook reported catch among fisheries and escapement. | 156 |
| Appendix E.42. | Percent distribution of White River Spring Yearling Chinook total fishing mortalities among fisheries and escapement. | 157 |
| Appendix E.43. | Percent distribution of Hoko Fall Fingerling Chinook reported catch among fisheries and escapement. | 158 |
| Appendix E.44. | Percent distribution of Hoko Fall Fingerling Chinook total fishing mortalities among fisheries and escapement. | 158 |
| Appendix E.45. | Percent distribution of Sooes Fall Fingerling Chinook reported catch among fisheries and escapement. | 159 |
| Appendix E.46. | Percent distribution of Sooes Fall Fingerling Chinook total fishing mortalities among fisheries and escapement. | 159 |
| Appendix E.47. | Percent distribution of Queets Fall Fingerling Chinook reported catch among fisheries and escapement. | 160 |
| Appendix E.48. | Percent distribution of Queets Fall Fingerling Chinook total fishing mortalities among fisheries and escapement. | 161 |
| Appendix E.49. | Percent distribution of Willamette Spring Chinook reported catch among fisheries and escapement. | 162 |
| Appendix E.50. | Percent distribution of Willamette Spring Chinook total fishing mortalities among fisheries and escapement. | 163 |
| Appendix E.51. | Percent distribution of Columbia Summer Chinook reported catch among fisheries and escapement. | 164 |
| Appendix E.52. | Percent distribution of Columbia Summer Chinook total fishing mortalities among fisheries and escapement. | 165 |
| Appendix E.53. | Percent distribution of Cowlitz Tule Chinook reported catch among fisheries and escapement. | 166 |
| Appendix E.54. | Percent distribution of Cowlitz Tule Chinook total fishing mortalities among fisheries and escapement. | 167 |
| Appendix E.55. | Percent distribution of Spring Creek Tule Chinook reported catch among fisheries and escapement. | 168 |
| Appendix E.56. | Percent distribution of Spring Creek Tule Chinook total fishing mortalities among fisheries and escapement. | 169 |
| Appendix E.57. | Percent distribution of Columbia Lower River Hatchery Chinook reported catch among fisheries and escapement. | 170 |
| Appendix E.58. | Percent distribution of Columbia Lower River Hatchery Chinook total fishing mortalities among fisheries and escapement. | 171 |
| Appendix E.59. | Percent distribution of Upriver Bright Chinook reported catch among fisheries and escapement. | 172 |
| Appendix E.60. | Percent distribution of Upriver Bright Chinook total fishing mortalities among fisheries and escapement. | 173 |
| Appendix E.61. | Percent distribution of Hanford Wild Chinook reported catch among fisheries and escapement. | 174 |

| | | |
|----------------|--|-----|
| Appendix E.62. | Percent distribution of Hanford Wild Chinook total fishing mortalities among fisheries and escapement..... | 174 |
| Appendix E.63. | Percent distribution of Lyons Ferry Chinook reported catch among fisheries and escapement..... | 175 |
| Appendix E.64. | Percent distribution of Lyons Ferry Chinook total fishing mortalities among fisheries and escapement..... | 175 |
| Appendix E.65. | Percent distribution of Lewis River Wild Chinook reported catch among fisheries and escapement..... | 176 |
| Appendix E.66. | Percent distribution of Lewis River Wild Chinook total fishing mortalities among fisheries and escapement..... | 177 |
| Appendix E.67. | Percent distribution of Salmon River Chinook reported catch among fisheries and escapement..... | 178 |
| Appendix E.68. | Percent distribution of Salmon River Chinook total fishing mortalities among fisheries and escapement..... | 179 |

Appendix E.1. Percent distribution of Alaska Spring Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr& Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|--------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1983 | 27.9% | 1.3% | 6.6% | 1.7% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 62.2% |
| 1984 | 23.0% | 2.6% | 13.7% | 0.9% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 59.4% |
| 1985 | 24.1% | 5.6% | 13.7% | 1.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 55.5% |
| 1986 | 25.1% | 5.2% | 11.9% | 0.6% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 57.1% |
| 1987 | 30.9% | 2.8% | 10.6% | 0.4% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 54.8% |
| 1988 | 29.9% | 2.0% | 14.5% | 1.1% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 52.2% |
| 1989 | 25.3% | 9.5% | 10.4% | 0.6% | 0.0% | 0.3% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 53.9% |
| 1990 | 37.0% | 2.4% | 13.4% | 1.7% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 45.4% |
| 1991 | 40.0% | 3.5% | 17.0% | 0.6% | 0.0% | 0.3% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 38.4% |
| 1992 | 26.1% | 6.8% | 20.2% | 0.4% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 46.2% |
| 1993 | 19.8% | 5.9% | 19.2% | 0.1% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 54.6% |
| 1994 | 16.2% | 16.1% | 13.5% | 0.4% | 0.0% | 0.4% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 53.2% |
| 1995 | 26.9% | 14.0% | 17.7% | 0.3% | 0.0% | 0.3% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 40.6% |
| 1996 | 24.4% | 10.3% | 30.4% | 0.0% | 0.0% | 0.3% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 34.4% |
| 1997 | 25.7% | 8.3% | 29.5% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 36.3% |
| 1998 | 28.8% | 10.3% | 25.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 35.5% |
| 1999 | 19.8% | 5.4% | 29.3% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 45.1% |
| 2000 | 22.9% | 5.2% | 24.2% | 0.0% | 0.0% | 0.1% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 47.4% |
| 2001 | 15.7% | 4.3% | 18.1% | 0.2% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 61.6% |
| 2002 | 11.2% | 3.9% | 16.0% | 0.7% | 0.0% | 0.0% | 0.8% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 67.2% |
| 2003 | 16.7% | 1.6% | 15.9% | 0.7% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 64.7% |
| 2004 | 15.5% | 5.4% | 13.8% | 0.4% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 64.4% |
| (83-84) | 25.4% | 2.0% | 10.2% | 1.3% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 60.8% |
| (85-98) | 27.2% | 7.3% | 17.6% | 0.5% | 0.0% | 0.2% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 47.0% |
| (99-04) | 17.0% | 4.3% | 19.5% | 0.3% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 58.4% |

Appendix E.2. Percent distribution of Alaska Spring Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1983 | 34.5% | 1.5% | 11.3% | 1.8% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 50.7% |
| 1984 | 27.5% | 2.6% | 17.9% | 1.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 50.7% |
| 1985 | 27.8% | 10.8% | 15.3% | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 45.2% |
| 1986 | 29.1% | 11.0% | 12.4% | 0.5% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 46.9% |
| 1987 | 40.4% | 5.3% | 9.9% | 0.4% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 43.5% |
| 1988 | 34.4% | 5.8% | 14.2% | 1.2% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 44.2% |
| 1989 | 29.6% | 16.4% | 10.8% | 0.6% | 0.0% | 0.2% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 42.3% |
| 1990 | 43.3% | 6.5% | 13.0% | 1.8% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 35.2% |
| 1991 | 42.0% | 8.6% | 16.1% | 0.6% | 0.0% | 0.3% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 32.2% |
| 1992 | 25.5% | 20.3% | 17.7% | 0.4% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 35.9% |
| 1993 | 23.7% | 9.4% | 19.5% | 0.2% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 46.9% |
| 1994 | 20.3% | 29.1% | 12.4% | 0.4% | 0.0% | 0.3% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 37.4% |
| 1995 | 32.4% | 14.8% | 17.8% | 0.3% | 0.0% | 0.4% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 34.2% |
| 1996 | 27.4% | 11.5% | 30.1% | 0.1% | 0.0% | 0.4% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 30.3% |
| 1997 | 26.9% | 10.9% | 29.5% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 32.4% |
| 1998 | 28.3% | 19.4% | 23.6% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 28.1% |
| 1999 | 22.1% | 8.1% | 31.1% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 38.3% |
| 2000 | 26.5% | 8.0% | 24.7% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 40.5% |
| 2001 | 18.7% | 6.6% | 18.9% | 0.2% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 55.5% |
| 2002 | 13.1% | 7.0% | 17.7% | 0.8% | 0.0% | 0.1% | 1.1% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 60.2% |
| 2003 | 17.9% | 4.3% | 18.5% | 0.7% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 57.9% |
| 2004 | 16.9% | 15.0% | 13.4% | 0.4% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 53.8% |
| (83-84) | 31.0% | 2.0% | 14.6% | 1.4% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 50.7% |
| (85-98) | 30.8% | 12.8% | 17.3% | 0.5% | 0.0% | 0.2% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 38.2% |
| (99-04) | 19.2% | 8.2% | 20.7% | 0.3% | 0.0% | 0.1% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 51.0% |

Appendix E.3. Percent distribution of Kitsumkalum River Summer Chinook reported catch among fisheries and escapement (NA=not available).

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|-----------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1984 | 50.8% | 0.0% | 0.0% | 18.5% | 0.0% | 30.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | NA ¹ |
| 1985 | 26.1% | 0.0% | 1.6% | 7.1% | 0.0% | 13.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 51.6% |
| 1986 | 8.9% | 0.0% | 0.0% | 14.1% | 0.0% | 8.9% | 2.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 65.7% |
| 1987 | 7.4% | 0.0% | 0.0% | 9.1% | 0.0% | 7.8% | 4.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 71.4% |
| 1988 | 17.4% | 0.6% | 1.9% | 3.1% | 0.0% | 23.0% | 7.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 46.6% |
| 1989 | 10.9% | 0.3% | 6.8% | 5.0% | 0.0% | 11.3% | 6.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 59.1% |
| 1990 | 10.7% | 0.0% | 2.8% | 6.6% | 0.3% | 7.1% | 7.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 65.0% |
| 1991 | 14.6% | 0.0% | 3.7% | 8.8% | 0.7% | 16.7% | 13.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 41.8% |
| 1992 | 13.9% | 0.0% | 1.9% | 7.0% | 0.0% | 9.4% | 6.6% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 60.7% |
| 1993 | 10.4% | 0.9% | 2.2% | 10.0% | 0.0% | 18.7% | 4.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 53.5% |
| 1994 | 11.1% | 0.0% | 0.0% | 5.6% | 0.0% | 19.0% | 6.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 57.9% |
| 1995 | 12.1% | 0.0% | 2.7% | 7.1% | 0.0% | 29.1% | 6.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 42.9% |
| 1996 | 8.5% | 0.2% | 6.0% | 0.0% | 0.0% | 18.5% | 5.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 61.9% |
| 1997 | 10.6% | 0.0% | 7.5% | 0.0% | 0.0% | 8.3% | 11.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 62.6% |
| 1998 | 8.6% | 0.0% | 3.1% | 0.0% | 0.0% | 1.2% | 5.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 81.6% |
| 1999 | 14.7% | 0.0% | 9.7% | 0.0% | 0.0% | 0.9% | 6.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 68.0% |
| 2000 | 6.8% | 0.0% | 6.8% | 0.0% | 0.0% | 9.8% | 5.5% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 70.7% |
| 2001 | 7.9% | 0.0% | 5.2% | 0.4% | 0.0% | 6.9% | 10.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 69.3% |
| 2002 | 12.2% | 0.2% | 5.2% | 1.4% | 0.0% | 2.4% | 14.6% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 63.7% |
| 2003 | 13.6% | 0.0% | 1.9% | 5.6% | 0.0% | 0.0% | 10.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 68.6% |
| 2004 | 9.7% | 1.0% | 3.8% | 1.4% | 0.0% | 1.2% | 12.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 69.9% |
| (85-98) | 12.2% | 0.1% | 2.9% | 6.0% | 0.1% | 13.8% | 6.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 58.7% |
| (99-04) | 10.8% | 0.2% | 5.4% | 1.5% | 0.0% | 3.5% | 10.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 68.4% |

1. Values represent estimates of catch distribution only for this year.

Appendix E.4. Percent distribution of Kitsumkalum River Summer Chinook total fishing mortalities among fisheries and escapement (NA=not available).

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|-----------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1984 | 52.6% | 0.0% | 0.0% | 21.1% | 0.0% | 26.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | NA ¹ |
| 1985 | 29.6% | 0.0% | 1.5% | 7.7% | 0.0% | 12.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 48.5% |
| 1986 | 10.2% | 0.0% | 0.0% | 13.9% | 0.0% | 8.8% | 2.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 64.8% |
| 1987 | 12.8% | 0.0% | 2.6% | 9.8% | 0.0% | 7.2% | 5.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 62.3% |
| 1988 | 23.4% | 2.4% | 4.9% | 7.3% | 0.0% | 18.0% | 7.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 36.6% |
| 1989 | 14.3% | 0.6% | 6.9% | 5.3% | 0.0% | 10.6% | 6.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 55.5% |
| 1990 | 11.8% | 0.0% | 3.3% | 7.7% | 0.3% | 6.8% | 7.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 62.1% |
| 1991 | 19.9% | 0.0% | 4.2% | 10.7% | 0.9% | 14.8% | 13.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 36.5% |
| 1992 | 15.4% | 0.0% | 2.0% | 7.9% | 0.0% | 9.1% | 6.9% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 58.3% |
| 1993 | 11.6% | 1.7% | 2.1% | 11.6% | 0.0% | 17.8% | 4.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 50.8% |
| 1994 | 13.3% | 0.0% | 0.0% | 6.7% | 0.0% | 17.8% | 8.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 54.1% |
| 1995 | 13.5% | 0.0% | 2.8% | 9.8% | 0.0% | 31.6% | 6.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 36.3% |
| 1996 | 10.1% | 0.2% | 6.4% | 0.2% | 0.0% | 20.5% | 5.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 57.0% |
| 1997 | 12.0% | 0.0% | 8.5% | 0.0% | 0.0% | 8.7% | 12.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 58.6% |
| 1998 | 10.4% | 0.0% | 3.3% | 0.0% | 0.0% | 1.4% | 6.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 78.7% |
| 1999 | 16.0% | 0.0% | 12.3% | 0.0% | 0.0% | 1.0% | 8.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 62.5% |
| 2000 | 8.6% | 0.0% | 8.9% | 0.0% | 0.0% | 9.9% | 7.2% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 65.1% |
| 2001 | 8.9% | 0.0% | 5.4% | 0.4% | 0.0% | 13.6% | 10.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 60.7% |
| 2002 | 13.0% | 0.6% | 6.1% | 1.4% | 0.0% | 4.7% | 18.4% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 55.3% |
| 2003 | 15.2% | 0.0% | 3.4% | 6.4% | 0.0% | 0.0% | 14.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 60.4% |
| 2004 | 9.7% | 3.1% | 3.8% | 1.4% | 0.0% | 2.1% | 14.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 64.9% |
| (85-98) | 14.9% | 0.3% | 3.5% | 7.0% | 0.1% | 13.3% | 6.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 54.3% |
| (99-04) | 11.9% | 0.6% | 6.6% | 1.6% | 0.0% | 5.2% | 12.4% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 61.5% |

¹ Values represent estimates of fishing mortality distribution only for this year.

Appendix E.5. Percent distribution of Robertson Creek Fall Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 18.4% | 0.8% | 0.6% | 11.6% | 10.8% | 7.7% | 0.3% | 8.0% | 1.7% | 2.2% | 5.2% | 0.0% | 0.1% | 0.0% | 32.3% |
| 1980 | 26.9% | 7.0% | 0.9% | 8.1% | 8.3% | 4.5% | 0.1% | 7.0% | 0.1% | 11.2% | 3.4% | 0.0% | 0.2% | 0.0% | 22.5% |
| 1981 | 29.7% | 1.6% | 0.8% | 12.2% | 8.2% | 4.9% | 0.5% | 5.3% | 0.6% | 13.5% | 5.7% | 0.0% | 0.4% | 0.0% | 16.5% |
| 1982 | 25.0% | 3.4% | 1.5% | 13.5% | 7.5% | 5.0% | 0.1% | 5.8% | 0.9% | 14.8% | 6.4% | 0.1% | 0.5% | 0.2% | 15.3% |
| 1983 | 36.0% | 3.3% | 0.6% | 10.4% | 8.0% | 2.4% | 0.3% | 5.3% | 0.3% | 18.2% | 4.6% | 0.0% | 0.2% | 0.0% | 10.4% |
| 1984 | 26.6% | 4.0% | 0.2% | 14.7% | 3.0% | 2.7% | 0.0% | 6.7% | 0.8% | 17.7% | 15.9% | 0.0% | 0.2% | 0.0% | 7.6% |
| 1985 | 14.1% | 5.8% | 0.0% | 17.7% | 0.5% | 4.5% | 0.0% | 2.0% | 0.8% | 3.6% | 17.7% | 0.0% | 2.0% | 0.0% | 31.3% |
| 1986 | 13.9% | 4.6% | 0.0% | 8.1% | 1.1% | 3.1% | 0.7% | 4.4% | 0.0% | 1.5% | 26.6% | 0.0% | 0.0% | 1.1% | 35.0% |
| 1987 | 6.5% | 1.5% | 0.6% | 6.1% | 2.9% | 2.4% | 0.5% | 2.2% | 0.5% | 1.1% | 20.9% | 0.0% | 0.3% | 0.1% | 54.3% |
| 1988 | 9.9% | 2.1% | 0.9% | 6.6% | 1.2% | 2.0% | 1.1% | 4.1% | 0.6% | 8.1% | 18.6% | 0.0% | 0.3% | 0.2% | 44.4% |
| 1989 | 8.0% | 2.5% | 0.4% | 7.8% | 0.8% | 1.1% | 1.0% | 1.6% | 0.8% | 20.5% | 18.5% | 0.0% | 0.1% | 0.1% | 36.9% |
| 1990 | 15.8% | 1.1% | 1.3% | 7.3% | 2.0% | 1.7% | 0.9% | 6.3% | 0.3% | 10.4% | 10.8% | 0.0% | 0.0% | 0.1% | 41.9% |
| 1991 | 16.9% | 1.1% | 3.1% | 9.1% | 2.7% | 0.6% | 0.8% | 4.4% | 0.3% | 14.9% | 13.6% | 0.0% | 0.0% | 0.1% | 32.3% |
| 1992 | 13.7% | 3.0% | 1.7% | 7.2% | 3.0% | 0.9% | 1.5% | 18.8% | 0.1% | 0.8% | 8.0% | 0.0% | 0.1% | 0.1% | 41.1% |
| 1993 | 13.9% | 1.0% | 2.5% | 7.1% | 2.0% | 0.4% | 1.4% | 13.7% | 0.5% | 8.4% | 15.7% | 0.0% | 0.0% | 0.1% | 33.2% |
| 1994 | 15.8% | 2.2% | 3.7% | 9.5% | 1.1% | 1.1% | 1.1% | 5.3% | 0.4% | 12.8% | 21.3% | 0.0% | 0.0% | 0.1% | 25.6% |
| 1995 | 15.3% | 0.0% | 4.0% | 3.1% | 0.3% | 0.3% | 0.9% | 1.5% | 1.4% | 7.3% | 12.5% | 0.0% | 0.2% | 0.0% | 53.2% |
| 1996 | 5.6% | 0.1% | 1.9% | 0.0% | 0.7% | 0.0% | 2.8% | 0.0% | 1.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 87.4% |
| 1997 | 10.7% | 3.2% | 3.9% | 4.5% | 1.8% | 0.4% | 2.8% | 0.1% | 0.5% | 6.5% | 20.0% | 0.1% | 0.0% | 0.0% | 45.1% |
| 1998 | 16.5% | 1.2% | 5.1% | 6.2% | 0.0% | 0.0% | 2.0% | 0.0% | 0.6% | 4.2% | 19.1% | 0.1% | 0.0% | 0.0% | 45.1% |
| 1999 | 12.2% | 0.4% | 7.9% | 3.3% | 0.2% | 0.0% | 2.9% | 0.0% | 0.8% | 7.0% | 22.3% | 0.0% | 0.0% | 0.0% | 42.9% |
| 2000 | 5.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.4% | 0.0% | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 88.4% |
| 2001 | 3.2% | 0.0% | 2.2% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 2.0% | 0.0% | 3.2% | 0.0% | 0.0% | 0.0% | 88.9% |
| 2002 | 12.4% | 0.3% | 1.7% | 3.0% | 0.2% | 0.0% | 4.2% | 0.4% | 0.7% | 8.6% | 8.9% | 0.0% | 0.0% | 0.0% | 59.6% |
| 2003 | 11.4% | 1.7% | 2.7% | 0.6% | 0.0% | 0.0% | 8.7% | 0.0% | 0.4% | 2.7% | 22.6% | 0.0% | 0.0% | 0.0% | 49.1% |
| 2004 | 10.1% | 6.4% | 2.3% | 1.9% | 0.0% | 0.0% | 4.2% | 0.1% | 1.1% | 10.6% | 12.7% | 0.0% | 0.0% | 0.1% | 50.4% |
| (79-84) | 27.1% | 3.3% | 0.8% | 11.8% | 7.6% | 4.5% | 0.2% | 6.4% | 0.7% | 12.9% | 6.9% | 0.0% | 0.3% | 0.0% | 17.4% |
| (85-98) | 12.6% | 2.1% | 2.1% | 7.2% | 1.4% | 1.3% | 1.3% | 4.6% | 0.6% | 7.1% | 16.0% | 0.0% | 0.2% | 0.1% | 43.3% |
| (99-04) | 9.2% | 1.5% | 2.8% | 1.5% | 0.1% | 0.0% | 4.0% | 0.1% | 1.3% | 4.8% | 11.6% | 0.0% | 0.0% | 0.0% | 63.2% |

Appendix E.6. Percent distribution of Robertson Creek Fall Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 21.1% | 0.7% | 0.7% | 13.0% | 11.9% | 7.0% | 0.3% | 8.9% | 1.5% | 2.0% | 4.8% | 0.0% | 0.1% | 0.0% | 27.8% |
| 1980 | 27.7% | 6.9% | 1.0% | 8.6% | 8.7% | 4.4% | 0.1% | 7.5% | 0.1% | 10.6% | 3.4% | 0.0% | 0.2% | 0.0% | 20.6% |
| 1981 | 32.9% | 1.5% | 1.0% | 13.1% | 8.9% | 4.4% | 0.5% | 5.8% | 0.6% | 11.9% | 5.2% | 0.0% | 0.5% | 0.0% | 13.7% |
| 1982 | 28.6% | 3.1% | 1.6% | 14.2% | 7.9% | 4.6% | 0.1% | 6.1% | 0.8% | 13.2% | 5.9% | 0.1% | 0.6% | 0.1% | 13.0% |
| 1983 | 40.6% | 3.0% | 0.6% | 10.1% | 7.7% | 2.2% | 0.3% | 5.1% | 0.3% | 16.5% | 4.4% | 0.0% | 0.2% | 0.0% | 9.1% |
| 1984 | 27.9% | 3.8% | 0.2% | 14.7% | 3.0% | 2.7% | 0.0% | 6.9% | 0.8% | 16.7% | 15.9% | 0.0% | 0.2% | 0.0% | 7.1% |
| 1985 | 14.9% | 16.8% | 0.0% | 16.0% | 0.4% | 3.7% | 0.0% | 1.8% | 0.7% | 2.9% | 15.5% | 0.0% | 1.9% | 0.0% | 25.4% |
| 1986 | 18.0% | 12.7% | 0.0% | 8.7% | 1.2% | 2.9% | 1.1% | 4.4% | 0.0% | 1.2% | 22.2% | 0.0% | 0.0% | 1.1% | 26.4% |
| 1987 | 10.2% | 3.4% | 1.1% | 7.5% | 3.5% | 2.3% | 0.6% | 2.7% | 0.5% | 1.0% | 19.8% | 0.0% | 0.3% | 0.1% | 47.1% |
| 1988 | 11.0% | 4.8% | 1.2% | 7.3% | 1.3% | 1.9% | 1.1% | 4.7% | 0.7% | 7.5% | 18.3% | 0.0% | 0.4% | 0.2% | 39.7% |
| 1989 | 11.0% | 6.9% | 0.5% | 9.0% | 1.0% | 1.1% | 1.0% | 1.9% | 0.8% | 18.3% | 17.2% | 0.0% | 0.1% | 0.1% | 31.0% |
| 1990 | 19.5% | 2.9% | 1.5% | 8.8% | 2.3% | 1.6% | 0.9% | 6.7% | 0.3% | 9.4% | 10.0% | 0.0% | 0.0% | 0.1% | 35.9% |
| 1991 | 20.0% | 2.4% | 3.3% | 9.8% | 2.9% | 0.6% | 0.8% | 4.8% | 0.3% | 13.6% | 13.0% | 0.0% | 0.0% | 0.1% | 28.5% |
| 1992 | 16.8% | 8.3% | 1.7% | 7.4% | 3.0% | 0.8% | 1.4% | 18.6% | 0.1% | 0.6% | 7.1% | 0.0% | 0.1% | 0.0% | 34.0% |
| 1993 | 16.0% | 2.3% | 2.5% | 7.6% | 2.1% | 0.4% | 1.4% | 14.4% | 0.5% | 7.7% | 15.1% | 0.0% | 0.0% | 0.1% | 29.9% |
| 1994 | 18.1% | 4.9% | 3.6% | 9.2% | 1.0% | 1.0% | 1.1% | 5.2% | 0.4% | 11.7% | 20.6% | 0.0% | 0.0% | 0.1% | 23.1% |
| 1995 | 17.5% | 0.0% | 4.6% | 3.7% | 0.4% | 0.5% | 1.1% | 1.9% | 1.5% | 6.8% | 13.3% | 0.0% | 0.2% | 0.0% | 48.6% |
| 1996 | 9.2% | 0.1% | 4.5% | 2.7% | 0.7% | 0.0% | 5.8% | 0.7% | 1.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 74.4% |
| 1997 | 13.8% | 8.2% | 4.4% | 5.0% | 2.0% | 0.4% | 3.1% | 0.2% | 0.6% | 6.0% | 18.1% | 0.1% | 0.0% | 0.0% | 38.1% |
| 1998 | 17.0% | 3.1% | 5.1% | 6.2% | 0.0% | 0.0% | 2.3% | 0.0% | 0.6% | 4.0% | 19.2% | 0.1% | 0.0% | 0.0% | 42.4% |
| 1999 | 12.9% | 0.8% | 8.1% | 3.3% | 0.2% | 0.0% | 3.2% | 0.0% | 0.8% | 6.7% | 23.1% | 0.0% | 0.0% | 0.0% | 40.8% |
| 2000 | 6.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.1% | 0.0% | 3.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 84.1% |
| 2001 | 4.6% | 0.0% | 4.0% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 2.5% | 0.0% | 4.1% | 0.0% | 0.0% | 0.0% | 84.1% |
| 2002 | 14.6% | 0.8% | 2.2% | 3.4% | 0.2% | 0.0% | 5.4% | 0.4% | 0.8% | 8.2% | 9.5% | 0.0% | 0.0% | 0.0% | 54.5% |
| 2003 | 12.2% | 4.6% | 3.3% | 0.7% | 0.0% | 0.0% | 11.2% | 0.0% | 0.5% | 2.4% | 23.0% | 0.0% | 0.0% | 0.0% | 42.1% |
| 2004 | 10.3% | 18.7% | 2.2% | 1.9% | 0.0% | 0.0% | 4.7% | 0.1% | 1.2% | 8.8% | 11.9% | 0.0% | 0.0% | 0.0% | 40.2% |
| (79-84) | 29.8% | 3.2% | 0.8% | 12.3% | 8.0% | 4.2% | 0.2% | 6.7% | 0.7% | 11.8% | 6.6% | 0.0% | 0.3% | 0.0% | 15.2% |
| (85-98) | 15.2% | 5.5% | 2.4% | 7.8% | 1.6% | 1.2% | 1.6% | 4.9% | 0.6% | 6.5% | 15.0% | 0.0% | 0.2% | 0.1% | 37.5% |
| (99-04) | 10.2% | 4.2% | 3.3% | 1.6% | 0.1% | 0.0% | 5.2% | 0.1% | 1.5% | 4.4% | 11.9% | 0.0% | 0.0% | 0.0% | 57.6% |

Appendix E.7. Percent distribution of Quinsam River Fall Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 6.3% | 6.8% | 0.9% | 7.3% | 13.6% | 25.7% | 4.1% | 0.0% | 9.3% | 5.8% | 0.0% | 0.0% | 0.0% | 0.0% | 20.1% |
| 1980 | 15.4% | 5.2% | 3.1% | 10.9% | 17.2% | 13.5% | 5.5% | 0.0% | 7.0% | 9.1% | 0.0% | 0.0% | 0.0% | 0.0% | 13.0% |
| 1981 | 12.7% | 2.8% | 1.9% | 15.4% | 14.2% | 12.2% | 7.5% | 0.7% | 13.9% | 7.6% | 0.0% | 0.0% | 0.0% | 0.0% | 11.2% |
| 1982 | 17.9% | 7.8% | 5.5% | 8.1% | 7.0% | 21.1% | 2.5% | 0.4% | 4.2% | 8.2% | 0.0% | 0.0% | 0.0% | 0.0% | 17.4% |
| 1983 | 22.0% | 1.6% | 2.1% | 15.4% | 12.1% | 17.8% | 2.8% | 0.7% | 4.9% | 8.8% | 0.0% | 0.0% | 0.0% | 0.0% | 11.7% |
| 1984 | 14.5% | 6.0% | 4.7% | 5.9% | 5.0% | 15.1% | 4.1% | 0.8% | 7.9% | 6.6% | 0.0% | 0.0% | 0.0% | 0.0% | 29.3% |
| 1985 | 25.9% | 5.8% | 4.4% | 5.1% | 3.6% | 11.1% | 1.0% | 0.1% | 4.4% | 8.3% | 0.0% | 0.0% | 0.0% | 0.0% | 30.2% |
| 1986 | 13.8% | 4.3% | 2.8% | 6.6% | 7.3% | 19.9% | 2.9% | 0.0% | 6.2% | 6.4% | 0.0% | 0.0% | 0.0% | 0.0% | 29.7% |
| 1987 | 10.8% | 3.7% | 2.8% | 6.3% | 6.1% | 17.3% | 6.6% | 0.4% | 4.0% | 7.3% | 0.4% | 0.0% | 0.0% | 0.0% | 34.3% |
| 1988 | 19.0% | 1.8% | 1.0% | 6.6% | 2.4% | 5.5% | 2.9% | 0.7% | 3.7% | 4.1% | 0.9% | 0.0% | 0.0% | 0.1% | 51.2% |
| 1989 | 12.6% | 2.8% | 2.8% | 3.9% | 1.9% | 4.9% | 3.2% | 0.3% | 7.3% | 13.0% | 0.0% | 0.0% | 0.1% | 0.0% | 47.1% |
| 1990 | 17.3% | 2.2% | 0.6% | 6.7% | 4.9% | 11.2% | 9.0% | 1.4% | 3.5% | 4.8% | 0.0% | 0.0% | 0.0% | 0.0% | 38.5% |
| 1991 | 10.1% | 2.8% | 1.4% | 5.7% | 9.1% | 10.2% | 11.9% | 0.5% | 4.4% | 3.5% | 0.8% | 0.0% | 0.0% | 0.0% | 39.7% |
| 1992 | 11.5% | 0.5% | 2.4% | 10.1% | 9.3% | 7.4% | 6.3% | 0.3% | 3.5% | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% | 46.2% |
| 1993 | 8.0% | 3.4% | 1.2% | 5.8% | 5.8% | 19.6% | 8.9% | 1.2% | 10.7% | 3.4% | 0.0% | 0.0% | 0.0% | 0.0% | 32.1% |
| 1994 | 5.3% | 6.0% | 4.0% | 9.3% | 1.3% | 14.0% | 5.0% | 0.0% | 6.0% | 4.0% | 0.0% | 0.0% | 0.0% | 0.0% | 45.0% |
| 1995 | 7.1% | 5.0% | 0.0% | 9.2% | 0.0% | 14.6% | 7.9% | 0.0% | 6.7% | 0.8% | 0.0% | 0.0% | 0.0% | 0.0% | 48.8% |
| 1996 | 6.8% | 0.4% | 0.0% | 0.0% | 0.0% | 17.4% | 4.5% | 0.0% | 6.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 64.5% |
| 1997 | 9.1% | 3.2% | 2.5% | 4.1% | 3.4% | 2.3% | 8.9% | 0.7% | 8.7% | 0.2% | 5.0% | 0.0% | 0.0% | 0.0% | 51.9% |
| 1998 | 14.2% | 2.2% | 2.0% | 0.0% | 0.0% | 0.4% | 8.7% | 0.0% | 5.4% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 66.8% |
| 1999 | 9.0% | 3.4% | 4.2% | 1.3% | 0.2% | 1.4% | 9.5% | 0.0% | 1.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 69.3% |
| 2000 | 13.1% | 2.2% | 5.0% | 0.3% | 0.0% | 0.0% | 5.6% | 0.0% | 2.7% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 70.6% |
| 2001 | 9.5% | 1.4% | 1.8% | 0.1% | 0.0% | 0.0% | 5.8% | 0.0% | 1.7% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 79.8% |
| 2002 | 14.7% | 3.1% | 0.9% | 0.4% | 0.1% | 0.0% | 11.8% | 0.0% | 2.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 66.2% |
| 2003 | 15.4% | 1.6% | 0.6% | 0.0% | 0.0% | 0.0% | 27.7% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 54.4% |
| 2004 | 8.4% | 13.3% | 1.6% | 0.3% | 0.0% | 0.9% | 15.4% | 0.0% | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 59.1% |
| (79-84) | 14.8% | 5.0% | 3.0% | 10.5% | 11.5% | 17.6% | 4.4% | 0.4% | 7.9% | 7.7% | 0.0% | 0.0% | 0.0% | 0.0% | 17.1% |
| (85-98) | 12.2% | 3.1% | 2.0% | 5.7% | 3.9% | 11.1% | 6.3% | 0.4% | 5.8% | 4.2% | 0.5% | 0.0% | 0.0% | 0.0% | 44.7% |
| (99-04) | 11.7% | 4.2% | 2.3% | 0.4% | 0.1% | 0.4% | 12.6% | 0.0% | 1.7% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 66.6% |

Appendix E.8. Percent distribution of Quinsam River Fall Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 8.4% | 6.5% | 1.5% | 8.7% | 15.3% | 24.0% | 3.9% | 0.1% | 8.5% | 5.6% | 0.0% | 0.0% | 0.0% | 0.0% | 17.6% |
| 1980 | 16.0% | 5.0% | 3.3% | 11.5% | 18.1% | 13.4% | 5.3% | 0.0% | 6.8% | 8.8% | 0.0% | 0.0% | 0.0% | 0.0% | 11.7% |
| 1981 | 13.3% | 2.6% | 2.0% | 16.5% | 14.8% | 11.7% | 7.5% | 0.7% | 13.5% | 7.1% | 0.0% | 0.0% | 0.0% | 0.0% | 10.1% |
| 1982 | 21.7% | 7.5% | 5.9% | 8.4% | 7.3% | 20.2% | 2.4% | 0.4% | 3.9% | 7.5% | 0.0% | 0.0% | 0.0% | 0.0% | 14.9% |
| 1983 | 25.9% | 1.5% | 2.6% | 15.2% | 11.9% | 17.0% | 2.9% | 0.7% | 4.5% | 8.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.9% |
| 1984 | 15.9% | 6.0% | 5.5% | 6.2% | 5.2% | 14.9% | 4.2% | 0.9% | 7.8% | 6.3% | 0.0% | 0.0% | 0.0% | 0.0% | 27.2% |
| 1985 | 27.4% | 12.8% | 4.2% | 4.7% | 3.3% | 10.0% | 1.0% | 0.1% | 4.0% | 7.2% | 0.0% | 0.0% | 0.0% | 0.0% | 25.3% |
| 1986 | 15.4% | 10.9% | 3.1% | 6.6% | 7.2% | 18.5% | 3.0% | 0.0% | 5.5% | 5.8% | 0.0% | 0.0% | 0.0% | 0.0% | 24.0% |
| 1987 | 16.0% | 10.4% | 2.8% | 6.8% | 6.7% | 14.4% | 5.7% | 0.4% | 3.4% | 6.0% | 0.3% | 0.0% | 0.0% | 0.0% | 27.0% |
| 1988 | 20.0% | 4.5% | 1.1% | 7.0% | 2.6% | 5.5% | 3.0% | 0.8% | 3.9% | 3.9% | 0.9% | 0.0% | 0.0% | 0.2% | 46.5% |
| 1989 | 14.2% | 8.1% | 2.8% | 4.1% | 2.0% | 4.6% | 3.2% | 0.3% | 7.6% | 11.9% | 0.0% | 0.0% | 0.1% | 0.0% | 41.2% |
| 1990 | 18.7% | 5.5% | 0.6% | 7.4% | 5.3% | 10.5% | 8.9% | 1.5% | 3.7% | 4.4% | 0.0% | 0.0% | 0.0% | 0.0% | 33.5% |
| 1991 | 11.4% | 7.9% | 1.4% | 5.9% | 9.4% | 9.1% | 11.3% | 0.5% | 4.5% | 3.2% | 0.7% | 0.0% | 0.0% | 0.0% | 34.6% |
| 1992 | 15.7% | 1.1% | 2.5% | 10.6% | 9.6% | 7.2% | 6.3% | 0.3% | 3.7% | 2.4% | 0.0% | 0.0% | 0.0% | 0.0% | 40.6% |
| 1993 | 8.9% | 7.1% | 1.3% | 6.5% | 6.5% | 18.1% | 8.6% | 1.3% | 11.3% | 2.9% | 0.0% | 0.0% | 0.0% | 0.0% | 27.5% |
| 1994 | 6.8% | 12.8% | 4.0% | 9.7% | 1.4% | 12.5% | 4.8% | 0.0% | 6.3% | 3.4% | 0.0% | 0.0% | 0.0% | 0.0% | 38.4% |
| 1995 | 8.6% | 5.1% | 0.0% | 11.3% | 0.0% | 16.8% | 9.6% | 0.0% | 6.5% | 2.1% | 0.0% | 0.0% | 0.0% | 0.0% | 40.1% |
| 1996 | 7.3% | 0.7% | 0.0% | 1.3% | 0.0% | 19.8% | 7.6% | 0.0% | 6.6% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 56.4% |
| 1997 | 10.1% | 5.8% | 2.9% | 4.3% | 3.5% | 2.3% | 10.9% | 0.8% | 8.9% | 1.4% | 4.7% | 0.0% | 0.0% | 0.0% | 44.4% |
| 1998 | 14.7% | 6.5% | 2.4% | 0.0% | 0.0% | 0.3% | 11.3% | 0.0% | 5.9% | 0.2% | 0.0% | 0.0% | 0.5% | 0.0% | 58.3% |
| 1999 | 10.3% | 7.2% | 5.2% | 1.4% | 0.2% | 1.7% | 11.2% | 0.0% | 1.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 61.1% |
| 2000 | 14.5% | 3.8% | 5.6% | 0.2% | 0.0% | 0.0% | 7.0% | 0.0% | 3.1% | 1.7% | 0.0% | 0.0% | 0.0% | 0.0% | 64.1% |
| 2001 | 10.6% | 2.8% | 2.0% | 0.1% | 0.0% | 0.0% | 7.3% | 0.0% | 1.8% | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 74.9% |
| 2002 | 15.3% | 7.0% | 0.9% | 0.4% | 0.1% | 0.0% | 14.4% | 0.0% | 3.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 58.7% |
| 2003 | 16.0% | 4.7% | 0.7% | 0.0% | 0.0% | 0.0% | 33.1% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 45.3% |
| 2004 | 7.1% | 30.9% | 1.4% | 0.2% | 0.0% | 1.2% | 16.5% | 0.0% | 1.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 41.7% |
| (79-84) | 16.9% | 4.9% | 3.5% | 11.1% | 12.1% | 16.9% | 4.4% | 0.5% | 7.5% | 7.2% | 0.0% | 0.0% | 0.0% | 0.0% | 15.2% |
| (85-98) | 13.9% | 7.1% | 2.1% | 6.2% | 4.1% | 10.7% | 6.8% | 0.4% | 5.8% | 3.9% | 0.5% | 0.0% | 0.0% | 0.0% | 38.4% |
| (99-04) | 12.3% | 9.4% | 2.6% | 0.4% | 0.1% | 0.5% | 14.9% | 0.0% | 1.9% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 57.6% |

Appendix E.9. Percent distribution of Puntledge River Summer Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 1.6% | 0.3% | 0.2% | 3.2% | 8.3% | 6.7% | 0.3% | 0.9% | 39.7% | 6.5% | 0.0% | 0.0% | 0.0% | 0.0% | 32.2% |
| 1980 | 2.4% | 0.0% | 0.4% | 2.0% | 5.9% | 4.4% | 1.3% | 4.9% | 38.6% | 5.9% | 0.0% | 0.0% | 0.0% | 0.0% | 34.2% |
| 1981 | 0.8% | 0.0% | 0.0% | 5.4% | 7.2% | 3.6% | 4.0% | 0.0% | 60.2% | 5.4% | 0.0% | 0.0% | 0.0% | 0.0% | 13.3% |
| 1982 | 0.9% | 0.3% | 0.0% | 2.2% | 12.8% | 5.5% | 1.0% | 1.6% | 19.2% | 14.7% | 0.0% | 0.0% | 0.0% | 0.0% | 41.7% |
| 1983 | 1.0% | 0.2% | 0.0% | 7.5% | 16.0% | 5.1% | 3.0% | 2.4% | 25.5% | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% | 36.6% |
| 1984 | 0.0% | 1.2% | 0.0% | 2.0% | 5.9% | 3.9% | 1.2% | 2.3% | 26.6% | 2.7% | 0.0% | 0.0% | 0.0% | 0.0% | 54.3% |
| 1985 | 10.9% | 0.8% | 2.3% | 6.2% | 1.6% | 8.5% | 6.2% | 0.0% | 33.3% | 6.2% | 0.0% | 0.0% | 0.0% | 0.0% | 24.0% |
| 1986 | 5.6% | 0.0% | 4.5% | 2.8% | 3.9% | 10.1% | 0.0% | 2.8% | 43.3% | 1.7% | 0.0% | 0.0% | 0.0% | 0.0% | 25.3% |
| 1987 | 2.7% | 0.7% | 0.0% | 12.1% | 2.0% | 6.7% | 10.1% | 0.0% | 16.8% | 0.0% | 4.7% | 0.0% | 0.0% | 0.0% | 44.3% |
| 1988 | 12.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.3% | 14.1% | 0.0% | 17.4% | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 51.1% |
| 1989 | 3.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 45.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 51.5% |
| 1990 | 9.4% | 0.0% | 0.0% | 0.0% | 3.5% | 11.8% | 3.5% | 0.0% | 9.4% | 4.7% | 0.0% | 0.0% | 0.0% | 0.0% | 57.6% |
| 1991 | 5.2% | 5.2% | 0.0% | 0.0% | 0.0% | 5.2% | 7.8% | 0.0% | 23.5% | 5.2% | 0.0% | 0.0% | 0.0% | 0.0% | 47.8% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.0% | 3.5% | 0.0% | 37.2% | 15.1% | 0.0% | 0.0% | 0.0% | 0.0% | 37.2% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.6% | 10.5% | 0.0% | 44.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 38.2% |
| 1994 | 7.1% | 0.0% | 0.0% | 0.0% | 0.0% | 7.1% | 0.0% | 0.0% | 53.6% | 3.6% | 0.0% | 0.0% | 0.0% | 0.0% | 28.6% |
| 1995 | 5.9% | 2.9% | 0.0% | 0.0% | 0.0% | 14.7% | 0.0% | 0.0% | 32.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 44.1% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.6% | 7.9% | 0.0% | 34.2% | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% | 52.6% |
| 1997 | 0.0% | 0.0% | 0.0% | 9.8% | 0.0% | 7.8% | 13.7% | 0.0% | 7.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 60.8% |
| 1998 | 21.2% | 6.1% | 0.0% | 0.0% | 0.0% | 0.0% | 15.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 57.6% |
| 1999 | 9.1% | 0.0% | 0.0% | 0.0% | 0.0% | 1.8% | 9.1% | 0.0% | 12.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 67.9% |
| 2000 | 1.6% | 0.8% | 0.0% | 1.6% | 0.0% | 0.0% | 0.0% | 0.0% | 25.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 71.1% |
| 2001 | 8.3% | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 4.9% | 1.5% | 7.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 77.4% |
| 2002 | 5.8% | 0.6% | 0.0% | 0.6% | 0.0% | 0.0% | 10.9% | 0.0% | 5.1% | 1.3% | 7.7% | 0.0% | 0.0% | 0.0% | 67.9% |
| 2003 | 3.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 37.7% | 0.0% | 11.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 46.6% |
| 2004 | 11.3% | 0.8% | 0.0% | 3.0% | 0.0% | 0.0% | 19.5% | 2.3% | 13.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 49.6% |
| (79-84) | 1.1% | 0.3% | 0.1% | 3.7% | 9.4% | 4.9% | 1.8% | 2.0% | 35.0% | 6.3% | 0.0% | 0.0% | 0.0% | 0.0% | 35.4% |
| (85-98) | 5.9% | 1.1% | 0.5% | 2.2% | 0.8% | 6.6% | 6.6% | 0.2% | 28.5% | 2.9% | 0.3% | 0.0% | 0.0% | 0.0% | 44.3% |
| (99-04) | 6.6% | 0.5% | 0.0% | 0.9% | 0.0% | 0.4% | 13.7% | 0.6% | 12.4% | 0.2% | 1.3% | 0.0% | 0.0% | 0.0% | 63.4% |

Appendix E.10. Percent distribution of Puntledge River Summer Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 2.0% | 0.3% | 0.3% | 4.4% | 10.5% | 6.5% | 0.3% | 1.2% | 38.7% | 6.3% | 0.0% | 0.0% | 0.0% | 0.0% | 29.6% |
| 1980 | 2.8% | 0.0% | 0.5% | 2.3% | 6.8% | 4.6% | 1.4% | 5.7% | 38.4% | 5.9% | 0.0% | 0.0% | 0.0% | 0.0% | 31.6% |
| 1981 | 0.9% | 0.0% | 0.0% | 6.6% | 8.8% | 3.3% | 4.0% | 0.0% | 58.9% | 5.3% | 0.0% | 0.0% | 0.0% | 0.0% | 12.1% |
| 1982 | 0.9% | 0.5% | 0.0% | 2.5% | 14.7% | 5.8% | 1.3% | 1.9% | 19.3% | 15.0% | 0.0% | 0.0% | 0.0% | 0.0% | 38.1% |
| 1983 | 2.0% | 0.2% | 0.0% | 8.2% | 17.0% | 5.1% | 3.1% | 2.6% | 25.5% | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% | 33.8% |
| 1984 | 0.0% | 1.1% | 0.0% | 2.6% | 6.6% | 4.0% | 1.5% | 2.6% | 27.6% | 2.9% | 0.0% | 0.0% | 0.0% | 0.0% | 51.1% |
| 1985 | 14.2% | 1.4% | 3.4% | 6.8% | 1.4% | 8.8% | 6.8% | 0.0% | 31.1% | 5.4% | 0.0% | 0.0% | 0.0% | 0.0% | 20.9% |
| 1986 | 6.0% | 0.0% | 5.5% | 3.0% | 4.5% | 10.1% | 0.0% | 3.0% | 43.7% | 1.5% | 0.0% | 0.0% | 0.0% | 0.0% | 22.6% |
| 1987 | 3.1% | 1.2% | 0.0% | 15.3% | 2.5% | 6.1% | 10.4% | 0.0% | 16.6% | 0.0% | 4.3% | 0.0% | 0.0% | 0.0% | 40.5% |
| 1988 | 11.9% | 0.0% | 0.0% | 0.0% | 0.0% | 5.0% | 15.8% | 0.0% | 19.8% | 1.0% | 0.0% | 0.0% | 0.0% | 0.0% | 46.5% |
| 1989 | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 52.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 44.7% |
| 1990 | 10.9% | 0.0% | 0.0% | 0.0% | 4.3% | 12.0% | 4.3% | 0.0% | 9.8% | 5.4% | 0.0% | 0.0% | 0.0% | 0.0% | 53.3% |
| 1991 | 5.6% | 14.0% | 0.0% | 0.0% | 0.0% | 4.2% | 8.4% | 0.0% | 24.5% | 4.9% | 0.0% | 0.0% | 0.0% | 0.0% | 38.5% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.3% | 3.1% | 0.0% | 42.7% | 13.5% | 0.0% | 0.0% | 0.0% | 0.0% | 33.3% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.9% | 10.6% | 0.0% | 49.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 34.1% |
| 1994 | 9.4% | 0.0% | 0.0% | 0.0% | 0.0% | 6.3% | 0.0% | 0.0% | 56.3% | 3.1% | 0.0% | 0.0% | 0.0% | 0.0% | 25.0% |
| 1995 | 5.1% | 2.6% | 0.0% | 0.0% | 0.0% | 15.4% | 0.0% | 0.0% | 35.9% | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% | 38.5% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.3% | 11.4% | 0.0% | 38.6% | 2.3% | 0.0% | 0.0% | 0.0% | 0.0% | 45.5% |
| 1997 | 0.0% | 0.0% | 0.0% | 10.2% | 0.0% | 8.5% | 18.6% | 0.0% | 8.5% | 1.7% | 0.0% | 0.0% | 0.0% | 0.0% | 52.5% |
| 1998 | 19.0% | 16.7% | 0.0% | 0.0% | 0.0% | 0.0% | 19.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 45.2% |
| 1999 | 9.8% | 0.0% | 0.0% | 0.0% | 0.0% | 2.2% | 12.6% | 0.0% | 14.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 61.2% |
| 2000 | 2.2% | 1.4% | 0.0% | 1.4% | 0.0% | 0.0% | 0.0% | 0.0% | 29.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 65.5% |
| 2001 | 10.2% | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 6.8% | 1.7% | 8.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 71.7% |
| 2002 | 7.1% | 1.0% | 0.0% | 0.5% | 0.0% | 0.0% | 11.1% | 0.0% | 5.6% | 13.6% | 7.6% | 0.0% | 0.0% | 0.0% | 53.5% |
| 2003 | 3.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 43.7% | 0.0% | 12.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 39.9% |
| 2004 | 13.2% | 1.1% | 0.0% | 2.9% | 0.0% | 0.0% | 27.6% | 1.7% | 15.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 37.9% |
| (79-84) | 1.5% | 0.3% | 0.1% | 4.4% | 10.7% | 4.9% | 1.9% | 2.3% | 34.7% | 6.3% | 0.0% | 0.0% | 0.0% | 0.0% | 32.7% |
| (85-98) | 6.3% | 2.6% | 0.6% | 2.5% | 0.9% | 6.5% | 7.8% | 0.2% | 30.7% | 3.0% | 0.3% | 0.0% | 0.0% | 0.0% | 38.7% |
| (99-04) | 7.6% | 0.8% | 0.0% | 0.8% | 0.0% | 0.4% | 17.0% | 0.6% | 14.3% | 2.3% | 1.3% | 0.0% | 0.0% | 0.0% | 55.0% |

Appendix E.11. Percent distribution of Big Qualicum River Fall Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 3.4% | 0.9% | 0.3% | 1.7% | 9.4% | 4.1% | 0.4% | 2.2% | 39.3% | 8.0% | 0.1% | 0.0% | 0.3% | 0.1% | 29.8% |
| 1980 | 1.4% | 1.6% | 0.4% | 4.3% | 6.6% | 3.4% | 1.3% | 4.2% | 39.2% | 9.4% | 0.0% | 0.1% | 0.3% | 0.2% | 27.6% |
| 1981 | 1.9% | 0.3% | 0.4% | 1.3% | 11.5% | 4.5% | 0.8% | 1.6% | 54.7% | 9.7% | 0.3% | 0.0% | 0.1% | 0.6% | 12.3% |
| 1982 | 4.5% | 0.4% | 1.2% | 4.5% | 5.8% | 8.5% | 0.4% | 4.3% | 25.6% | 12.1% | 0.0% | 0.0% | 1.1% | 0.7% | 30.9% |
| 1983 | 5.4% | 0.3% | 0.3% | 4.9% | 6.8% | 4.6% | 1.0% | 1.1% | 36.6% | 14.6% | 0.0% | 0.0% | 0.0% | 0.6% | 23.7% |
| 1984 | 1.4% | 0.4% | 0.0% | 1.4% | 6.6% | 3.6% | 5.8% | 1.4% | 52.3% | 6.2% | 0.0% | 0.0% | 0.0% | 0.0% | 20.7% |
| 1985 | 3.9% | 0.3% | 0.6% | 1.7% | 3.7% | 6.8% | 1.7% | 1.4% | 35.6% | 12.4% | 0.0% | 0.0% | 2.6% | 0.0% | 29.3% |
| 1986 | 1.9% | 0.3% | 0.0% | 0.8% | 12.8% | 8.3% | 2.9% | 1.4% | 45.4% | 7.5% | 0.0% | 0.0% | 0.0% | 0.0% | 18.8% |
| 1987 | 8.8% | 0.0% | 1.0% | 4.0% | 2.5% | 2.6% | 2.7% | 4.2% | 31.7% | 5.2% | 0.0% | 0.8% | 0.7% | 0.0% | 35.8% |
| 1988 | 2.8% | 0.5% | 0.0% | 2.3% | 1.3% | 10.2% | 1.3% | 2.8% | 32.1% | 4.8% | 2.0% | 0.0% | 1.0% | 0.0% | 38.9% |
| 1989 | 4.2% | 1.6% | 0.6% | 3.2% | 0.6% | 1.0% | 1.8% | 4.8% | 39.0% | 8.2% | 0.0% | 0.2% | 0.0% | 1.0% | 34.0% |
| 1990 | 4.8% | 1.9% | 0.0% | 6.0% | 1.6% | 6.7% | 2.4% | 3.0% | 22.7% | 11.3% | 0.0% | 0.2% | 0.0% | 1.9% | 37.5% |
| 1991 | 2.4% | 1.3% | 0.0% | 2.1% | 1.1% | 2.9% | 1.9% | 1.9% | 44.7% | 5.6% | 0.0% | 0.5% | 0.5% | 0.0% | 35.0% |
| 1992 | 2.3% | 0.0% | 2.5% | 5.4% | 5.9% | 1.6% | 7.7% | 3.4% | 41.3% | 3.9% | 0.0% | 0.0% | 0.4% | 0.0% | 25.5% |
| 1993 | 1.2% | 1.2% | 0.0% | 1.5% | 3.9% | 2.9% | 3.2% | 1.7% | 45.0% | 6.8% | 0.0% | 0.0% | 0.0% | 1.0% | 31.5% |
| 1994 | 4.5% | 0.0% | 0.0% | 1.6% | 1.6% | 3.7% | 2.0% | 2.8% | 34.6% | 2.4% | 0.0% | 0.0% | 2.8% | 0.0% | 43.9% |
| 1995 | 7.0% | 0.0% | 0.0% | 1.5% | 0.0% | 7.0% | 2.5% | 0.0% | 21.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 60.5% |
| 1996 | 2.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 1.1% | 0.0% | 46.8% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 47.5% |
| 1997 | 3.0% | 0.0% | 0.0% | 5.0% | 1.5% | 1.5% | 2.0% | 0.0% | 30.5% | 0.5% | 4.5% | 0.0% | 0.0% | 0.0% | 51.5% |
| 1998 | 7.6% | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 6.5% | 0.0% | 21.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 64.1% |
| 1999 | 6.0% | 2.6% | 0.0% | 2.1% | 2.6% | 0.0% | 2.1% | 0.0% | 12.3% | 0.0% | 3.8% | 0.0% | 0.9% | 0.0% | 67.7% |
| 2000 | 14.2% | 0.9% | 0.0% | 0.0% | 0.0% | 0.5% | 3.2% | 0.0% | 11.5% | 0.0% | 0.0% | 0.0% | 3.2% | 0.0% | 66.5% |
| 2001 | 4.0% | 6.8% | 0.0% | 0.0% | 0.0% | 0.0% | 5.1% | 0.6% | 10.2% | 0.0% | 0.0% | 0.0% | 1.7% | 0.0% | 71.5% |
| 2002 | 10.4% | 0.0% | 3.1% | 2.8% | 0.0% | 0.0% | 7.6% | 2.4% | 9.7% | 0.3% | 1.7% | 0.0% | 2.1% | 1.0% | 58.7% |
| 2003 | 8.1% | 0.4% | 1.7% | 0.0% | 0.0% | 0.0% | 20.8% | 3.4% | 7.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 58.5% |
| 2004 | 6.1% | 0.0% | 0.2% | 3.5% | 0.0% | 0.0% | 4.5% | 0.9% | 6.8% | 0.0% | 0.0% | 0.5% | 1.2% | 0.0% | 76.2% |
| (79-84) | 3.0% | 0.7% | 0.4% | 3.0% | 7.8% | 4.8% | 1.6% | 2.5% | 41.3% | 10.0% | 0.1% | 0.0% | 0.3% | 0.4% | 24.2% |
| (85-98) | 4.1% | 0.5% | 0.3% | 2.5% | 2.6% | 4.0% | 2.8% | 2.0% | 35.1% | 4.9% | 0.5% | 0.1% | 0.6% | 0.4% | 39.6% |
| (99-04) | 8.1% | 1.8% | 0.8% | 1.4% | 0.4% | 0.1% | 7.2% | 1.2% | 9.6% | 0.1% | 0.9% | 0.1% | 1.5% | 0.2% | 66.5% |

Appendix E.12. Percent distribution of Big Qualicum Fall Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 4.3% | 0.9% | 0.4% | 2.2% | 11.7% | 4.0% | 0.4% | 2.8% | 38.0% | 7.6% | 0.1% | 0.0% | 0.3% | 0.1% | 27.1% |
| 1980 | 1.5% | 1.7% | 0.4% | 5.0% | 7.5% | 3.4% | 1.3% | 5.0% | 38.7% | 9.3% | 0.0% | 0.2% | 0.3% | 0.2% | 25.5% |
| 1981 | 2.4% | 0.3% | 0.4% | 1.6% | 13.4% | 4.5% | 0.8% | 1.9% | 53.1% | 9.4% | 0.3% | 0.0% | 0.2% | 0.6% | 11.1% |
| 1982 | 5.7% | 0.5% | 1.4% | 4.9% | 6.4% | 8.4% | 0.4% | 4.9% | 25.2% | 11.8% | 0.0% | 0.0% | 1.1% | 0.6% | 28.7% |
| 1983 | 5.5% | 0.3% | 0.7% | 5.0% | 7.2% | 4.8% | 1.2% | 1.2% | 37.6% | 14.1% | 0.0% | 0.0% | 0.0% | 1.0% | 21.5% |
| 1984 | 2.3% | 0.4% | 0.0% | 1.6% | 7.2% | 3.6% | 6.5% | 1.6% | 52.0% | 6.1% | 0.0% | 0.0% | 0.0% | 0.0% | 18.6% |
| 1985 | 6.8% | 1.1% | 0.9% | 2.1% | 4.3% | 6.5% | 2.1% | 1.6% | 34.3% | 12.0% | 0.0% | 0.0% | 3.3% | 0.0% | 25.0% |
| 1986 | 3.2% | 1.4% | 0.0% | 0.8% | 13.7% | 7.8% | 2.9% | 1.4% | 45.3% | 7.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.7% |
| 1987 | 10.6% | 0.0% | 1.0% | 4.3% | 2.8% | 2.5% | 2.9% | 4.8% | 31.6% | 5.0% | 0.0% | 0.9% | 0.8% | 0.0% | 33.0% |
| 1988 | 3.0% | 2.0% | 0.0% | 2.6% | 1.3% | 10.0% | 1.3% | 3.3% | 35.4% | 4.3% | 2.0% | 0.0% | 1.5% | 0.0% | 33.3% |
| 1989 | 4.5% | 4.6% | 0.8% | 3.6% | 0.5% | 0.8% | 1.8% | 5.1% | 41.1% | 7.3% | 0.0% | 0.3% | 0.0% | 1.0% | 28.4% |
| 1990 | 5.1% | 5.0% | 0.0% | 7.0% | 1.8% | 6.5% | 2.6% | 3.2% | 24.3% | 10.5% | 0.0% | 0.1% | 0.0% | 1.9% | 31.9% |
| 1991 | 3.2% | 3.6% | 0.0% | 2.4% | 1.3% | 2.7% | 1.9% | 2.1% | 47.7% | 5.0% | 0.0% | 0.5% | 0.4% | 0.0% | 29.1% |
| 1992 | 4.0% | 0.0% | 2.7% | 6.1% | 6.2% | 1.5% | 7.5% | 3.5% | 43.7% | 3.4% | 0.0% | 0.0% | 0.4% | 0.0% | 21.0% |
| 1993 | 1.6% | 2.8% | 0.0% | 1.6% | 4.7% | 2.6% | 3.0% | 1.8% | 48.4% | 6.1% | 0.0% | 0.0% | 0.0% | 1.2% | 26.1% |
| 1994 | 5.1% | 0.0% | 0.0% | 1.8% | 1.8% | 3.3% | 1.8% | 2.9% | 38.2% | 2.2% | 0.0% | 0.0% | 2.9% | 0.0% | 39.7% |
| 1995 | 7.4% | 0.0% | 0.0% | 2.2% | 0.0% | 8.7% | 3.5% | 0.0% | 22.6% | 3.0% | 0.0% | 0.0% | 0.0% | 0.0% | 52.6% |
| 1996 | 3.3% | 0.0% | 0.0% | 0.6% | 0.0% | 0.9% | 1.5% | 0.3% | 51.8% | 0.3% | 0.0% | 0.0% | 0.0% | 1.2% | 40.0% |
| 1997 | 3.9% | 0.0% | 0.0% | 5.7% | 1.7% | 1.7% | 2.6% | 0.0% | 31.9% | 3.1% | 4.4% | 0.0% | 0.0% | 0.0% | 45.0% |
| 1998 | 8.0% | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 9.0% | 0.0% | 23.4% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 58.0% |
| 1999 | 6.8% | 6.4% | 0.0% | 2.6% | 3.0% | 0.0% | 2.6% | 0.0% | 13.6% | 0.0% | 4.2% | 0.0% | 0.8% | 0.0% | 60.0% |
| 2000 | 16.6% | 2.1% | 0.0% | 0.0% | 0.0% | 0.4% | 4.1% | 0.0% | 12.9% | 0.0% | 0.0% | 0.0% | 3.7% | 0.0% | 60.2% |
| 2001 | 4.5% | 16.6% | 0.0% | 0.0% | 0.0% | 0.0% | 6.1% | 0.5% | 10.4% | 0.0% | 0.0% | 0.0% | 1.8% | 0.0% | 60.2% |
| 2002 | 11.4% | 0.0% | 3.3% | 3.0% | 0.0% | 0.0% | 9.0% | 2.1% | 10.8% | 5.1% | 1.8% | 0.0% | 2.1% | 0.9% | 50.6% |
| 2003 | 8.5% | 1.8% | 2.1% | 0.0% | 0.0% | 0.0% | 27.0% | 3.2% | 8.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 49.1% |
| 2004 | 7.0% | 0.0% | 0.4% | 4.0% | 0.0% | 0.0% | 8.6% | 1.1% | 8.6% | 0.0% | 0.0% | 0.4% | 1.5% | 0.0% | 68.4% |
| (79-84) | 3.6% | 0.7% | 0.6% | 3.4% | 8.9% | 4.8% | 1.8% | 2.9% | 40.8% | 9.7% | 0.1% | 0.0% | 0.3% | 0.4% | 22.1% |
| (85-98) | 5.0% | 1.5% | 0.4% | 2.9% | 2.9% | 4.0% | 3.2% | 2.2% | 37.1% | 5.0% | 0.5% | 0.1% | 0.7% | 0.4% | 34.3% |
| (99-04) | 9.1% | 4.5% | 1.0% | 1.6% | 0.5% | 0.1% | 9.6% | 1.1% | 10.7% | 0.8% | 1.0% | 0.1% | 1.6% | 0.1% | 58.1% |

Appendix E.13. Percent distribution of Cowichan River Fall Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 4.6% | 0.3% | 1.3% | 52.1% | 12.9% | 0.0% | 0.7% | 3.0% | 2.2% | 21.6% |
| 1991 | 0.1% | 0.0% | 0.0% | 0.2% | 0.2% | 0.6% | 1.5% | 3.2% | 57.3% | 4.8% | 0.7% | 0.9% | 3.6% | 0.8% | 26.0% |
| 1992 | 0.1% | 0.0% | 0.0% | 0.4% | 1.1% | 1.2% | 0.9% | 9.6% | 63.1% | 4.3% | 1.4% | 0.3% | 1.3% | 1.3% | 15.1% |
| 1993 | 0.2% | 0.0% | 0.0% | 0.1% | 0.5% | 0.6% | 1.5% | 7.8% | 59.6% | 3.4% | 1.6% | 0.6% | 0.9% | 0.5% | 22.8% |
| 1994 | 0.6% | 0.0% | 0.0% | 0.4% | 0.2% | 2.3% | 0.0% | 4.1% | 37.9% | 6.3% | 0.9% | 0.4% | 3.7% | 0.5% | 42.7% |
| 1995 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 0.0% | 4.0% | 33.2% | 0.5% | 0.6% | 0.0% | 2.2% | 0.8% | 57.3% |
| 1996 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 42.6% | 0.4% | 1.1% | 0.0% | 0.9% | 3.7% | 50.6% |
| 1997 | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.6% | 2.8% | 25.3% | 0.2% | 1.1% | 0.0% | 3.5% | 2.9% | 62.2% |
| 1998 | 3.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 0.5% | 26.7% | 0.3% | 1.5% | 0.0% | 2.8% | 0.0% | 63.7% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.0% | 0.0% | 38.7% | 1.2% | 4.1% | 1.0% | 6.8% | 0.7% | 46.5% |
| 2000 | 1.2% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 19.8% | 0.0% | 5.3% | 0.0% | 4.2% | 1.3% | 66.8% |
| 2001 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 11.3% | 23.4% | 0.3% | 0.0% | 0.2% | 14.9% | 0.9% | 48.0% |
| 2002 | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.7% | 4.1% | 27.7% | 0.1% | 0.7% | 0.7% | 3.0% | 3.6% | 56.2% |
| 2003 | 2.0% | 0.3% | 0.0% | 2.2% | 3.1% | 0.0% | 6.7% | 9.0% | 25.8% | 0.0% | 11.2% | 0.6% | 5.6% | 2.5% | 30.9% |
| 2004 | 0.0% | 0.3% | 0.0% | 0.6% | 0.0% | 0.0% | 4.5% | 16.6% | 21.4% | 2.6% | 14.4% | 2.6% | 6.4% | 1.9% | 28.8% |
| (90-98) | 0.7% | 0.0% | 0.0% | 0.1% | 0.4% | 1.3% | 0.6% | 3.7% | 44.2% | 3.7% | 1.0% | 0.3% | 2.4% | 1.4% | 40.2% |
| (99-04) | 0.8% | 0.1% | 0.0% | 0.5% | 0.5% | 0.0% | 2.6% | 7.1% | 26.1% | 0.7% | 6.0% | 0.8% | 6.8% | 1.8% | 46.2% |

Appendix E.14. Percent distribution of Cowichan River Fall Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1990 | 0.0% | 0.0% | 0.0% | 0.1% | 1.4% | 3.6% | 0.6% | 2.8% | 58.6% | 9.9% | 0.1% | 0.8% | 4.5% | 2.5% | 15.1% |
| 1991 | 0.1% | 0.0% | 0.0% | 0.2% | 0.4% | 0.6% | 1.4% | 4.3% | 62.1% | 4.2% | 0.7% | 0.8% | 3.7% | 0.8% | 20.7% |
| 1992 | 0.1% | 0.1% | 0.0% | 0.4% | 1.1% | 1.0% | 0.9% | 9.8% | 66.8% | 3.7% | 1.2% | 0.3% | 1.4% | 1.2% | 12.0% |
| 1993 | 0.3% | 0.0% | 0.0% | 0.1% | 0.5% | 0.5% | 1.4% | 8.2% | 63.7% | 3.0% | 1.4% | 0.6% | 0.9% | 0.5% | 18.9% |
| 1994 | 0.6% | 0.0% | 0.0% | 0.4% | 0.3% | 2.3% | 0.0% | 4.4% | 42.9% | 6.3% | 0.8% | 0.4% | 4.6% | 0.7% | 36.4% |
| 1995 | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 0.0% | 5.6% | 37.1% | 1.5% | 0.6% | 0.0% | 2.5% | 0.9% | 49.9% |
| 1996 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.3% | 47.6% | 0.5% | 1.1% | 0.0% | 1.1% | 4.7% | 43.8% |
| 1997 | 1.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.7% | 3.5% | 29.3% | 1.1% | 1.1% | 0.0% | 4.2% | 3.4% | 55.0% |
| 1998 | 4.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.5% | 30.8% | 0.5% | 1.6% | 0.0% | 3.9% | 0.0% | 57.6% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 0.0% | 43.5% | 1.0% | 4.1% | 1.0% | 9.0% | 0.6% | 39.4% |
| 2000 | 1.6% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 23.4% | 0.0% | 5.8% | 0.0% | 5.3% | 2.4% | 59.9% |
| 2001 | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 10.7% | 26.0% | 0.3% | 0.0% | 0.1% | 17.7% | 2.9% | 40.9% |
| 2002 | 1.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.2% | 3.8% | 33.0% | 0.1% | 0.9% | 0.7% | 3.2% | 5.1% | 48.6% |
| 2003 | 2.1% | 0.7% | 0.0% | 2.3% | 4.1% | 0.0% | 8.0% | 8.0% | 28.2% | 0.0% | 11.6% | 0.5% | 6.6% | 3.0% | 25.1% |
| 2004 | 0.0% | 0.8% | 0.0% | 0.5% | 0.0% | 0.0% | 6.3% | 15.1% | 24.1% | 2.4% | 14.8% | 2.6% | 7.4% | 2.1% | 23.8% |
| (90-98) | 0.8% | 0.0% | 0.0% | 0.1% | 0.4% | 1.1% | 0.7% | 4.4% | 48.8% | 3.4% | 1.0% | 0.3% | 3.0% | 1.6% | 34.4% |
| (99-04) | 0.9% | 0.3% | 0.0% | 0.5% | 0.7% | 0.0% | 3.3% | 6.5% | 29.7% | 0.6% | 6.2% | 0.8% | 8.2% | 2.7% | 39.6% |

Appendix E.15. Percent distribution of Chilliwack River Fall Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1985 | 0.5% | 0.0% | 0.0% | 0.3% | 2.3% | 0.8% | 0.2% | 34.5% | 28.9% | 5.9% | 0.0% | 4.0% | 4.2% | 3.6% | 14.6% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.8% | 2.5% | 1.5% | 0.2% | 19.5% | 28.2% | 12.6% | 0.0% | 2.6% | 4.1% | 5.9% | 22.2% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.7% | 0.4% | 0.3% | 0.3% | 16.2% | 35.3% | 2.2% | 0.5% | 3.7% | 3.9% | 2.7% | 33.5% |
| 1988 | 0.4% | 0.1% | 0.0% | 0.2% | 0.0% | 0.1% | 0.0% | 17.9% | 19.7% | 2.2% | 0.0% | 4.2% | 3.0% | 1.8% | 50.3% |
| 1989 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 19.5% | 17.4% | 3.7% | 0.0% | 5.3% | 3.7% | 1.4% | 48.3% |
| 1990 | 0.9% | 0.0% | 0.0% | 0.0% | 0.2% | 1.5% | 0.3% | 9.4% | 15.3% | 4.3% | 2.4% | 6.2% | 12.2% | 5.6% | 41.9% |
| 1991 | 0.2% | 0.1% | 0.0% | 0.4% | 0.2% | 1.0% | 0.2% | 18.3% | 21.9% | 4.2% | 0.7% | 13.4% | 5.3% | 4.6% | 29.5% |
| 1992 | 0.3% | 0.0% | 0.0% | 0.1% | 0.6% | 0.3% | 0.2% | 18.0% | 16.1% | 1.0% | 0.1% | 8.3% | 0.9% | 3.3% | 50.8% |
| 1993 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 11.9% | 14.7% | 1.5% | 0.4% | 7.1% | 0.0% | 0.9% | 63.0% |
| 1994 | 0.3% | 0.2% | 0.0% | 0.7% | 0.3% | 1.6% | 0.0% | 6.5% | 13.6% | 4.4% | 2.5% | 1.6% | 3.6% | 3.6% | 61.1% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 0.2% | 8.8% | 6.5% | 0.6% | 0.5% | 1.2% | 1.1% | 1.7% | 78.9% |
| 1996 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 0.0% | 0.0% | 15.7% | 1.1% | 0.5% | 4.5% | 0.9% | 2.8% | 73.1% |
| 1997 | 0.7% | 0.0% | 0.0% | 0.1% | 0.4% | 0.6% | 0.6% | 10.0% | 15.1% | 1.5% | 2.0% | 4.9% | 2.3% | 3.3% | 58.5% |
| 1998 | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.2% | 3.9% | 0.3% | 0.3% | 3.0% | 0.3% | 0.4% | 91.1% |
| 1999 | 0.1% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.2% | 0.3% | 10.3% | 0.5% | 1.9% | 11.6% | 0.7% | 0.9% | 73.5% |
| 2000 | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 5.0% | 5.8% | 0.0% | 2.0% | 3.8% | 0.5% | 0.4% | 81.8% |
| 2001 | 0.1% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 3.5% | 9.2% | 0.1% | 1.6% | 6.2% | 0.9% | 2.8% | 75.2% |
| 2002 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 8.5% | 7.6% | 0.2% | 2.9% | 7.1% | 0.3% | 2.5% | 70.4% |
| 2003 | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 5.3% | 11.5% | 0.2% | 5.8% | 7.0% | 0.2% | 1.3% | 68.3% |
| 2004 | 0.1% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 4.7% | 6.7% | 0.5% | 2.2% | 5.8% | 0.1% | 0.9% | 78.8% |
| (85-98) | 0.3% | 0.0% | 0.0% | 0.2% | 0.5% | 0.7% | 0.2% | 13.6% | 18.0% | 3.2% | 0.7% | 5.0% | 3.3% | 3.0% | 51.2% |
| (99-04) | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 4.6% | 8.5% | 0.2% | 2.7% | 6.9% | 0.5% | 1.5% | 74.7% |

Appendix E.16. Percent distribution of Chilliwack River Fall Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1985 | 1.1% | 0.1% | 0.0% | 0.4% | 2.3% | 0.7% | 0.2% | 34.3% | 28.8% | 5.7% | 0.0% | 3.9% | 4.9% | 4.5% | 13.1% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.8% | 2.6% | 1.5% | 0.2% | 20.6% | 28.5% | 11.7% | 0.0% | 2.8% | 5.1% | 7.2% | 19.0% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.8% | 0.5% | 0.3% | 0.3% | 19.0% | 35.8% | 2.0% | 0.5% | 3.9% | 4.0% | 2.9% | 29.9% |
| 1988 | 0.4% | 0.2% | 0.0% | 0.2% | 0.0% | 0.1% | 0.0% | 18.6% | 20.3% | 2.2% | 0.0% | 4.3% | 4.1% | 2.7% | 46.9% |
| 1989 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 24.1% | 20.9% | 3.3% | 0.0% | 6.0% | 3.7% | 1.5% | 39.7% |
| 1990 | 1.0% | 0.0% | 0.0% | 0.0% | 0.1% | 1.3% | 0.4% | 11.5% | 16.1% | 3.7% | 2.2% | 6.6% | 17.1% | 7.1% | 32.9% |
| 1991 | 0.3% | 0.2% | 0.0% | 0.4% | 0.2% | 0.9% | 0.2% | 20.0% | 24.3% | 3.6% | 0.7% | 13.8% | 6.1% | 5.2% | 24.3% |
| 1992 | 0.3% | 0.0% | 0.0% | 0.1% | 0.7% | 0.3% | 0.2% | 20.3% | 18.3% | 0.9% | 0.1% | 8.8% | 0.9% | 3.5% | 45.6% |
| 1993 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 13.5% | 17.0% | 1.4% | 0.4% | 7.6% | 0.0% | 1.0% | 58.6% |
| 1994 | 0.4% | 0.3% | 0.0% | 0.9% | 0.4% | 1.7% | 0.0% | 8.1% | 15.0% | 5.0% | 2.7% | 1.6% | 5.4% | 5.3% | 53.3% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 0.2% | 13.1% | 7.5% | 0.9% | 0.5% | 1.1% | 1.4% | 2.5% | 72.0% |
| 1996 | 0.2% | 0.0% | 0.0% | 0.1% | 0.0% | 1.4% | 0.0% | 2.1% | 18.1% | 1.3% | 0.6% | 4.3% | 1.2% | 4.6% | 66.1% |
| 1997 | 0.8% | 0.0% | 0.0% | 0.2% | 0.4% | 0.6% | 0.8% | 12.4% | 16.8% | 1.8% | 1.9% | 5.5% | 2.5% | 3.9% | 52.4% |
| 1998 | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.2% | 4.5% | 0.3% | 0.3% | 3.4% | 0.3% | 0.9% | 89.3% |
| 1999 | 0.1% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.3% | 0.3% | 12.1% | 0.5% | 1.9% | 13.6% | 0.7% | 1.0% | 69.4% |
| 2000 | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 5.5% | 6.6% | 0.0% | 2.4% | 4.5% | 0.7% | 1.1% | 78.4% |
| 2001 | 0.1% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 3.6% | 10.5% | 0.1% | 1.8% | 7.1% | 1.2% | 5.7% | 69.2% |
| 2002 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.2% | 8.7% | 8.9% | 0.2% | 3.4% | 8.3% | 0.4% | 3.2% | 66.4% |
| 2003 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 5.5% | 12.4% | 0.2% | 6.9% | 7.9% | 0.3% | 1.6% | 64.8% |
| 2004 | 0.2% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 4.9% | 7.4% | 0.5% | 2.6% | 6.7% | 0.1% | 1.3% | 76.2% |
| (85-98) | 0.4% | 0.1% | 0.0% | 0.3% | 0.5% | 0.7% | 0.2% | 15.6% | 19.4% | 3.1% | 0.7% | 5.3% | 4.1% | 3.8% | 45.9% |
| (99-04) | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 4.8% | 9.7% | 0.2% | 3.2% | 8.0% | 0.6% | 2.3% | 70.7% |

Appendix E.17. Percent distribution of Nooksack Spring Fingerling Chinook reported catch among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1996 | 1.4% | 0.0% | 0.0% | 0.0% | 0.0% | 5.1% | 1.3% | 0.0% | 16.8% | 0.2% | 4.2% | 0.7% | 0.3% | 6.4% | 63.6% |
| 1997 | 3.5% | 0.2% | 0.7% | 0.2% | 0.1% | 0.4% | 0.2% | 1.6% | 10.3% | 0.1% | 2.9% | 0.5% | 1.3% | 5.2% | 73.0% |
| 1998 | 8.1% | 0.2% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 1.7% | 2.9% | 0.0% | 2.3% | 0.2% | 0.1% | 0.6% | 83.6% |
| 1999 | 1.6% | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 1.1% | 3.6% | 0.0% | 5.5% | 1.3% | 0.0% | 0.7% | 84.2% |
| 2000 | 4.6% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 19.5% | 12.6% | 0.0% | 4.6% | 0.2% | 0.2% | 0.4% | 57.6% |
| 2001 | 1.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.9% | 4.5% | 0.0% | 7.4% | 1.0% | 0.8% | 0.7% | 75.2% |
| 2002 | 5.7% | 0.0% | 0.5% | 0.8% | 0.0% | 0.0% | 1.0% | 17.5% | 1.4% | 0.0% | 1.5% | 0.3% | 0.2% | 0.9% | 70.2% |
| 2003 | 2.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 14.1% | 5.7% | 0.0% | 4.4% | 0.0% | 1.3% | 1.9% | 68.7% |
| 2004 | 1.4% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 27.5% | 5.0% | 0.0% | 6.4% | 3.0% | 0.0% | 1.6% | 54.8% |
| (96-98) | 4.3% | 0.1% | 0.2% | 0.1% | 0.0% | 1.9% | 0.5% | 1.1% | 10.0% | 0.1% | 3.1% | 0.5% | 0.6% | 4.1% | 73.4% |
| (99-04) | 2.9% | 0.2% | 0.1% | 0.2% | 0.0% | 0.0% | 0.5% | 14.8% | 5.5% | 0.0% | 5.0% | 1.0% | 0.4% | 1.0% | 68.5% |

Appendix E.18. Percent distribution of Nooksack Spring Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1996 | 3.3% | 0.0% | 0.2% | 0.0% | 0.0% | 5.8% | 1.7% | 0.7% | 18.5% | 0.5% | 4.1% | 0.7% | 0.3% | 9.4% | 54.9% |
| 1997 | 4.0% | 0.4% | 0.8% | 0.3% | 0.0% | 0.4% | 0.2% | 2.0% | 11.4% | 0.9% | 2.9% | 0.6% | 1.3% | 6.4% | 68.4% |
| 1998 | 8.8% | 0.5% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 1.8% | 3.3% | 0.0% | 2.5% | 0.2% | 0.1% | 1.1% | 81.4% |
| 1999 | 2.0% | 2.3% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 1.1% | 4.3% | 0.0% | 5.9% | 1.5% | 0.0% | 1.1% | 80.5% |
| 2000 | 5.3% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 19.9% | 14.3% | 0.0% | 5.3% | 0.2% | 0.2% | 0.7% | 53.8% |
| 2001 | 1.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.0% | 5.3% | 0.0% | 8.3% | 1.2% | 0.8% | 1.6% | 71.9% |
| 2002 | 6.5% | 0.0% | 0.6% | 0.8% | 0.0% | 0.0% | 1.2% | 17.6% | 1.9% | 0.0% | 1.7% | 0.2% | 0.2% | 1.2% | 68.0% |
| 2003 | 3.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.6% | 14.5% | 6.9% | 0.0% | 5.2% | 0.0% | 1.2% | 3.1% | 64.3% |
| 2004 | 1.8% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 27.5% | 6.4% | 0.0% | 7.2% | 3.5% | 0.0% | 2.3% | 50.9% |
| (96-98) | 5.4% | 0.3% | 0.3% | 0.1% | 0.0% | 2.1% | 0.6% | 1.5% | 11.1% | 0.5% | 3.2% | 0.5% | 0.6% | 5.6% | 68.2% |
| (99-04) | 3.4% | 0.4% | 0.1% | 0.2% | 0.0% | 0.0% | 0.7% | 14.9% | 6.5% | 0.0% | 5.6% | 1.1% | 0.4% | 1.7% | 64.9% |

Appendix E.19. Percent distribution of Nooksack Spring Yearling Chinook reported catch among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.9% | 4.7% | 0.0% | 0.0% | 0.0% | 1.6% | 84.8% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.0% | 0.0% | 0.0% | 0.0% | 13.8% | 6.9% | 73.3% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.9% | 0.0% | 0.0% | 14.6% | 9.8% | 0.0% | 2.4% | 4.9% | 34.1% | 29.3% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 2.1% | 32.6% | 5.6% | 7.0% | 2.1% | 8.4% | 5.3% | 36.1% |
| 1992 | 0.4% | 0.4% | 0.0% | 0.0% | 0.9% | 0.6% | 0.4% | 17.4% | 12.3% | 1.1% | 2.3% | 0.9% | 0.4% | 7.8% | 55.3% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 4.4% | 14.7% | 6.0% | 7.6% | 0.8% | 5.3% | 11.5% | 49.2% |
| 1994 | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.1% | 34.2% | 1.0% | 0.0% | 0.2% | 6.3% | 3.3% | 49.3% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 22.8% | 0.0% | 0.0% | 0.0% | 2.9% | 7.0% | 67.3% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 0.0% | 12.4% | 0.0% | 3.2% | 0.5% | 0.0% | 3.2% | 79.6% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 14.2% | 2.7% | 5.3% | 0.0% | 3.5% | 15.9% | 58.4% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.5% | 3.5% | 0.0% | 15.9% | 0.9% | 6.2% | 0.0% | 4.4% | 5.3% | 60.2% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.8% | 25.4% | 0.0% | 1.1% | 2.8% | 5.0% | 1.1% | 61.9% |
| (86-98) | 0.1% | 0.0% | 0.0% | 0.0% | 0.1% | 0.9% | 0.5% | 2.6% | 17.2% | 2.9% | 2.9% | 0.6% | 4.5% | 9.3% | 58.4% |
| (1999) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.8% | 25.4% | 0.0% | 1.1% | 2.8% | 5.0% | 1.1% | 61.9% |

¹ No data are shown for 2000-2004 because of lack of coded-wire tagging of broods from 1997-2000.

Appendix E.20. Percent distribution of Nooksack Spring Yearling Chinook total fishing mortalities among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 2.1% | 11.8% | 4.6% | 0.8% | 0.4% | 8.0% | 3.8% | 68.1% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.1% | 0.0% | 0.0% | 0.0% | 14.5% | 8.9% | 68.5% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 4.2% | 0.0% | 8.5% | 26.8% | 8.5% | 1.4% | 1.4% | 2.8% | 28.2% | 16.9% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 2.4% | 36.9% | 5.7% | 6.8% | 2.4% | 7.7% | 6.8% | 30.7% |
| 1992 | 2.0% | 0.9% | 0.0% | 0.0% | 1.0% | 0.6% | 0.4% | 19.5% | 13.7% | 1.0% | 2.3% | 1.0% | 0.4% | 9.7% | 47.4% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 4.8% | 17.6% | 5.7% | 7.7% | 0.8% | 5.1% | 12.3% | 45.6% |
| 1994 | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.1% | 35.8% | 0.9% | 0.0% | 0.2% | 6.0% | 3.8% | 47.5% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 24.5% | 0.5% | 0.0% | 0.0% | 3.1% | 12.0% | 59.9% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.0% | 0.5% | 14.6% | 0.0% | 3.5% | 0.5% | 0.0% | 5.5% | 74.4% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 15.6% | 2.3% | 5.5% | 0.0% | 3.1% | 21.9% | 51.6% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.9% | 5.4% | 0.0% | 17.1% | 1.6% | 6.2% | 0.0% | 3.9% | 9.3% | 52.7% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.6% | 28.0% | 0.0% | 1.6% | 3.1% | 4.7% | 2.1% | 58.0% |
| (86-98) | 0.2% | 0.1% | 0.0% | 0.0% | 0.3% | 0.9% | 0.6% | 3.9% | 20.2% | 2.8% | 3.1% | 0.6% | 5.0% | 11.1% | 51.2% |
| (1999) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.6% | 28.0% | 0.0% | 1.6% | 3.1% | 4.7% | 2.1% | 58.0% |

¹ No data are shown for 2000-2004 because of lack of coded-wire tagging of broods from 1997-2000.

Appendix E.21. Percent distribution of Skagit Spring Fingerling Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1997 | 1.0% | 0.0% | 0.0% | 0.4% | 0.6% | 1.5% | 0.9% | 1.4% | 8.7% | 0.2% | 4.0% | 0.0% | 1.4% | 7.3% | 72.5% |
| 1998 | 2.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 1.1% | 0.0% | 9.4% | 0.3% | 3.0% | 0.0% | 1.7% | 2.6% | 79.4% |
| 1999 | 0.5% | 0.6% | 0.0% | 0.2% | 0.0% | 0.1% | 0.4% | 0.5% | 4.7% | 0.0% | 5.8% | 0.3% | 1.3% | 1.7% | 83.9% |
| 2000 | 1.5% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.5% | 5.5% | 9.9% | 0.1% | 6.3% | 0.0% | 0.2% | 2.4% | 73.2% |
| 2001 | 1.3% | 0.2% | 0.3% | 0.2% | 0.0% | 0.0% | 1.4% | 4.8% | 6.0% | 0.0% | 6.0% | 0.2% | 0.7% | 4.2% | 74.8% |
| 2002 | 2.7% | 0.0% | 0.5% | 0.2% | 0.0% | 0.1% | 0.6% | 4.7% | 5.6% | 0.0% | 3.1% | 0.3% | 0.6% | 2.7% | 78.8% |
| 2003 | 2.0% | 0.0% | 0.8% | 1.1% | 0.0% | 0.1% | 3.9% | 21.3% | 4.5% | 0.0% | 2.3% | 1.2% | 0.8% | 1.1% | 60.9% |
| 2004 | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 10.4% | 9.0% | 0.0% | 3.0% | 2.3% | 1.3% | 1.5% | 72.0% |
| (97-98) | 1.5% | 0.0% | 0.0% | 0.2% | 0.3% | 1.1% | 1.0% | 0.7% | 9.1% | 0.3% | 3.5% | 0.0% | 1.6% | 4.9% | 75.9% |
| (99-04) | 1.3% | 0.1% | 0.3% | 0.3% | 0.0% | 0.1% | 1.1% | 7.9% | 6.6% | 0.0% | 4.4% | 0.7% | 0.8% | 2.3% | 73.9% |

Appendix E.22. Percent distribution of Skagit Spring Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1997 | 1.2% | 0.0% | 0.0% | 0.4% | 0.5% | 1.8% | 1.2% | 1.6% | 9.9% | 1.1% | 4.3% | 0.0% | 1.3% | 8.9% | 67.5% |
| 1998 | 2.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 1.8% | 0.0% | 10.8% | 0.3% | 3.4% | 0.0% | 1.5% | 6.3% | 73.0% |
| 1999 | 0.9% | 1.4% | 0.0% | 0.2% | 0.0% | 0.1% | 0.6% | 0.6% | 5.9% | 0.0% | 6.3% | 0.4% | 1.3% | 2.8% | 79.6% |
| 2000 | 2.0% | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 0.7% | 5.9% | 11.7% | 0.2% | 7.0% | 0.0% | 0.2% | 5.0% | 66.8% |
| 2001 | 1.7% | 0.3% | 0.3% | 0.2% | 0.0% | 0.0% | 1.7% | 4.6% | 6.8% | 0.0% | 6.5% | 0.1% | 0.6% | 9.2% | 67.8% |
| 2002 | 2.9% | 0.0% | 0.6% | 0.3% | 0.0% | 0.1% | 0.8% | 4.7% | 7.3% | 0.0% | 3.5% | 0.3% | 0.6% | 3.9% | 74.9% |
| 2003 | 2.2% | 0.0% | 0.9% | 1.1% | 0.0% | 0.1% | 5.0% | 21.4% | 5.5% | 0.0% | 2.7% | 1.3% | 0.8% | 1.5% | 57.5% |
| 2004 | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 10.5% | 11.1% | 0.0% | 3.5% | 2.6% | 1.3% | 2.1% | 68.4% |
| (97-98) | 1.7% | 0.0% | 0.0% | 0.2% | 0.3% | 1.3% | 1.5% | 0.8% | 10.3% | 0.7% | 3.8% | 0.0% | 1.4% | 7.6% | 70.3% |
| (99-04) | 1.6% | 0.3% | 0.4% | 0.4% | 0.0% | 0.0% | 1.5% | 7.9% | 8.1% | 0.0% | 4.9% | 0.8% | 0.8% | 4.1% | 69.2% |

Appendix E.23. Percent distribution of Skagit Spring Yearling Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.7% | 29.2% | 26.7% | 0.0% | 0.0% | 10.0% | 15.8% | 11.7% |
| 1986 | 1.4% | 0.0% | 0.0% | 0.0% | 4.3% | 6.6% | 0.0% | 6.2% | 41.7% | 2.8% | 5.7% | 0.0% | 3.3% | 7.6% | 20.4% |
| 1987 | 0.0% | 0.0% | 0.0% | 4.6% | 0.0% | 6.5% | 0.0% | 3.7% | 10.2% | 5.6% | 0.0% | 1.9% | 24.1% | 20.4% | 23.1% |
| 1988 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.9% | 0.0% | 1.8% | 14.9% | 7.7% | 9.6% | 1.8% | 20.6% | 14.5% | 23.2% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 0.1% | 0.0% | 3.4% | 17.5% | 3.3% | 1.8% | 4.3% | 30.4% | 8.4% | 29.9% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 1.9% | 1.0% | 4.9% | 14.0% | 4.0% | 8.7% | 3.4% | 15.4% | 22.9% | 23.3% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 0.0% | 2.0% | 19.6% | 1.6% | 10.2% | 0.0% | 2.4% | 20.9% | 42.2% |
| 1998 | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 3.5% | 1.3% | 9.1% | 0.0% | 7.2% | 0.0% | 3.2% | 17.2% | 57.8% |
| 1999 | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 1.2% | 7.7% | 0.0% | 4.5% | 0.2% | 1.1% | 9.1% | 75.4% |
| 2000 | 0.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 6.4% | 16.1% | 0.0% | 3.6% | 0.0% | 1.5% | 15.3% | 55.8% |
| 2001 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.2% | 11.6% | 0.0% | 2.8% | 3.2% | 2.0% | 10.8% | 66.4% |
| 2002 | 1.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 12.2% | 0.0% | 10.2% | 0.0% | 1.6% | 8.9% | 65.0% |
| 2003 | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 0.0% | 0.6% | 22.7% | 9.0% | 0.0% | 13.1% | 0.1% | 0.7% | 6.2% | 46.9% |
| 2004 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 12.4% | 4.6% | 0.0% | 4.1% | 0.8% | 1.0% | 4.0% | 72.0% |
| (85-98) | 0.2% | 0.0% | 0.0% | 0.6% | 0.7% | 2.8% | 0.6% | 3.7% | 19.5% | 6.4% | 5.4% | 1.4% | 13.7% | 16.0% | 29.0% |
| (99-04) | 0.5% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.4% | 7.8% | 10.2% | 0.0% | 6.4% | 0.7% | 1.3% | 9.1% | 63.6% |

Appendix E.24. Percent distribution of Skagit Spring Yearling Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 0.0% | 6.9% | 29.2% | 24.6% | 0.0% | 0.0% | 9.2% | 18.5% | 10.8% |
| 1986 | 1.8% | 0.0% | 0.0% | 0.0% | 4.0% | 6.6% | 0.0% | 6.2% | 41.6% | 2.7% | 5.8% | 0.0% | 3.1% | 9.3% | 19.0% |
| 1987 | 0.0% | 0.0% | 0.0% | 4.9% | 0.0% | 4.9% | 0.0% | 3.1% | 7.4% | 4.3% | 0.0% | 1.2% | 19.0% | 39.9% | 15.3% |
| 1988 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.5% | 0.0% | 2.4% | 17.6% | 7.1% | 9.3% | 2.1% | 19.5% | 16.2% | 20.3% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 0.1% | 0.0% | 4.0% | 19.5% | 3.3% | 1.9% | 4.7% | 28.2% | 10.4% | 26.9% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 1.9% | 1.1% | 5.1% | 14.8% | 3.7% | 8.6% | 3.7% | 14.6% | 24.6% | 21.6% |
| 1997 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 1.0% | 2.6% | 19.3% | 2.8% | 9.0% | 0.0% | 1.8% | 31.1% | 31.1% |
| 1998 | 0.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 4.0% | 1.2% | 10.1% | 0.2% | 7.1% | 0.0% | 3.0% | 21.1% | 52.4% |
| 1999 | 0.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 1.2% | 8.1% | 0.0% | 4.6% | 0.2% | 1.0% | 12.7% | 71.3% |
| 2000 | 0.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 6.0% | 17.1% | 0.0% | 3.8% | 0.0% | 1.4% | 19.5% | 50.9% |
| 2001 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.8% | 11.0% | 0.0% | 2.8% | 2.8% | 1.6% | 26.7% | 52.2% |
| 2002 | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 15.5% | 0.0% | 10.9% | 0.0% | 1.4% | 14.1% | 56.3% |
| 2003 | 0.0% | 0.0% | 0.0% | 0.8% | 0.0% | 0.0% | 0.7% | 21.8% | 10.7% | 0.0% | 14.8% | 0.1% | 0.6% | 8.4% | 42.2% |
| 2004 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 12.2% | 5.4% | 0.0% | 4.6% | 0.8% | 1.0% | 4.9% | 69.7% |
| (85-98) | 0.3% | 0.0% | 0.0% | 0.6% | 0.7% | 2.6% | 0.8% | 3.9% | 19.9% | 6.1% | 5.2% | 1.5% | 12.3% | 21.4% | 24.7% |
| (99-04) | 0.4% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.5% | 7.5% | 11.3% | 0.0% | 6.9% | 0.6% | 1.2% | 14.4% | 57.1% |

Appendix E.25. Percent distribution of Samish Fall Fingerling Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1989 | 0.0% | 0.0% | 0.0% | 0.2% | 0.2% | 0.2% | 0.3% | 6.8% | 17.2% | 3.5% | 1.9% | 7.4% | 36.2% | 9.7% | 16.5% |
| 1990 | 2.1% | 0.0% | 0.0% | 0.5% | 0.1% | 0.2% | 0.0% | 18.5% | 12.9% | 1.3% | 2.0% | 9.0% | 30.5% | 7.4% | 15.4% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.3% | 0.0% | 13.5% | 11.4% | 2.7% | 3.2% | 8.9% | 23.2% | 10.9% | 25.8% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.5% | 11.4% | 14.6% | 2.1% | 0.9% | 10.2% | 15.6% | 17.2% | 27.4% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.3% | 0.2% | 0.5% | 0.3% | 12.3% | 19.0% | 2.3% | 8.5% | 3.9% | 16.5% | 12.7% | 23.6% |
| 1994 | 0.2% | 0.0% | 0.0% | 0.4% | 0.0% | 0.4% | 0.0% | 11.8% | 13.8% | 1.9% | 5.4% | 2.2% | 38.5% | 3.9% | 21.2% |
| 1995 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 5.8% | 5.1% | 0.3% | 3.4% | 3.4% | 27.2% | 15.0% | 38.8% |
| 1996 | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 10.7% | 0.1% | 0.7% | 1.9% | 33.9% | 24.1% | 28.1% |
| 1997 | 0.5% | 0.2% | 0.0% | 0.3% | 0.7% | 0.8% | 0.3% | 2.0% | 8.2% | 0.1% | 1.8% | 0.9% | 34.5% | 9.8% | 40.0% |
| 1998 | 3.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.7% | 10.9% | 0.0% | 1.7% | 0.7% | 44.2% | 4.1% | 33.3% |
| 1999 | 3.7% | 0.0% | 0.0% | 1.2% | 0.0% | 0.0% | 3.3% | 1.6% | 11.0% | 0.0% | 10.2% | 1.6% | 38.6% | 3.7% | 25.2% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 11.4% | 6.5% | 0.0% | 9.5% | 0.4% | 37.6% | 1.5% | 33.1% |
| 2001 | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.1% | 0.3% | 4.7% | 8.2% | 0.0% | 6.8% | 2.4% | 38.7% | 4.0% | 34.6% |
| 2002 | 0.9% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 8.9% | 7.5% | 0.0% | 4.2% | 2.9% | 37.8% | 5.2% | 32.4% |
| 2003 | 0.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 14.2% | 5.7% | 0.3% | 4.4% | 6.0% | 37.7% | 2.4% | 28.7% |
| 2004 | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.1% | 4.5% | 0.0% | 7.3% | 10.6% | 31.5% | 6.1% | 32.5% |
| (89-98) | 0.6% | 0.0% | 0.0% | 0.2% | 0.1% | 0.4% | 0.1% | 8.4% | 12.4% | 1.4% | 3.0% | 4.9% | 30.0% | 11.5% | 27.0% |
| (99-04) | 1.0% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.6% | 8.0% | 7.2% | 0.0% | 7.1% | 4.0% | 37.0% | 3.8% | 31.1% |

Appendix E.26. Percent distribution of Samish Fall Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1989 | 0.2% | 0.0% | 0.0% | 0.2% | 0.2% | 0.2% | 0.2% | 9.1% | 18.4% | 3.1% | 1.8% | 8.0% | 33.3% | 11.0% | 14.3% |
| 1990 | 2.1% | 0.0% | 0.0% | 0.5% | 0.1% | 0.2% | 0.0% | 19.9% | 13.5% | 1.3% | 2.0% | 9.3% | 28.7% | 8.2% | 14.2% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.4% | 0.0% | 14.6% | 12.4% | 2.5% | 3.2% | 9.4% | 21.7% | 12.1% | 23.6% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.6% | 11.6% | 15.3% | 1.8% | 0.8% | 9.9% | 14.2% | 23.8% | 21.8% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.3% | 0.2% | 0.4% | 0.3% | 14.0% | 21.7% | 2.0% | 8.0% | 4.1% | 15.3% | 13.6% | 20.1% |
| 1994 | 0.5% | 0.0% | 0.0% | 0.5% | 0.0% | 0.4% | 0.0% | 13.1% | 15.1% | 1.9% | 5.5% | 2.1% | 37.0% | 4.6% | 19.3% |
| 1995 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.0% | 7.3% | 5.3% | 0.7% | 3.3% | 3.0% | 24.3% | 22.6% | 32.3% |
| 1996 | 0.0% | 0.1% | 0.0% | 0.1% | 0.0% | 0.4% | 0.0% | 1.0% | 11.4% | 0.2% | 0.7% | 1.7% | 32.6% | 29.1% | 22.9% |
| 1997 | 0.6% | 0.4% | 0.0% | 0.4% | 0.8% | 0.8% | 0.4% | 2.5% | 9.3% | 0.4% | 1.7% | 1.1% | 33.6% | 11.7% | 36.5% |
| 1998 | 3.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.7% | 11.9% | 0.0% | 1.8% | 0.8% | 43.2% | 5.5% | 31.6% |
| 1999 | 4.0% | 0.0% | 0.0% | 1.5% | 0.0% | 0.0% | 3.6% | 1.5% | 12.4% | 0.0% | 10.5% | 1.8% | 36.4% | 5.8% | 22.5% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 10.4% | 6.6% | 0.0% | 9.6% | 0.3% | 40.3% | 6.9% | 26.0% |
| 2001 | 0.0% | 0.7% | 0.0% | 0.0% | 0.0% | 0.2% | 0.3% | 4.3% | 9.5% | 0.0% | 7.1% | 2.6% | 37.1% | 7.6% | 30.5% |
| 2002 | 0.9% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 8.5% | 9.2% | 0.0% | 4.7% | 3.0% | 36.2% | 6.9% | 30.1% |
| 2003 | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 14.2% | 6.8% | 0.6% | 5.1% | 6.3% | 36.5% | 3.0% | 26.8% |
| 2004 | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.9% | 5.5% | 0.0% | 7.8% | 11.8% | 30.1% | 8.2% | 29.1% |
| (89-98) | 0.7% | 0.0% | 0.0% | 0.2% | 0.1% | 0.4% | 0.1% | 9.5% | 13.4% | 1.4% | 2.9% | 4.9% | 28.4% | 14.2% | 23.7% |
| (99-04) | 1.1% | 0.1% | 0.0% | 0.3% | 0.0% | 0.0% | 0.7% | 7.6% | 8.3% | 0.1% | 7.5% | 4.3% | 36.1% | 6.4% | 27.5% |

Appendix E.27. Percent distribution of Skagit Summer Fingerling Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1998 | 3.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.7% | 1.7% | 1.7% | 0.0% | 2.3% | 0.0% | 0.0% | 1.2% | 87.8% |
| 1999 | 7.1% | 2.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.1% | 0.0% | 20.2% | 0.0% | 1.2% | 0.0% | 61.9% |
| 2000 | 5.8% | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.2% | 8.0% | 0.0% | 8.9% | 0.0% | 2.2% | 5.3% | 66.7% |
| 2001 | 6.3% | 6.2% | 0.8% | 0.0% | 0.0% | 0.0% | 1.7% | 6.7% | 9.7% | 0.0% | 8.3% | 0.1% | 0.7% | 1.2% | 58.3% |
| 2002 | 12.9% | 0.0% | 0.9% | 0.9% | 0.0% | 0.1% | 1.5% | 6.4% | 3.3% | 0.2% | 1.1% | 0.1% | 0.9% | 0.0% | 71.7% |
| 2003 | 5.5% | 0.1% | 0.0% | 3.4% | 0.0% | 0.0% | 10.5% | 10.9% | 4.5% | 0.1% | 6.8% | 0.3% | 0.6% | 0.6% | 56.7% |
| 2004 | 5.3% | 0.0% | 0.0% | 2.5% | 0.0% | 0.0% | 1.6% | 10.4% | 1.7% | 0.0% | 1.6% | 0.8% | 1.1% | 0.5% | 74.5% |
| (1998) | 3.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.7% | 1.7% | 1.7% | 0.0% | 2.3% | 0.0% | 0.0% | 1.2% | 87.8% |
| (99-04) | 7.2% | 1.6% | 0.3% | 1.1% | 0.0% | 0.0% | 2.5% | 6.1% | 5.7% | 0.0% | 7.8% | 0.2% | 1.1% | 1.3% | 65.0% |

Appendix E.28. Percent distribution of Skagit Summer Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1998 | 4.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.3% | 1.7% | 2.8% | 0.0% | 2.3% | 0.0% | 0.0% | 1.7% | 85.3% |
| 1999 | 10.1% | 5.1% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 1.5% | 8.1% | 0.0% | 19.2% | 0.0% | 1.0% | 2.0% | 52.5% |
| 2000 | 10.1% | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.2% | 9.0% | 0.0% | 9.4% | 0.0% | 1.8% | 11.5% | 54.0% |
| 2001 | 8.4% | 13.9% | 0.9% | 0.0% | 0.0% | 0.0% | 1.9% | 5.8% | 10.0% | 0.0% | 8.1% | 0.1% | 0.6% | 2.3% | 48.1% |
| 2002 | 13.4% | 0.0% | 0.8% | 1.0% | 0.0% | 0.1% | 1.9% | 6.2% | 4.0% | 2.8% | 1.2% | 0.1% | 0.9% | 0.0% | 67.5% |
| 2003 | 5.8% | 0.5% | 0.0% | 3.5% | 0.0% | 0.0% | 13.0% | 10.6% | 5.1% | 0.2% | 7.8% | 0.3% | 0.6% | 0.7% | 52.0% |
| 2004 | 6.0% | 0.0% | 0.0% | 3.1% | 0.0% | 0.0% | 2.3% | 10.6% | 2.3% | 0.0% | 1.8% | 0.9% | 1.0% | 0.8% | 71.3% |
| (1998) | 4.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.3% | 1.7% | 2.8% | 0.0% | 2.3% | 0.0% | 0.0% | 1.7% | 85.3% |
| (99-04) | 9.0% | 3.4% | 0.4% | 1.3% | 0.0% | 0.0% | 3.2% | 6.3% | 6.4% | 0.5% | 7.9% | 0.2% | 1.0% | 2.9% | 57.6% |

Appendix E.29. Percent distribution of Stillaguamish Fall Fingerling Chinook reported catch among fisheries and escapement (NA=not available).

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|-----------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1984 | 0.0% | 0.0% | 0.0% | 3.6% | 19.3% | 2.4% | 3.6% | 7.2% | 15.7% | 24.1% | 0.0% | 0.0% | 4.8% | 19.3% | NA ² |
| 1985 | 7.3% | 0.0% | 0.0% | 4.2% | 0.0% | 4.2% | 0.0% | 30.2% | 10.4% | 11.5% | 9.4% | 0.0% | 9.4% | 13.5% | NA ² |
| 1986 | 4.2% | 0.0% | 0.0% | 0.0% | 0.0% | 4.2% | 0.0% | 30.2% | 18.8% | 0.0% | 0.0% | 0.0% | 15.6% | 19.8% | NA ² |
| 1990 | 0.4% | 0.0% | 0.0% | 0.7% | 6.2% | 4.0% | 0.0% | 16.2% | 7.5% | 4.2% | 4.9% | 4.2% | 7.1% | 10.4% | 13.3% |
| 1991 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.4% | 5.9% | 4.4% | 0.9% | 2.6% | 5.1% | 6.9% | 7.9% | 68.6% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 2.4% | 0.0% | 17.0% | 5.1% | 2.5% | 4.0% | 5.7% | 11.9% | 28.1% | 22.5% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.6% | 0.4% | 1.0% | 1.4% | 11.5% | 8.7% | 1.4% | 9.6% | 5.4% | 1.5% | 22.5% | 38.1% |
| 1994 | 2.4% | 0.0% | 0.0% | 0.7% | 0.0% | 1.3% | 0.0% | 6.7% | 7.8% | 0.9% | 5.3% | 0.0% | 2.4% | 5.8% | 66.8% |
| 1995 | 2.3% | 0.0% | 0.0% | 0.0% | 0.0% | 9.8% | 0.0% | 2.3% | 4.1% | 1.0% | 9.6% | 1.0% | 2.3% | 13.7% | 52.9% |
| 1996 | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% | 8.3% | 1.4% | 0.0% | 6.3% | 0.6% | 7.6% | 0.0% | 0.3% | 19.4% | 57.7% |
| 1997 | 8.3% | 0.4% | 0.0% | 0.5% | 0.0% | 1.3% | 1.0% | 6.5% | 4.5% | 0.0% | 4.7% | 0.0% | 1.8% | 14.3% | 53.2% |
| 1998 | 12.7% | 0.3% | 0.4% | 1.2% | 0.0% | 0.0% | 0.8% | 1.3% | 2.2% | 0.1% | 2.9% | 0.0% | 2.4% | 2.5% | 80.2% |
| 1999 | 0.9% | 2.2% | 0.0% | 0.0% | 0.0% | 0.7% | 0.4% | 1.5% | 7.9% | 0.0% | 10.6% | 0.0% | 0.4% | 3.5% | 79.8% |
| 2000 | 4.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.6% | 2.0% | 0.0% | 1.5% | 0.5% | 0.1% | 1.6% | 80.7% |
| 2001 | 2.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.2% | 5.6% | 0.0% | 4.9% | 0.3% | 1.4% | 10.5% | 70.8% |
| (90-98) | 2.8% | 0.1% | 0.0% | 0.5% | 0.9% | 3.2% | 0.5% | 8.0% | 5.7% | 1.4% | 5.7% | 2.5% | 4.2% | 14.0% | 50.4% |
| (99-01) | 2.5% | 0.7% | 0.0% | 0.0% | 0.0% | 0.2% | 0.1% | 4.4% | 5.2% | 0.0% | 5.7% | 0.3% | 0.6% | 5.2% | 77.1% |

Appendix E.30. Percent distribution of Stillaguamish Fall Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|-----------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1984 | 0.9% | 0.0% | 0.0% | 3.7% | 16.8% | 1.9% | 2.8% | 10.3% | 13.1% | 19.6% | 0.0% | 0.0% | 4.7% | 26.2% | NA ² |
| 1985 | 7.1% | 0.0% | 0.0% | 4.5% | 0.0% | 3.6% | 0.0% | 31.3% | 8.9% | 9.8% | 8.9% | 0.0% | 8.0% | 17.9% | NA ² |
| 1986 | 5.7% | 0.0% | 0.0% | 0.0% | 0.0% | 3.8% | 0.0% | 29.5% | 19.0% | 0.0% | 0.0% | 0.0% | 14.3% | 21.0% | NA ² |
| 1990 | 0.6% | 0.0% | 0.0% | 0.8% | 6.1% | 3.6% | 0.0% | 17.0% | 8.4% | 3.8% | 4.8% | 5.1% | 7.4% | 13.0% | 11.0% |
| 1991 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.5% | 6.8% | 5.1% | 1.0% | 2.6% | 5.9% | 6.9% | 10.0% | 64.2% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 2.1% | 0.0% | 16.6% | 4.9% | 2.0% | 3.4% | 5.3% | 10.4% | 38.7% | 16.3% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.9% | 0.5% | 1.0% | 1.3% | 13.8% | 10.0% | 1.3% | 9.1% | 5.9% | 1.4% | 23.5% | 33.5% |
| 1994 | 2.9% | 0.0% | 0.0% | 0.6% | 0.0% | 1.3% | 0.0% | 7.3% | 8.6% | 1.0% | 5.7% | 0.0% | 2.3% | 7.1% | 63.2% |
| 1995 | 2.4% | 0.0% | 0.0% | 0.0% | 0.0% | 10.7% | 0.0% | 3.8% | 4.4% | 1.8% | 8.9% | 0.8% | 2.2% | 24.2% | 40.1% |
| 1996 | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 9.2% | 2.1% | 1.1% | 6.7% | 0.6% | 7.3% | 0.0% | 0.3% | 26.0% | 47.5% |
| 1997 | 9.0% | 0.8% | 0.0% | 0.4% | 0.0% | 1.4% | 1.2% | 7.2% | 4.7% | 0.3% | 4.6% | 0.0% | 1.7% | 17.2% | 47.9% |
| 1998 | 14.0% | 1.0% | 0.5% | 2.2% | 0.0% | 0.0% | 1.1% | 1.2% | 2.4% | 0.1% | 3.0% | 0.0% | 2.3% | 4.0% | 76.5% |
| 1999 | 1.0% | 9.3% | 0.0% | 0.0% | 0.0% | 0.6% | 0.4% | 1.4% | 8.6% | 0.0% | 10.3% | 0.0% | 0.4% | 4.9% | 73.0% |
| 2000 | 5.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.8% | 2.3% | 0.0% | 1.7% | 0.5% | 0.1% | 2.6% | 77.9% |
| 2001 | 1.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.1% | 5.5% | 0.0% | 5.1% | 0.3% | 1.3% | 16.4% | 65.6% |
| (90-98) | 3.0% | 0.2% | 0.0% | 0.5% | 0.9% | 3.3% | 0.7% | 8.8% | 6.2% | 1.4% | 5.5% | 2.7% | 3.9% | 18.3% | 44.5% |
| (99-01) | 2.7% | 3.1% | 0.0% | 0.0% | 0.0% | 0.2% | 0.1% | 4.4% | 5.5% | 0.0% | 5.7% | 0.3% | 0.6% | 8.0% | 72.2% |

¹ No data are shown for 2002-2004 because of lack of coded-wire tagging of broods from 1999-2000.

² Values represent estimates of catch or total fishing mortality distribution only for this year.

Appendix E.31. Percent distribution of Nisqually Fall Fingerling Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|-----------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1983 | 0.0% | 0.0% | 0.0% | 2.5% | 0.0% | 0.0% | 0.0% | 16.6% | 12.6% | 6.1% | 0.0% | 4.6% | 11.1% | 46.5% | NA ¹ |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 30.7% | 1.6% | 2.7% | 0.0% | 1.6% | 40.4% | 23.1% | NA ¹ |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 31.7% | 0.0% | 6.4% | 3.1% | 8.0% | 33.3% | 17.5% | NA ¹ |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 15.7% | 13.0% | 1.7% | 0.0% | 0.0% | 35.7% | 14.8% | 19.1% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 1.3% | 0.0% | 10.7% | 13.3% | 0.7% | 0.0% | 5.3% | 35.3% | 18.7% | 12.7% |
| 1988 | 0.0% | 0.0% | 0.0% | 0.7% | 2.2% | 0.7% | 2.2% | 5.4% | 17.7% | 4.7% | 0.0% | 8.7% | 17.3% | 10.5% | 30.0% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 0.7% | 0.0% | 4.4% | 2.5% | 3.6% | 6.3% | 13.3% | 42.6% | 18.3% | 8.0% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 22.5% | 3.1% | 0.2% | 5.8% | 10.2% | 37.7% | 12.2% | 8.2% |
| 1991 | 0.0% | 0.0% | 0.0% | 2.1% | 0.0% | 0.0% | 0.0% | 8.2% | 3.3% | 2.5% | 2.1% | 16.5% | 23.0% | 24.3% | 18.1% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.8% | 7.6% | 2.9% | 2.6% | 4.2% | 7.6% | 18.2% | 16.7% | 39.3% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.0% | 0.0% | 12.4% | 3.9% | 2.2% | 1.8% | 2.9% | 22.4% | 19.2% | 34.3% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 4.5% | 2.4% | 2.4% | 0.5% | 0.8% | 22.0% | 21.2% | 46.2% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.2% | 5.4% | 1.7% | 0.1% | 3.1% | 2.7% | 32.4% | 24.4% | 29.7% |
| 1996 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 1.0% | 0.0% | 0.0% | 3.3% | 0.0% | 1.1% | 1.7% | 42.0% | 21.3% | 29.4% |
| 1997 | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.3% | 0.6% | 2.4% | 0.6% | 0.0% | 4.5% | 0.8% | 18.9% | 24.4% | 47.0% |
| 1998 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.5% | 1.5% | 0.0% | 0.7% | 0.5% | 36.4% | 12.0% | 47.9% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 2.9% | 0.0% | 2.7% | 2.8% | 43.9% | 19.6% | 27.7% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 13.4% | 3.2% | 0.0% | 5.6% | 1.7% | 44.9% | 17.5% | 13.7% |
| 2001 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.1% | 1.9% | 0.0% | 3.8% | 4.2% | 29.2% | 15.6% | 42.0% |
| 2002 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.0% | 1.0% | 0.0% | 2.2% | 3.5% | 42.6% | 11.1% | 32.7% |
| 2003 | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 5.8% | 1.3% | 0.0% | 5.3% | 4.1% | 42.1% | 12.4% | 28.5% |
| 2004 | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.7% | 1.2% | 0.0% | 1.3% | 6.6% | 31.9% | 8.6% | 44.7% |
| (86-98) | 0.0% | 0.0% | 0.0% | 0.2% | 0.3% | 0.4% | 0.3% | 7.7% | 5.3% | 1.6% | 2.3% | 5.5% | 29.5% | 18.3% | 28.5% |
| (99-04) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 5.9% | 1.9% | 0.0% | 3.5% | 3.8% | 39.1% | 14.1% | 31.5% |

¹ Values represent estimates of catch distribution only for this year.

Appendix E.32. Percent distribution of Nisqually Fall Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|-----------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1983 | 0.0% | 0.0% | 0.0% | 1.7% | 0.0% | 0.0% | 0.0% | 15.1% | 8.9% | 4.8% | 0.0% | 3.1% | 9.3% | 57.0% | NA ¹ |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 30.3% | 1.4% | 2.6% | 0.0% | 1.8% | 37.1% | 26.8% | NA ¹ |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 29.6% | 0.0% | 5.0% | 3.7% | 7.4% | 32.1% | 22.2% | NA ¹ |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.4% | 12.5% | 1.6% | 0.0% | 0.0% | 32.8% | 19.5% | 17.2% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.0% | 2.7% | 1.1% | 0.0% | 14.4% | 11.8% | 0.5% | 0.0% | 5.9% | 29.9% | 23.5% | 10.2% |
| 1988 | 0.0% | 0.0% | 0.0% | 0.8% | 2.1% | 0.8% | 2.6% | 5.8% | 18.6% | 3.7% | 0.0% | 8.1% | 16.0% | 19.7% | 21.8% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.6% | 0.0% | 5.4% | 3.0% | 3.2% | 6.0% | 14.6% | 40.4% | 19.1% | 7.2% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 23.4% | 3.2% | 0.1% | 5.9% | 10.4% | 35.6% | 13.6% | 7.6% |
| 1991 | 0.0% | 0.0% | 0.0% | 2.2% | 0.0% | 0.0% | 0.0% | 9.1% | 3.6% | 2.2% | 1.8% | 17.2% | 21.2% | 26.6% | 16.1% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 1.0% | 7.2% | 2.9% | 1.9% | 3.7% | 7.0% | 18.4% | 28.5% | 29.3% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.0% | 14.7% | 4.5% | 2.0% | 1.7% | 3.2% | 21.6% | 21.8% | 29.6% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 4.2% | 2.3% | 2.4% | 0.4% | 0.6% | 17.8% | 39.9% | 32.3% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.3% | 8.0% | 2.0% | 0.3% | 3.0% | 2.4% | 30.3% | 27.7% | 25.8% |
| 1996 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 0.0% | 0.7% | 3.6% | 0.0% | 1.2% | 1.6% | 38.9% | 26.3% | 26.4% |
| 1997 | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 0.3% | 0.8% | 2.8% | 0.7% | 0.3% | 4.3% | 0.8% | 17.4% | 31.9% | 40.2% |
| 1998 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.4% | 1.5% | 0.0% | 0.7% | 0.5% | 31.5% | 26.4% | 38.3% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 3.3% | 0.0% | 2.6% | 3.1% | 41.8% | 24.1% | 24.6% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 11.9% | 3.1% | 0.0% | 5.6% | 1.6% | 36.9% | 29.9% | 11.0% |
| 2001 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.8% | 2.1% | 0.0% | 3.8% | 4.4% | 26.0% | 26.1% | 34.6% |
| 2002 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.8% | 1.2% | 0.0% | 2.4% | 3.8% | 40.7% | 15.6% | 29.5% |
| 2003 | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 5.5% | 1.5% | 0.0% | 5.9% | 4.4% | 39.6% | 16.6% | 25.6% |
| 2004 | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.7% | 1.5% | 0.0% | 1.4% | 7.2% | 29.9% | 14.7% | 39.5% |
| (86-98) | 0.0% | 0.0% | 0.0% | 0.3% | 0.4% | 0.4% | 0.4% | 8.7% | 5.4% | 1.4% | 2.2% | 5.6% | 27.1% | 25.0% | 23.2% |
| (99-04) | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 5.5% | 2.1% | 0.0% | 3.6% | 4.1% | 35.8% | 21.2% | 27.5% |

¹ Values represent estimates of fishing mortality distribution only for this year.

Appendix E.33. Percent distribution of George Adams Fall Fingerling Chinook among fisheries reported catch and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1982 | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.3% | 0.0% | 20.8% | 4.4% | 0.4% | 0.0% | 3.0% | 38.1% | 10.7% | 21.9% |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 1.6% | 1.6% | 0.0% | 15.7% | 3.5% | 4.2% | 0.5% | 0.2% | 29.8% | 25.8% | 17.2% |
| 1984 | 0.0% | 0.1% | 0.0% | 0.5% | 3.2% | 0.7% | 0.4% | 18.1% | 5.7% | 1.2% | 0.0% | 2.2% | 31.3% | 20.6% | 15.9% |
| 1989 | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 8.5% | 3.8% | 4.6% | 1.7% | 12.9% | 38.6% | 17.2% | 12.2% |
| 1990 | 0.1% | 0.0% | 0.0% | 0.4% | 0.3% | 0.5% | 0.0% | 19.3% | 4.7% | 1.0% | 5.0% | 15.0% | 28.4% | 18.4% | 6.8% |
| 1991 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 18.4% | 2.2% | 0.4% | 4.5% | 8.6% | 33.3% | 18.0% | 14.4% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 15.6% | 2.1% | 5.2% | 0.0% | 20.3% | 9.4% | 39.6% | 7.3% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 33.9% | 4.3% | 0.0% | 7.8% | 8.7% | 4.3% | 22.6% | 18.3% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.0% | 0.0% | 0.0% | 0.0% | 14.0% | 7.0% | 72.1% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 0.0% | 7.9% | 3.9% | 0.5% | 3.9% | 1.0% | 4.4% | 18.7% | 57.6% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.4% | 0.0% | 0.0% | 12.6% | 0.0% | 4.7% | 5.9% | 0.0% | 13.8% | 60.6% |
| 1997 | 1.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.2% | 3.0% | 0.3% | 1.4% | 3.0% | 0.8% | 18.8% | 66.5% |
| 1998 | 0.7% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.7% | 0.0% | 1.1% | 1.8% | 1.8% | 7.2% | 86.4% |
| 1999 | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 2.5% | 0.0% | 9.0% | 4.9% | 2.9% | 10.9% | 68.5% |
| 2000 | 0.4% | 0.0% | 0.0% | 0.2% | 0.0% | 0.1% | 0.0% | 18.9% | 3.4% | 0.0% | 10.6% | 3.5% | 0.4% | 17.5% | 45.1% |
| 2001 | 0.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 11.7% | 2.9% | 0.0% | 2.7% | 6.3% | 10.9% | 9.9% | 54.4% |
| 2002 | 1.5% | 0.0% | 0.0% | 1.0% | 0.0% | 0.0% | 0.0% | 11.4% | 1.8% | 0.0% | 7.0% | 4.2% | 11.5% | 15.5% | 46.0% |
| 2003 | 0.5% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 11.6% | 2.4% | 0.0% | 3.4% | 6.4% | 10.1% | 18.0% | 47.5% |
| 2004 | 0.5% | 0.1% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 14.8% | 1.9% | 0.0% | 3.5% | 6.0% | 11.5% | 7.1% | 54.4% |
| (82-84) | 0.0% | 0.0% | 0.0% | 0.2% | 1.7% | 0.8% | 0.1% | 18.2% | 4.5% | 1.9% | 0.2% | 1.8% | 33.1% | 19.0% | 18.4% |
| (89-98) | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 10.8% | 4.4% | 1.2% | 3.0% | 7.7% | 13.5% | 18.1% | 40.2% |
| (99-04) | 0.7% | 0.0% | 0.0% | 0.2% | 0.0% | 0.1% | 0.1% | 11.5% | 2.5% | 0.0% | 6.0% | 5.2% | 7.9% | 13.1% | 52.6% |

Appendix E.34. Percent distribution of George Adams Fall Fingerling Chinook total fishing among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1982 | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.3% | 0.0% | 21.6% | 4.3% | 0.5% | 0.0% | 2.9% | 36.7% | 12.8% | 20.3% |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 1.1% | 0.0% | 12.6% | 2.4% | 3.1% | 0.3% | 0.1% | 25.7% | 42.4% | 11.0% |
| 1984 | 0.0% | 0.1% | 0.0% | 0.6% | 3.2% | 0.7% | 0.5% | 18.2% | 5.6% | 1.1% | 0.0% | 2.3% | 30.6% | 22.5% | 14.6% |
| 1989 | 0.0% | 0.7% | 0.0% | 0.1% | 0.1% | 0.3% | 0.0% | 10.2% | 3.9% | 4.0% | 1.8% | 13.1% | 35.6% | 19.9% | 10.3% |
| 1990 | 0.8% | 0.0% | 0.0% | 0.5% | 0.4% | 0.5% | 0.0% | 21.2% | 4.9% | 1.0% | 4.6% | 15.5% | 25.9% | 18.9% | 5.9% |
| 1991 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 19.4% | 2.3% | 0.4% | 4.5% | 8.7% | 31.6% | 19.7% | 13.3% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 16.6% | 1.8% | 4.6% | 0.0% | 20.3% | 8.3% | 41.5% | 6.5% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 33.6% | 5.1% | 0.0% | 7.3% | 8.0% | 4.4% | 26.3% | 15.3% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.3% | 0.0% | 0.0% | 0.0% | 16.7% | 10.4% | 64.6% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.3% | 0.0% | 9.7% | 4.3% | 1.2% | 3.9% | 0.8% | 4.3% | 28.3% | 45.3% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.7% | 0.0% | 1.3% | 14.3% | 0.0% | 4.6% | 5.7% | 0.0% | 15.9% | 55.5% |
| 1997 | 2.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.5% | 3.0% | 0.8% | 1.3% | 3.0% | 0.8% | 24.2% | 60.5% |
| 1998 | 0.7% | 0.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.9% | 0.0% | 1.2% | 1.7% | 2.0% | 27.0% | 65.6% |
| 1999 | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 3.1% | 0.0% | 9.1% | 5.8% | 2.8% | 14.3% | 63.5% |
| 2000 | 0.4% | 0.0% | 0.0% | 0.2% | 0.0% | 0.2% | 0.0% | 18.4% | 3.6% | 0.0% | 11.3% | 3.5% | 0.3% | 23.0% | 39.0% |
| 2001 | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 11.0% | 3.3% | 0.0% | 2.8% | 6.9% | 10.4% | 16.7% | 47.4% |
| 2002 | 1.8% | 0.0% | 0.0% | 1.1% | 0.0% | 0.0% | 0.0% | 11.1% | 2.3% | 0.0% | 8.0% | 4.5% | 11.2% | 18.4% | 41.7% |
| 2003 | 0.6% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 11.3% | 2.8% | 0.0% | 3.9% | 6.8% | 9.7% | 21.7% | 42.6% |
| 2004 | 0.6% | 0.2% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 14.7% | 2.2% | 0.0% | 3.8% | 6.6% | 11.6% | 10.7% | 49.1% |
| (82-84) | 0.0% | 0.0% | 0.0% | 0.2% | 1.7% | 0.7% | 0.2% | 17.4% | 4.1% | 1.6% | 0.1% | 1.8% | 31.0% | 25.9% | 15.3% |
| (89-98) | 0.4% | 0.1% | 0.0% | 0.1% | 0.0% | 0.6% | 0.0% | 11.7% | 4.9% | 1.2% | 2.9% | 7.7% | 12.9% | 23.2% | 34.3% |
| (99-04) | 0.8% | 0.1% | 0.0% | 0.2% | 0.0% | 0.1% | 0.1% | 11.2% | 2.9% | 0.0% | 6.5% | 5.7% | 7.7% | 17.5% | 47.2% |

Appendix E.35. Percent distribution of South Puget Sound Fall Fingerling Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1982 | 0.2% | 0.0% | 0.0% | 0.1% | 0.8% | 0.4% | 0.1% | 23.0% | 13.8% | 1.6% | 0.1% | 2.8% | 24.8% | 21.3% | 10.9% |
| 1983 | 0.1% | 0.0% | 0.0% | 0.7% | 1.8% | 0.6% | 0.1% | 17.3% | 4.6% | 2.6% | 0.3% | 1.6% | 27.4% | 28.6% | 14.3% |
| 1984 | 0.1% | 0.2% | 0.0% | 0.7% | 1.4% | 0.2% | 0.1% | 20.5% | 8.5% | 1.0% | 0.3% | 1.4% | 24.6% | 22.5% | 18.5% |
| 1985 | 0.8% | 0.0% | 0.0% | 0.0% | 0.3% | 0.4% | 0.2% | 18.7% | 6.3% | 1.6% | 0.8% | 1.9% | 29.3% | 18.2% | 21.6% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 18.4% | 7.5% | 1.7% | 0.0% | 4.0% | 10.7% | 22.4% | 34.0% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 12.7% | 12.7% | 3.9% | 0.0% | 7.2% | 13.9% | 10.9% | 38.8% |
| 1988 | 0.1% | 0.0% | 0.0% | 0.2% | 0.5% | 0.8% | 0.5% | 5.5% | 7.5% | 3.8% | 4.2% | 7.1% | 26.4% | 14.7% | 28.7% |
| 1989 | 0.1% | 0.0% | 0.0% | 0.2% | 0.3% | 0.1% | 0.0% | 7.4% | 4.5% | 3.9% | 2.5% | 11.0% | 21.4% | 16.1% | 32.3% |
| 1990 | 0.0% | 0.0% | 0.1% | 0.3% | 0.3% | 0.3% | 0.0% | 22.7% | 3.6% | 1.0% | 4.3% | 9.0% | 23.7% | 12.5% | 22.3% |
| 1991 | 0.4% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 15.1% | 1.8% | 1.0% | 2.6% | 11.6% | 26.5% | 13.1% | 27.7% |
| 1992 | 0.6% | 0.1% | 0.0% | 0.0% | 0.9% | 0.5% | 0.0% | 17.2% | 3.7% | 2.5% | 2.2% | 9.1% | 23.7% | 18.0% | 21.5% |
| 1993 | 0.2% | 0.1% | 0.0% | 0.0% | 0.1% | 0.6% | 0.0% | 15.7% | 3.8% | 2.2% | 4.6% | 5.5% | 15.7% | 21.0% | 30.4% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.2% | 0.0% | 8.9% | 3.0% | 4.1% | 1.3% | 0.7% | 16.3% | 10.0% | 55.0% |
| 1995 | 0.2% | 0.0% | 0.0% | 0.1% | 0.0% | 0.9% | 0.0% | 3.7% | 1.8% | 0.2% | 1.1% | 1.3% | 5.6% | 11.7% | 73.4% |
| 1996 | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.1% | 0.0% | 4.1% | 0.1% | 1.8% | 2.9% | 6.3% | 14.8% | 69.4% |
| 1997 | 0.5% | 0.0% | 0.0% | 0.3% | 0.0% | 0.5% | 0.0% | 5.2% | 1.8% | 0.0% | 1.5% | 1.6% | 2.9% | 13.2% | 72.5% |
| 1998 | 1.3% | 0.0% | 0.0% | 0.9% | 0.0% | 0.0% | 0.2% | 0.5% | 1.7% | 0.0% | 0.8% | 1.0% | 8.0% | 6.3% | 79.3% |
| 1999 | 0.5% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.7% | 2.4% | 0.0% | 4.0% | 3.0% | 9.2% | 5.3% | 74.8% |
| 2000 | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.1% | 1.9% | 0.0% | 4.1% | 0.3% | 12.2% | 6.7% | 65.3% |
| 2001 | 0.1% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 7.5% | 3.4% | 0.0% | 4.5% | 4.1% | 11.3% | 8.9% | 59.8% |
| 2002 | 0.8% | 0.0% | 0.0% | 0.5% | 0.0% | 0.1% | 0.2% | 11.5% | 3.7% | 0.0% | 2.1% | 4.1% | 18.7% | 7.0% | 51.4% |
| 2003 | 0.6% | 0.0% | 0.0% | 0.7% | 0.0% | 0.0% | 0.0% | 13.2% | 3.4% | 0.0% | 12.0% | 4.4% | 12.8% | 9.5% | 43.3% |
| 2004 | 0.4% | 0.1% | 0.0% | 0.6% | 0.0% | 0.0% | 0.2% | 17.2% | 2.0% | 0.0% | 4.9% | 10.0% | 14.5% | 10.6% | 39.7% |
| (82-84) | 0.2% | 0.1% | 0.0% | 0.5% | 1.4% | 0.4% | 0.1% | 20.3% | 8.9% | 1.8% | 0.2% | 2.0% | 25.6% | 24.1% | 14.5% |
| (85-98) | 0.3% | 0.0% | 0.0% | 0.2% | 0.2% | 0.4% | 0.1% | 10.8% | 4.6% | 1.9% | 2.0% | 5.3% | 16.5% | 14.5% | 43.3% |
| (99-04) | 0.4% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.1% | 9.9% | 2.8% | 0.0% | 5.3% | 4.3% | 13.1% | 8.0% | 55.7% |

Appendix E.36. Percent distribution of South Puget Sound Fall Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1982 | 0.2% | 0.0% | 0.0% | 0.2% | 1.0% | 0.3% | 0.1% | 24.6% | 12.5% | 1.5% | 0.1% | 2.7% | 23.2% | 24.1% | 9.3% |
| 1983 | 0.1% | 0.0% | 0.0% | 0.7% | 1.8% | 0.5% | 0.1% | 16.8% | 3.9% | 2.3% | 0.2% | 1.6% | 25.3% | 35.6% | 11.2% |
| 1984 | 0.1% | 0.2% | 0.0% | 0.7% | 1.4% | 0.2% | 0.1% | 20.8% | 8.3% | 0.9% | 0.3% | 1.5% | 23.9% | 24.8% | 16.9% |
| 1985 | 0.8% | 0.0% | 0.0% | 0.0% | 0.3% | 0.3% | 0.2% | 18.6% | 6.2% | 1.6% | 0.9% | 1.9% | 28.4% | 20.7% | 20.2% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 18.5% | 7.1% | 1.6% | 0.0% | 4.0% | 9.9% | 28.0% | 29.6% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 21.3% | 10.5% | 3.4% | 0.0% | 8.9% | 11.8% | 15.3% | 28.8% |
| 1988 | 0.4% | 0.0% | 0.0% | 0.2% | 1.0% | 0.6% | 0.4% | 10.2% | 9.3% | 3.0% | 3.3% | 7.6% | 22.1% | 22.6% | 19.3% |
| 1989 | 0.1% | 0.0% | 0.0% | 0.3% | 0.4% | 0.1% | 0.0% | 8.8% | 5.2% | 3.6% | 2.4% | 12.2% | 20.5% | 17.4% | 28.9% |
| 1990 | 0.0% | 0.1% | 0.1% | 0.3% | 0.3% | 0.3% | 0.0% | 23.9% | 3.8% | 0.9% | 4.3% | 9.2% | 22.4% | 13.9% | 20.5% |
| 1991 | 0.5% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 16.5% | 1.9% | 0.9% | 2.6% | 12.3% | 25.2% | 14.5% | 25.4% |
| 1992 | 0.6% | 0.2% | 0.0% | 0.0% | 0.9% | 0.5% | 0.0% | 17.4% | 3.8% | 2.4% | 2.1% | 9.1% | 21.1% | 24.0% | 17.9% |
| 1993 | 0.3% | 0.1% | 0.0% | 0.0% | 0.1% | 0.6% | 0.0% | 18.2% | 4.5% | 2.0% | 4.3% | 5.9% | 14.7% | 22.7% | 26.5% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.2% | 0.0% | 9.4% | 3.3% | 4.8% | 1.3% | 0.6% | 15.5% | 17.4% | 46.9% |
| 1995 | 0.2% | 0.0% | 0.0% | 0.1% | 0.0% | 1.1% | 0.0% | 5.4% | 2.1% | 0.7% | 1.2% | 1.3% | 5.8% | 17.3% | 64.9% |
| 1996 | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.2% | 0.9% | 4.8% | 0.2% | 1.8% | 2.8% | 6.3% | 17.9% | 64.5% |
| 1997 | 0.5% | 0.0% | 0.0% | 0.3% | 0.0% | 0.6% | 0.0% | 6.2% | 2.0% | 0.3% | 1.5% | 1.7% | 2.8% | 16.3% | 67.7% |
| 1998 | 1.4% | 0.0% | 0.0% | 0.9% | 0.0% | 0.0% | 0.3% | 0.5% | 1.8% | 0.0% | 0.8% | 1.1% | 8.0% | 11.9% | 73.2% |
| 1999 | 0.6% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.7% | 3.0% | 0.0% | 4.3% | 3.5% | 9.3% | 7.9% | 70.5% |
| 2000 | 0.4% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.0% | 2.2% | 0.0% | 4.6% | 0.3% | 11.8% | 13.9% | 57.5% |
| 2001 | 0.1% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 7.2% | 4.0% | 0.0% | 4.7% | 4.6% | 10.8% | 14.0% | 54.0% |
| 2002 | 0.9% | 0.0% | 0.0% | 0.5% | 0.0% | 0.2% | 0.3% | 11.3% | 4.7% | 0.0% | 2.4% | 4.5% | 17.9% | 9.9% | 47.5% |
| 2003 | 0.6% | 0.0% | 0.0% | 0.8% | 0.0% | 0.0% | 0.0% | 12.6% | 4.1% | 0.0% | 13.5% | 4.7% | 12.2% | 12.1% | 39.4% |
| 2004 | 0.4% | 0.2% | 0.0% | 0.6% | 0.0% | 0.0% | 0.2% | 16.5% | 2.3% | 0.0% | 5.2% | 10.4% | 13.4% | 15.7% | 34.9% |
| (82-84) | 0.2% | 0.1% | 0.0% | 0.6% | 1.4% | 0.3% | 0.1% | 20.7% | 8.2% | 1.6% | 0.2% | 2.0% | 24.1% | 28.2% | 12.4% |
| (85-98) | 0.4% | 0.0% | 0.0% | 0.2% | 0.2% | 0.4% | 0.1% | 12.6% | 4.7% | 1.8% | 1.9% | 5.6% | 15.3% | 18.6% | 38.2% |
| (99-04) | 0.5% | 0.1% | 0.0% | 0.3% | 0.0% | 0.0% | 0.2% | 9.5% | 3.4% | 0.0% | 5.8% | 4.7% | 12.6% | 12.3% | 50.6% |

Appendix E.37. Percent distribution of South Puget Sound Fall Yearling Chinook reported catch among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1982 | 0.0% | 0.0% | 0.0% | 0.0% | 2.5% | 0.0% | 0.0% | 2.8% | 3.2% | 0.0% | 0.0% | 1.1% | 14.5% | 67.5% | 8.5% |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 1.8% | 0.0% | 0.0% | 5.8% | 0.5% | 0.0% | 0.0% | 0.0% | 9.8% | 76.3% | 5.8% |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.3% | 1.6% | 0.0% | 0.0% | 0.0% | 33.6% | 43.3% | 14.2% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.1% | 0.0% | 0.3% | 0.0% | 0.5% | 0.0% | 1.4% | 32.3% | 54.7% | 10.6% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.6% | 0.7% | 0.0% | 0.0% | 3.7% | 12.8% | 57.6% | 19.6% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.6% | 0.8% | 0.0% | 1.2% | 4.6% | 28.5% | 49.1% | 11.2% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 1.1% | 0.0% | 0.0% | 1.4% | 10.4% | 57.7% | 28.0% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 0.5% | 2.2% | 0.7% | 0.0% | 15.6% | 63.3% | 16.9% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.4% | 2.6% | 0.0% | 2.0% | 0.4% | 10.4% | 68.2% | 10.0% |
| 1996 | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.7% | 0.0% | 1.3% | 0.7% | 3.2% | 89.3% | 3.3% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.5% | 1.1% | 0.0% | 0.4% | 1.3% | 4.0% | 66.6% | 25.2% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.2% | 5.6% | 82.2% | 10.0% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 15.0% | 0.0% | 0.0% | 7.5% | 2.5% | 70.0% | 5.0% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.9% | 6.3% | 11.4% | 67.1% | 6.3% |
| 2001 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.5% | 0.0% | 0.0% | 0.0% | 3.0% | 0.0% | 74.6% | 17.9% |
| 2002 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 83.3% | 16.7% |
| 2004 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.1% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 11.3% | 85.8% |
| (82-84) | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 0.0% | 0.0% | 5.3% | 1.8% | 0.0% | 0.0% | 0.4% | 19.3% | 62.4% | 9.5% |
| (90-98) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.3% | 0.9% | 0.3% | 0.6% | 1.7% | 13.6% | 65.4% | 15.0% |
| (99-04) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 3.0% | 0.0% | 1.8% | 3.5% | 2.8% | 61.3% | 26.3% |

Appendix E.38. Percent distribution of South Puget Sound Fall Yearling Chinook for total fishing mortalities among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1982 | 0.0% | 0.0% | 0.0% | 0.0% | 2.2% | 0.0% | 0.0% | 3.8% | 2.7% | 0.0% | 0.0% | 0.8% | 12.7% | 71.4% | 6.5% |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 1.8% | 0.0% | 0.0% | 5.5% | 0.4% | 0.0% | 0.0% | 0.0% | 8.8% | 78.8% | 4.7% |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.0% | 1.8% | 0.0% | 0.0% | 0.0% | 31.7% | 46.5% | 12.9% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.1% | 0.0% | 0.8% | 0.1% | 0.4% | 0.0% | 1.6% | 30.5% | 56.9% | 9.5% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.4% | 0.6% | 0.0% | 0.0% | 3.5% | 11.4% | 62.5% | 16.5% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.0% | 0.9% | 0.0% | 1.2% | 4.8% | 27.0% | 51.5% | 9.6% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 1.0% | 0.0% | 0.0% | 1.2% | 6.7% | 75.0% | 15.0% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.0% | 0.6% | 2.3% | 0.6% | 0.0% | 14.5% | 67.0% | 14.0% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.9% | 2.0% | 0.4% | 1.6% | 0.3% | 8.2% | 74.7% | 6.9% |
| 1996 | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 1.9% | 0.0% | 1.2% | 0.6% | 2.8% | 90.0% | 2.8% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 1.0% | 0.0% | 0.3% | 1.2% | 3.4% | 72.0% | 20.6% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.7% | 4.3% | 86.1% | 7.8% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.6% | 0.0% | 0.0% | 3.8% | 1.0% | 84.8% | 1.9% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.2% | 6.2% | 9.3% | 71.1% | 5.2% |
| 2001 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.3% | 0.0% | 0.0% | 0.0% | 2.2% | 0.0% | 81.3% | 13.2% |
| 2002 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 89.5% | 10.5% |
| 2004 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 15.0% | 82.3% |
| (82-84) | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 0.0% | 5.4% | 1.7% | 0.0% | 0.0% | 0.3% | 17.7% | 65.6% | 8.0% |
| (90-98) | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.3% | 0.9% | 0.3% | 0.6% | 1.6% | 12.1% | 70.6% | 11.4% |
| (99-04) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 1.7% | 0.0% | 1.6% | 2.6% | 2.0% | 68.3% | 22.6% |

¹ No data are shown for 2003 because of lack of coded-wire tagging of broods from 1998 and 2000, for both landed catch and total mortality.

Appendix E.39. Percent distribution of Squaxin Pens Fall Yearling Chinook reported catch among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement ² |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|-------------------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 3.4% | 0.7% | 1.2% | 0.6% | 4.1% | 33.5% | 56.3% | NA ² |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.4% | 1.6% | 0.6% | 0.0% | 9.1% | 34.0% | 50.3% | NA ² |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.5% | 2.4% | 3.6% | 1.3% | 0.8% | 7.4% | 23.5% | 60.1% | NA ² |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.0% | 0.0% | 11.2% | 6.2% | 1.6% | 2.7% | 15.6% | 3.9% | 57.7% | NA ² |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 31.5% | 7.5% | 4.5% | 6.0% | 8.3% | 28.6% | 13.5% | NA ² |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 60.9% | 39.1% | NA ² |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.9% | 0.0% | 0.0% | 1.1% | 4.8% | 92.1% | NA ² |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.4% | 0.0% | 0.0% | 0.0% | 2.8% | 8.0% | 85.7% | NA ² |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.0% | 3.0% | 94.0% | NA ² |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 83.3% | NA ² |
| (90-98) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.1% | 6.3% | 2.4% | 1.0% | 1.1% | 5.7% | 22.3% | 61.0% | NA ² |
| (1999) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 83.3% | NA ² |

¹ No data are shown for 2000-2003 because of lack of coded-wire tagging of broods from 1998-2000.

² Values represent estimates of catch distribution only because escapement data is of insufficient quality.

Appendix E.40. Percent distribution of Squaxin Pens Fall Yearling Chinook total fishing mortalities among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement ² |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|-------------------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 3.3% | 0.8% | 1.0% | 0.6% | 4.2% | 32.2% | 57.8% | NA ² |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.4% | 1.7% | 0.5% | 0.0% | 9.2% | 31.8% | 52.4% | NA ² |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.4% | 2.1% | 3.1% | 0.9% | 0.6% | 6.2% | 22.9% | 63.5% | NA ² |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.0% | 12.1% | 6.7% | 1.5% | 2.3% | 14.7% | 4.1% | 57.7% | NA ² |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 29.0% | 7.2% | 5.2% | 6.0% | 7.8% | 25.7% | 19.1% | NA ² |
| 1995 ³ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.4% | 0.0% | 0.0% | 0.0% | 23.8% | 75.4% | NA ² |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.8% | 0.0% | 0.0% | 0.9% | 5.3% | 91.9% | NA ² |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.0% | 0.0% | 0.0% | 0.0% | 2.1% | 6.4% | 88.4% | NA ² |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.4% | 2.4% | 95.3% | NA ² |
| 1999 ³ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 0.5% | 0.0% | 0.5% | 1.0% | 0.5% | 95.4% | NA ² |
| (90-98) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 6.0% | 2.4% | 1.0% | 1.1% | 5.3% | 17.2% | 66.8% | NA ² |
| (1999) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 0.5% | 0.0% | 0.5% | 1.0% | 0.5% | 95.4% | NA ² |

¹ No data are shown for 2000-2003 because of lack of coded-wire tagging of broods from 1998-2000.

² Values represent estimates of catch distribution only because escapement data is of insufficient quality.

³ Relatively high age-2 survival, combined with relatively few total catch recoveries of CWTs, result in large estimates of sublegal CNR mortality in 1995 and 1999.

Appendix E.41. Percent distribution of White River Spring Yearling Chinook reported catch among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|-----------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1982 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.4% | 0.0% | 0.0% | 0.0% | 74.1% | 23.5% | NA ¹ |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.6% | 0.0% | 4.3% | 0.0% | 0.0% | 0.0% | 1.6% | 11.3% | 59.7% | 21.5% |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 5.8% | 0.0% | 0.0% | 4.5% | 5.2% | 0.0% | 0.0% | 2.6% | 9.0% | 25.2% | 47.7% |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.9% | 2.2% | 0.0% | 30.8% | 50.6% | 13.5% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.6% | 2.4% | 2.0% | 0.0% | 0.4% | 15.3% | 52.3% | 26.8% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.5% | 0.4% | 0.0% | 3.3% | 11.3% | 42.3% | 41.2% |
| 1988 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 2.5% | 0.2% | 0.8% | 1.3% | 13.0% | 48.4% | 33.6% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 1.2% | 1.0% | 0.0% | 6.0% | 13.6% | 41.1% | 35.8% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.9% | 0.4% | 0.6% | 0.0% | 5.2% | 15.4% | 44.6% | 31.8% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 1.3% | 0.0% | 1.3% | 4.1% | 10.8% | 38.1% | 43.6% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 2.4% | 1.9% | 2.3% | 0.8% | 2.4% | 7.8% | 45.5% | 36.2% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 0.0% | 2.9% | 3.6% | 30.6% | 62.2% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.8% | 0.9% | 0.0% | 0.0% | 1.4% | 45.2% | 50.7% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.8% | 29.4% | 69.3% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.0% | 0.0% | 0.0% | 0.3% | 42.9% | 55.9% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.7% | 40.4% | 55.8% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.6% | 1.6% | 27.0% | 69.8% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.4% | 2.4% | 0.0% | 0.0% | 0.0% | 0.0% | 30.5% | 64.6% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.7% | 0.0% | 0.0% | 0.0% | 0.0% | 2.4% | 37.6% | 55.3% |
| (83-84) | 0.0% | 0.0% | 0.0% | 0.0% | 2.9% | 0.8% | 0.0% | 4.4% | 2.6% | 0.0% | 0.0% | 2.1% | 10.2% | 42.5% | 34.6% |
| (85-98) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 1.1% | 0.7% | 0.4% | 1.9% | 9.2% | 41.3% | 44.7% |
| (99-00) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.6% | 1.2% | 0.0% | 0.0% | 0.0% | 1.2% | 34.1% | 60.0% |

¹ No data are shown for 2001 to 2003 because of lack of coded-wire tagging of broods from 1998-2000.

² Values represent estimates of catch distribution only for this year because escapement data is of insufficient quality.

Appendix E.42. Percent distribution of White River Spring Yearling Chinook total fishing mortalities among fisheries and escapement.

| Catch ¹ Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|----------------------------|-----------------|---------------|-----------------|----------------|------------------|--------------|----------------|---------------|----------------|-----------------|-----------------|---------------|-------------|---------------|-----------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1982 | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.0% | 0.0% | 1.9% | 1.9% | 0.0% | 0.0% | 0.9% | 60.4% | 33.9% | NA ¹ |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 0.0% | 4.3% | 0.0% | 0.0% | 0.0% | 1.4% | 10.4% | 63.5% | 19.0% |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 4.8% | 0.0% | 0.0% | 3.9% | 4.4% | 0.0% | 0.0% | 1.8% | 7.0% | 45.6% | 32.5% |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.5% | 1.8% | 0.0% | 25.7% | 60.3% | 9.6% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.6% | 2.3% | 2.0% | 0.0% | 0.4% | 14.1% | 56.5% | 23.6% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.0% | 0.4% | 0.0% | 2.5% | 8.2% | 61.9% | 25.9% |
| 1988 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 2.9% | 0.2% | 0.8% | 1.4% | 12.6% | 52.3% | 29.6% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 1.3% | 1.0% | 0.0% | 6.3% | 12.3% | 46.5% | 31.4% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 0.4% | 0.6% | 0.0% | 5.5% | 13.7% | 50.6% | 27.2% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 1.3% | 0.0% | 1.3% | 4.1% | 9.8% | 46.0% | 36.7% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 2.7% | 2.1% | 2.1% | 0.7% | 2.7% | 7.5% | 49.0% | 32.9% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 2.8% | 3.1% | 39.6% | 53.9% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 0.8% | 0.0% | 0.0% | 1.6% | 52.4% | 43.3% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 0.9% | 41.3% | 57.2% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 0.0% | 0.0% | 0.0% | 0.3% | 48.5% | 50.1% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.5% | 49.5% | 47.0% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 1.4% | 33.3% | 63.8% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.9% | 1.9% | 0.0% | 0.0% | 0.0% | 0.0% | 45.2% | 51.0% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.2% | 0.0% | 0.0% | 0.0% | 0.0% | 2.1% | 44.2% | 49.5% |
| (83-84) | 0.0% | 0.0% | 0.0% | 0.0% | 2.4% | 0.7% | 0.0% | 4.1% | 2.2% | 0.0% | 0.0% | 1.6% | 8.7% | 54.6% | 25.8% |
| (85-98) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 1.1% | 0.7% | 0.3% | 1.9% | 8.2% | 49.1% | 38.0% |
| (99-00) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.1% | 1.0% | 0.0% | 0.0% | 0.0% | 1.1% | 44.7% | 50.2% |

¹ No data are shown for 2001 to 2003 because of lack of coded-wire tagging of broods from 1998-2000.

² Values represent estimates of total fishing mortality distribution only for this year because escapement data is of insufficient quality.

Appendix E.43. Percent distribution of Hoko Fall Fingerling Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1989 | 4.8% | 0.8% | 0.0% | 7.6% | 0.4% | 6.0% | 0.0% | 10.8% | 1.6% | 15.3% | 0.0% | 0.8% | 0.8% | 21.7% | 29.3% |
| 1990 | 15.8% | 1.9% | 0.5% | 8.0% | 0.7% | 2.4% | 0.0% | 17.0% | 0.8% | 1.9% | 0.0% | 0.5% | 1.0% | 14.4% | 35.1% |
| 1991 | 15.2% | 0.0% | 0.0% | 5.0% | 1.1% | 0.3% | 0.6% | 6.9% | 0.4% | 0.6% | 0.5% | 0.2% | 1.0% | 8.2% | 59.8% |
| 1992 | 7.7% | 1.7% | 1.2% | 4.4% | 1.2% | 1.4% | 0.7% | 9.8% | 0.5% | 0.0% | 2.1% | 0.0% | 0.2% | 2.4% | 66.6% |
| 1993 | 6.6% | 0.0% | 2.0% | 6.6% | 0.0% | 3.3% | 0.0% | 14.9% | 0.3% | 2.0% | 0.0% | 0.0% | 0.3% | 4.6% | 59.4% |
| 1994 | 13.6% | 2.1% | 2.4% | 14.8% | 0.6% | 1.5% | 0.0% | 11.4% | 2.1% | 1.5% | 2.1% | 0.0% | 0.0% | 0.0% | 47.9% |
| 1995 | 12.6% | 0.0% | 4.1% | 6.2% | 0.0% | 0.3% | 0.4% | 2.9% | 0.8% | 0.1% | 0.0% | 0.0% | 0.0% | 0.7% | 71.9% |
| 1996 | 10.6% | 0.0% | 3.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 85.2% |
| 1997 | 13.9% | 0.0% | 0.0% | 1.7% | 0.2% | 0.0% | 0.6% | 0.9% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 0.5% | 81.7% |
| 1998 | 9.0% | 0.0% | 0.4% | 5.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 84.1% |
| 1999 | 6.6% | 0.0% | 0.7% | 4.3% | 0.0% | 0.0% | 0.6% | 0.0% | 0.3% | 0.0% | 1.4% | 0.0% | 0.1% | 0.0% | 86.0% |
| 2000 | 4.4% | 0.2% | 1.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 1.2% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 91.7% |
| 2001 | 6.0% | 0.0% | 1.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 90.1% |
| 2002 | 17.2% | 0.0% | 0.9% | 3.7% | 0.3% | 0.0% | 4.7% | 1.5% | 1.9% | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 68.7% |
| 2003 | 13.6% | 0.1% | 2.7% | 3.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.1% | 0.0% | 1.8% | 0.0% | 0.0% | 0.0% | 76.7% |
| 2004 | 11.4% | 0.0% | 1.1% | 8.5% | 0.0% | 0.0% | 0.8% | 0.6% | 1.9% | 0.0% | 1.0% | 0.0% | 0.0% | 0.8% | 74.0% |
| (89-98) | 11.0% | 0.7% | 1.4% | 6.0% | 0.4% | 1.5% | 0.2% | 7.5% | 0.7% | 2.1% | 0.6% | 0.2% | 0.4% | 5.3% | 62.1% |
| (99-04) | 9.9% | 0.1% | 1.5% | 3.2% | 0.0% | 0.0% | 1.0% | 0.4% | 1.5% | 0.0% | 0.7% | 0.1% | 0.0% | 0.4% | 81.2% |

Appendix E.44. Percent distribution of Hoko Fall Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1989 | 11.7% | 3.1% | 0.3% | 8.5% | 1.1% | 4.8% | 0.0% | 13.7% | 1.7% | 11.4% | 0.0% | 0.6% | 0.6% | 21.7% | 20.8% |
| 1990 | 18.5% | 4.8% | 0.6% | 8.4% | 0.9% | 2.0% | 0.0% | 16.9% | 0.7% | 1.6% | 0.0% | 0.6% | 0.9% | 14.1% | 30.1% |
| 1991 | 18.8% | 0.0% | 0.1% | 5.2% | 1.1% | 0.3% | 0.5% | 7.0% | 0.4% | 0.6% | 0.4% | 0.2% | 1.0% | 8.8% | 55.5% |
| 1992 | 8.6% | 4.9% | 1.6% | 5.5% | 1.1% | 1.4% | 0.6% | 10.3% | 0.6% | 0.0% | 2.1% | 0.0% | 0.2% | 2.7% | 60.4% |
| 1993 | 12.3% | 1.1% | 2.3% | 7.7% | 0.0% | 2.9% | 0.0% | 14.9% | 0.6% | 1.7% | 0.0% | 0.0% | 0.3% | 4.9% | 51.4% |
| 1994 | 20.8% | 4.8% | 2.8% | 13.5% | 0.5% | 1.3% | 0.0% | 10.7% | 2.0% | 1.5% | 1.8% | 0.0% | 0.0% | 0.0% | 40.4% |
| 1995 | 16.4% | 0.0% | 4.7% | 7.8% | 0.0% | 0.4% | 0.5% | 3.7% | 0.8% | 0.1% | 0.0% | 0.0% | 0.0% | 1.0% | 64.6% |
| 1996 | 14.1% | 0.0% | 4.4% | 0.7% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 79.1% |
| 1997 | 16.5% | 0.0% | 0.0% | 1.8% | 0.2% | 0.0% | 0.7% | 1.1% | 0.0% | 0.1% | 0.5% | 0.0% | 0.0% | 0.4% | 78.6% |
| 1998 | 10.0% | 0.0% | 0.3% | 6.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 82.8% |
| 1999 | 8.0% | 0.0% | 0.7% | 4.7% | 0.0% | 0.0% | 0.7% | 0.0% | 0.3% | 0.0% | 1.5% | 0.0% | 0.1% | 0.0% | 84.1% |
| 2000 | 5.9% | 0.2% | 2.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 1.3% | 0.0% | 0.0% | 0.8% | 0.0% | 0.0% | 88.7% |
| 2001 | 8.9% | 0.0% | 3.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 85.6% |
| 2002 | 19.6% | 0.0% | 1.0% | 4.0% | 0.3% | 0.0% | 5.7% | 1.4% | 2.5% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 64.5% |
| 2003 | 15.0% | 0.2% | 2.9% | 3.3% | 0.0% | 0.0% | 0.0% | 0.0% | 2.5% | 0.0% | 2.2% | 0.0% | 0.0% | 0.0% | 74.0% |
| 2004 | 13.5% | 0.0% | 1.2% | 9.7% | 0.0% | 0.0% | 1.1% | 0.6% | 2.3% | 0.0% | 1.1% | 0.0% | 0.0% | 0.9% | 69.5% |
| (89-98) | 14.8% | 1.9% | 1.7% | 6.6% | 0.5% | 1.3% | 0.2% | 8.0% | 0.7% | 1.7% | 0.5% | 0.1% | 0.3% | 5.4% | 56.4% |
| (99-04) | 11.8% | 0.1% | 1.9% | 3.6% | 0.0% | 0.0% | 1.2% | 0.4% | 1.8% | 0.0% | 0.8% | 0.1% | 0.0% | 0.4% | 77.7% |

Appendix E.45. Percent distribution of Sooes Fall Fingerling Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1989 | 7.0% | 1.3% | 0.0% | 0.0% | 0.0% | 4.4% | 0.0% | 1.9% | 0.0% | 1.9% | 8.2% | 0.0% | 0.0% | 0.0% | 75.3% |
| 1990 | 9.9% | 2.8% | 4.3% | 14.2% | 1.4% | 0.7% | 0.0% | 17.7% | 7.1% | 2.1% | 0.0% | 1.4% | 0.0% | 3.5% | 34.8% |
| 1991 | 11.9% | 0.0% | 0.0% | 9.9% | 0.0% | 1.7% | 0.0% | 5.2% | 0.0% | 2.0% | 0.0% | 0.0% | 0.0% | 4.9% | 64.3% |
| 1992 | 8.5% | 0.0% | 0.0% | 9.5% | 2.0% | 0.0% | 0.0% | 19.3% | 1.0% | 3.4% | 1.7% | 0.3% | 0.0% | 2.4% | 51.9% |
| 1993 | 4.6% | 0.0% | 0.0% | 7.6% | 2.1% | 2.1% | 2.1% | 16.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.8% | 64.1% |
| 1994 | 17.0% | 3.0% | 4.0% | 10.5% | 1.0% | 0.0% | 1.0% | 8.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 55.5% |
| 1995 | 8.5% | 0.0% | 0.0% | 4.6% | 0.0% | 0.7% | 0.0% | 9.8% | 0.0% | 0.0% | 0.0% | 0.0% | 2.6% | 0.0% | 73.9% |
| 1996 | 8.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 90.3% |
| 1997 | 10.3% | 0.0% | 5.2% | 5.5% | 0.7% | 0.3% | 0.0% | 0.0% | 1.4% | 0.0% | 2.8% | 1.0% | 23.4% | 0.0% | 49.3% |
| 1998 | 9.0% | 0.0% | 1.5% | 17.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 72.0% |
| 1999 | 12.3% | 0.0% | 12.3% | 4.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.0% | 70.5% |
| 2000 | 0.0% | 0.0% | 2.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 10.7% | 0.0% | 0.0% | 0.0% | 86.9% |
| 2001 | 6.1% | 0.0% | 2.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 0.0% | 1.7% | 0.0% | 0.0% | 0.0% | 88.8% |
| 2002 | 10.9% | 0.2% | 1.3% | 1.7% | 0.0% | 0.0% | 1.9% | 0.6% | 1.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 81.6% |
| 2003 | 11.7% | 0.1% | 0.0% | 4.5% | 0.0% | 0.0% | 6.1% | 0.0% | 0.9% | 0.0% | 0.0% | 0.0% | 24.3% | 1.3% | 51.1% |
| 2004 | 17.4% | 0.5% | 2.0% | 14.5% | 0.0% | 0.0% | 0.0% | 0.8% | 0.0% | 0.0% | 1.9% | 0.0% | 0.9% | 0.9% | 61.0% |
| (89-98) | 9.5% | 0.7% | 1.5% | 7.9% | 0.7% | 1.0% | 0.3% | 7.8% | 0.9% | 0.9% | 1.3% | 0.3% | 2.6% | 1.2% | 63.1% |
| (99-04) | 9.7% | 0.1% | 3.3% | 4.1% | 0.0% | 0.0% | 1.3% | 0.2% | 0.5% | 0.0% | 2.4% | 0.0% | 4.4% | 0.5% | 73.3% |

Appendix E.46. Percent distribution of Sooes Fall Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1989 | 11.0% | 3.7% | 0.5% | 3.1% | 0.0% | 3.7% | 0.0% | 4.7% | 0.0% | 2.1% | 7.3% | 0.0% | 0.0% | 1.6% | 62.3% |
| 1990 | 11.6% | 7.0% | 4.1% | 16.3% | 1.7% | 0.6% | 0.0% | 17.4% | 6.4% | 1.7% | 0.0% | 1.7% | 0.0% | 2.9% | 28.5% |
| 1991 | 14.1% | 0.0% | 0.3% | 10.6% | 0.3% | 1.6% | 0.0% | 7.2% | 0.0% | 1.9% | 0.0% | 0.0% | 0.0% | 5.1% | 59.0% |
| 1992 | 11.0% | 0.3% | 0.3% | 10.7% | 2.1% | 0.0% | 0.0% | 20.4% | 1.2% | 3.0% | 1.5% | 0.3% | 0.0% | 2.4% | 46.6% |
| 1993 | 7.5% | 0.4% | 0.0% | 7.9% | 2.0% | 2.0% | 2.0% | 16.9% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 1.2% | 59.8% |
| 1994 | 21.0% | 7.4% | 3.5% | 9.6% | 0.9% | 0.0% | 0.9% | 7.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.0% | 48.5% |
| 1995 | 14.9% | 0.0% | 0.0% | 6.1% | 0.0% | 1.1% | 0.0% | 12.7% | 0.0% | 0.6% | 0.0% | 0.0% | 2.2% | 0.0% | 62.4% |
| 1996 | 15.5% | 0.0% | 0.0% | 0.9% | 0.0% | 0.4% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 82.3% |
| 1997 | 12.0% | 0.0% | 5.8% | 5.8% | 0.6% | 0.3% | 0.0% | 0.0% | 1.3% | 0.3% | 2.6% | 1.0% | 23.7% | 0.0% | 46.4% |
| 1998 | 10.3% | 0.0% | 1.8% | 19.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 68.7% |
| 1999 | 13.5% | 0.0% | 13.5% | 4.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 67.4% |
| 2000 | 0.0% | 0.0% | 5.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 13.3% | 0.0% | 0.0% | 0.0% | 81.1% |
| 2001 | 9.6% | 0.0% | 2.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.6% | 0.0% | 2.2% | 0.0% | 0.0% | 0.0% | 83.7% |
| 2002 | 13.4% | 0.4% | 1.6% | 2.0% | 0.0% | 0.0% | 2.5% | 0.5% | 1.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 77.5% |
| 2003 | 13.5% | 0.4% | 0.0% | 5.1% | 0.0% | 0.0% | 7.8% | 0.0% | 1.1% | 0.0% | 0.0% | 0.0% | 23.5% | 1.4% | 47.3% |
| 2004 | 19.5% | 1.4% | 2.2% | 15.9% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 0.0% | 2.1% | 0.0% | 0.8% | 0.9% | 56.5% |
| (89-98) | 12.9% | 1.9% | 1.6% | 9.0% | 0.8% | 1.0% | 0.3% | 8.7% | 0.9% | 1.0% | 1.1% | 0.3% | 2.7% | 1.4% | 56.5% |
| (99-04) | 11.6% | 0.3% | 4.3% | 4.6% | 0.0% | 0.0% | 1.7% | 0.2% | 0.7% | 0.0% | 2.9% | 0.0% | 4.3% | 0.5% | 68.9% |

Appendix E.47. Percent distribution of Queets Fall Fingerling Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1981 | 9.5% | 0.0% | 0.0% | 13.7% | 2.1% | 2.1% | 0.0% | 11.6% | 0.0% | 1.1% | 0.0% | 1.1% | 31.6% | 3.2% | 24.2% |
| 1982 | 11.8% | 2.4% | 0.0% | 22.9% | 0.0% | 0.8% | 1.2% | 12.2% | 0.0% | 0.0% | 0.0% | 0.0% | 25.7% | 0.0% | 22.9% |
| 1983 | 33.3% | 0.0% | 0.0% | 6.8% | 0.0% | 0.8% | 0.0% | 7.6% | 0.0% | 2.3% | 0.0% | 0.8% | 25.8% | 0.0% | 22.7% |
| 1984 | 16.1% | 0.7% | 0.0% | 19.6% | 0.0% | 0.0% | 2.1% | 7.7% | 0.0% | 0.0% | 0.0% | 2.1% | 28.7% | 0.0% | 23.1% |
| 1985 | 15.6% | 0.0% | 0.0% | 31.6% | 0.0% | 0.0% | 0.0% | 2.0% | 0.0% | 1.6% | 0.0% | 0.0% | 14.4% | 1.2% | 33.6% |
| 1986 | 17.3% | 0.0% | 1.1% | 11.6% | 1.8% | 0.0% | 0.0% | 7.0% | 0.0% | 1.1% | 0.0% | 0.0% | 9.9% | 0.0% | 50.4% |
| 1987 | 22.3% | 0.2% | 0.0% | 11.7% | 0.9% | 0.6% | 0.9% | 0.7% | 0.0% | 0.0% | 0.0% | 0.6% | 22.7% | 0.6% | 38.7% |
| 1988 | 14.6% | 0.8% | 1.6% | 7.8% | 2.5% | 0.4% | 0.0% | 4.0% | 0.0% | 0.0% | 1.1% | 0.0% | 16.6% | 3.3% | 47.3% |
| 1989 | 11.1% | 0.0% | 0.0% | 9.1% | 0.5% | 0.2% | 1.1% | 7.6% | 0.0% | 0.0% | 0.0% | 0.0% | 27.8% | 1.6% | 41.1% |
| 1990 | 12.6% | 0.0% | 0.0% | 5.5% | 0.3% | 0.3% | 1.8% | 6.6% | 0.0% | 0.0% | 0.0% | 0.0% | 13.9% | 0.0% | 58.9% |
| 1991 | 20.5% | 0.2% | 1.1% | 9.7% | 0.0% | 0.0% | 1.3% | 4.8% | 0.0% | 0.0% | 0.0% | 0.0% | 15.7% | 0.5% | 46.3% |
| 1992 | 8.3% | 0.8% | 2.2% | 7.7% | 0.0% | 0.2% | 1.9% | 17.5% | 0.0% | 0.0% | 0.0% | 0.0% | 19.2% | 0.8% | 41.4% |
| 1993 | 15.6% | 0.0% | 0.7% | 14.1% | 0.3% | 0.0% | 2.1% | 12.1% | 0.0% | 0.0% | 0.0% | 0.5% | 16.1% | 2.8% | 35.7% |
| 1994 | 16.1% | 0.3% | 0.5% | 21.7% | 0.2% | 0.4% | 1.5% | 4.1% | 0.3% | 0.0% | 1.0% | 0.0% | 21.4% | 0.0% | 32.4% |
| 1995 | 17.5% | 0.0% | 1.6% | 6.1% | 0.0% | 0.1% | 2.0% | 0.7% | 0.3% | 0.0% | 0.4% | 0.7% | 33.8% | 0.0% | 36.7% |
| 1996 | 10.4% | 0.0% | 1.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 17.5% | 0.6% | 70.2% |
| 1997 | 34.4% | 0.3% | 0.0% | 6.0% | 0.8% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 20.9% | 0.0% | 37.4% |
| 1998 | 23.7% | 0.0% | 3.0% | 19.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 12.1% | 5.2% | 37.0% |
| 1999 | 9.3% | 0.0% | 1.4% | 1.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.8% | 0.3% | 78.4% |
| 2000 | 23.7% | 0.0% | 10.0% | 10.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.0% | 0.0% | 51.6% |
| 2001 | 23.7% | 0.0% | 5.9% | 3.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 42.4% | 0.7% | 22.6% |
| 2002 | 26.2% | 0.0% | 3.4% | 1.8% | 0.0% | 0.0% | 2.6% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 25.6% | 0.3% | 39.9% |
| 2003 | 19.7% | 0.1% | 3.4% | 9.9% | 0.0% | 0.0% | 9.6% | 0.0% | 0.0% | 0.0% | 1.2% | 0.0% | 20.1% | 0.6% | 35.4% |
| 2004 | 22.9% | 0.6% | 4.6% | 9.8% | 0.0% | 0.0% | 12.7% | 1.6% | 0.0% | 0.0% | 0.0% | 0.1% | 17.8% | 0.2% | 29.7% |
| (81-84) | 17.7% | 0.8% | 0.0% | 15.7% | 0.5% | 0.9% | 0.8% | 9.8% | 0.0% | 0.8% | 0.0% | 1.0% | 27.9% | 0.8% | 23.2% |
| (85-98) | 17.1% | 0.2% | 0.9% | 11.6% | 0.5% | 0.2% | 0.9% | 4.8% | 0.0% | 0.2% | 0.2% | 0.1% | 18.7% | 1.2% | 43.4% |
| (99-04) | 20.9% | 0.1% | 4.8% | 6.3% | 0.0% | 0.0% | 4.1% | 0.3% | 0.0% | 0.0% | 0.2% | 0.2% | 19.8% | 0.3% | 42.9% |

Appendix E.48. Percent distribution of Queets Fall Fingerling Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1981 | 12.9% | 0.0% | 0.0% | 18.1% | 1.7% | 1.7% | 0.0% | 12.9% | 0.0% | 0.9% | 0.0% | 1.7% | 26.7% | 3.4% | 19.8% |
| 1982 | 14.2% | 2.2% | 0.0% | 24.0% | 0.0% | 0.7% | 1.1% | 12.0% | 0.0% | 0.0% | 0.0% | 0.0% | 24.7% | 0.0% | 21.0% |
| 1983 | 50.5% | 0.0% | 0.0% | 5.5% | 0.0% | 0.5% | 0.0% | 5.5% | 0.0% | 1.6% | 0.0% | 0.5% | 19.2% | 0.0% | 16.5% |
| 1984 | 20.9% | 0.6% | 0.0% | 20.2% | 0.0% | 0.0% | 2.5% | 7.4% | 0.0% | 0.0% | 0.0% | 2.5% | 25.8% | 0.0% | 20.2% |
| 1985 | 20.2% | 0.0% | 0.0% | 33.6% | 0.0% | 0.0% | 0.0% | 2.1% | 0.0% | 1.4% | 0.0% | 0.0% | 12.3% | 1.7% | 28.8% |
| 1986 | 26.8% | 0.0% | 1.2% | 11.0% | 1.5% | 0.0% | 0.0% | 6.8% | 0.0% | 0.9% | 0.0% | 0.0% | 9.2% | 0.0% | 42.6% |
| 1987 | 28.7% | 0.5% | 0.0% | 11.7% | 0.8% | 0.5% | 1.0% | 1.3% | 0.0% | 0.0% | 0.0% | 0.5% | 20.2% | 0.7% | 34.2% |
| 1988 | 17.5% | 2.4% | 1.6% | 9.4% | 2.4% | 0.4% | 0.1% | 5.5% | 0.0% | 0.0% | 1.0% | 0.0% | 14.8% | 3.4% | 41.5% |
| 1989 | 17.0% | 0.2% | 0.2% | 10.6% | 0.6% | 0.3% | 1.1% | 8.9% | 0.0% | 0.0% | 0.0% | 0.0% | 24.3% | 1.7% | 35.3% |
| 1990 | 15.5% | 0.1% | 0.1% | 6.4% | 0.3% | 0.3% | 1.9% | 7.1% | 0.0% | 0.0% | 0.0% | 0.0% | 13.3% | 0.0% | 54.9% |
| 1991 | 24.5% | 0.3% | 1.2% | 10.1% | 0.0% | 0.0% | 1.4% | 5.0% | 0.0% | 0.0% | 0.0% | 0.0% | 14.6% | 0.5% | 42.5% |
| 1992 | 15.4% | 2.2% | 2.4% | 8.6% | 0.0% | 0.1% | 1.8% | 17.9% | 0.0% | 0.0% | 0.0% | 0.0% | 16.2% | 0.8% | 34.4% |
| 1993 | 20.0% | 0.0% | 0.7% | 15.3% | 0.3% | 0.0% | 2.0% | 13.0% | 0.0% | 0.0% | 0.0% | 0.4% | 14.3% | 2.9% | 31.1% |
| 1994 | 24.8% | 0.6% | 0.4% | 20.9% | 0.2% | 0.3% | 1.5% | 4.0% | 0.2% | 0.0% | 1.0% | 0.0% | 18.4% | 0.0% | 27.6% |
| 1995 | 22.4% | 0.0% | 1.8% | 7.5% | 0.0% | 0.2% | 2.5% | 0.8% | 0.2% | 0.0% | 0.4% | 0.7% | 30.5% | 0.0% | 32.7% |
| 1996 | 18.9% | 0.0% | 1.5% | 1.1% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 15.8% | 0.5% | 61.8% |
| 1997 | 38.4% | 0.5% | 0.0% | 6.1% | 0.7% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 19.5% | 0.0% | 34.6% |
| 1998 | 25.6% | 0.0% | 3.1% | 19.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 11.5% | 5.3% | 34.8% |
| 1999 | 13.1% | 0.0% | 1.9% | 2.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.5% | 0.3% | 74.1% |
| 2000 | 27.7% | 0.0% | 12.7% | 11.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.6% | 0.0% | 44.8% |
| 2001 | 29.7% | 0.0% | 6.7% | 4.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 38.0% | 0.6% | 19.8% |
| 2002 | 29.8% | 0.0% | 3.6% | 1.9% | 0.0% | 0.0% | 3.2% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 24.1% | 0.3% | 36.9% |
| 2003 | 21.1% | 0.1% | 3.7% | 10.4% | 0.0% | 0.0% | 12.1% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 18.5% | 0.7% | 32.1% |
| 2004 | 23.8% | 1.7% | 4.4% | 9.4% | 0.0% | 0.0% | 15.8% | 1.5% | 0.0% | 0.0% | 0.0% | 0.1% | 16.2% | 0.2% | 27.0% |
| (81-84) | 24.6% | 0.7% | 0.0% | 17.0% | 0.4% | 0.8% | 0.9% | 9.4% | 0.0% | 0.6% | 0.0% | 1.2% | 24.1% | 0.9% | 19.4% |
| (85-98) | 22.6% | 0.5% | 1.0% | 12.3% | 0.5% | 0.2% | 1.0% | 5.2% | 0.0% | 0.2% | 0.2% | 0.1% | 16.8% | 1.2% | 38.3% |
| (99-04) | 24.2% | 0.3% | 5.5% | 6.5% | 0.0% | 0.0% | 5.2% | 0.2% | 0.0% | 0.0% | 0.2% | 0.2% | 18.2% | 0.3% | 39.1% |

Appendix E.49. Percent distribution of Willamette Spring Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1980 | 6.5% | 0.9% | 0.3% | 11.0% | 0.3% | 0.8% | 0.1% | 4.7% | 0.0% | 0.1% | 0.0% | 0.9% | 0.6% | 15.8% | 57.9% |
| 1981 | 8.7% | 1.1% | 0.2% | 12.0% | 0.8% | 0.2% | 0.0% | 2.7% | 0.0% | 0.0% | 0.0% | 0.7% | 3.1% | 18.4% | 52.2% |
| 1982 | 4.1% | 1.1% | 0.1% | 6.6% | 0.1% | 0.3% | 0.1% | 4.1% | 0.0% | 0.0% | 0.0% | 1.1% | 7.3% | 24.9% | 50.1% |
| 1983 | 12.8% | 0.1% | 0.0% | 12.0% | 0.3% | 0.0% | 0.0% | 1.9% | 0.8% | 0.0% | 0.0% | 1.9% | 6.5% | 21.2% | 42.6% |
| 1984 | 4.0% | 0.3% | 0.3% | 2.1% | 0.1% | 0.1% | 0.1% | 1.9% | 0.1% | 0.0% | 0.0% | 1.0% | 6.2% | 23.9% | 59.8% |
| 1985 | 5.1% | 0.1% | 0.0% | 0.5% | 0.2% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.3% | 18.3% | 20.5% | 54.6% |
| 1986 | 3.1% | 0.4% | 0.0% | 6.6% | 0.6% | 2.5% | 0.0% | 5.5% | 0.0% | 0.0% | 0.6% | 0.0% | 9.2% | 17.1% | 54.4% |
| 1987 | 9.8% | 0.0% | 0.6% | 13.3% | 0.8% | 1.1% | 0.0% | 0.9% | 0.0% | 0.0% | 1.3% | 2.4% | 6.3% | 27.0% | 36.5% |
| 1988 | 8.6% | 0.2% | 0.4% | 6.2% | 0.6% | 0.1% | 0.0% | 3.1% | 0.0% | 0.0% | 0.0% | 2.2% | 6.9% | 28.8% | 42.9% |
| 1989 | 4.4% | 0.0% | 0.2% | 1.8% | 0.0% | 0.1% | 0.0% | 1.4% | 0.5% | 0.2% | 0.5% | 1.5% | 12.6% | 20.3% | 56.6% |
| 1990 | 6.3% | 0.3% | 0.2% | 1.4% | 0.2% | 0.5% | 0.2% | 2.1% | 0.0% | 0.1% | 0.7% | 1.3% | 17.0% | 27.7% | 42.0% |
| 1991 | 3.1% | 1.2% | 0.6% | 1.7% | 0.0% | 0.2% | 0.0% | 0.4% | 0.2% | 0.0% | 0.2% | 0.7% | 6.0% | 42.8% | 43.0% |
| 1992 | 3.5% | 1.3% | 0.2% | 1.7% | 0.0% | 0.2% | 0.2% | 2.7% | 0.0% | 0.1% | 0.2% | 2.4% | 5.8% | 31.3% | 50.4% |
| 1993 | 8.1% | 0.0% | 0.0% | 1.3% | 0.0% | 0.0% | 0.1% | 1.4% | 0.0% | 0.0% | 0.2% | 1.5% | 0.8% | 43.1% | 43.5% |
| 1994 | 4.1% | 0.3% | 0.9% | 0.7% | 0.2% | 0.2% | 0.1% | 0.6% | 0.0% | 0.0% | 0.0% | 0.2% | 5.1% | 38.9% | 48.7% |
| 1995 | 2.8% | 0.1% | 0.3% | 1.0% | 0.0% | 0.3% | 0.0% | 0.3% | 0.0% | 0.0% | 0.1% | 0.0% | 0.3% | 43.8% | 50.9% |
| 1996 | 2.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 1.2% | 7.9% | 88.6% |
| 1997 | 3.6% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.8% | 15.8% | 79.0% |
| 1998 | 4.2% | 0.1% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.1% | 0.4% | 16.4% | 78.5% |
| 1999 | 4.3% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 0.8% | 14.7% | 79.3% |
| 2000 | 7.8% | 0.1% | 0.4% | 0.1% | 0.0% | 0.0% | 0.3% | 0.3% | 0.0% | 0.0% | 0.3% | 0.3% | 2.3% | 29.8% | 58.2% |
| 2001 | 1.4% | 0.0% | 0.1% | 0.1% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.1% | 0.3% | 3.5% | 23.2% | 70.9% |
| 2002 | 1.8% | 0.1% | 0.1% | 0.6% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 0.0% | 0.0% | 0.8% | 15.9% | 20.4% | 59.6% |
| 2003 | 4.8% | 0.0% | 0.1% | 0.4% | 0.0% | 0.0% | 0.3% | 2.3% | 0.0% | 0.0% | 0.6% | 0.3% | 1.5% | 15.7% | 74.1% |
| 2004 | 3.2% | 0.4% | 0.1% | 0.6% | 0.0% | 0.0% | 0.0% | 5.9% | 0.0% | 0.0% | 0.0% | 1.7% | 6.7% | 18.5% | 63.0% |
| (80-84) | 7.2% | 0.7% | 0.2% | 8.7% | 0.3% | 0.3% | 0.1% | 3.1% | 0.2% | 0.0% | 0.0% | 1.1% | 4.7% | 20.8% | 52.5% |
| (85-98) | 4.9% | 0.3% | 0.3% | 2.6% | 0.2% | 0.4% | 0.0% | 1.4% | 0.0% | 0.0% | 0.3% | 0.9% | 6.5% | 27.2% | 55.0% |
| (99-04) | 3.9% | 0.1% | 0.2% | 0.3% | 0.0% | 0.0% | 0.1% | 1.6% | 0.0% | 0.0% | 0.3% | 0.6% | 5.1% | 20.4% | 67.5% |

Appendix E.50. Percent distribution of Willamette Spring Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1980 | 8.7% | 0.9% | 0.3% | 14.2% | 0.4% | 0.8% | 0.1% | 5.8% | 0.0% | 0.1% | 0.0% | 1.1% | 0.7% | 15.2% | 51.5% |
| 1981 | 10.7% | 1.1% | 0.3% | 14.8% | 0.9% | 0.2% | 0.0% | 3.3% | 0.0% | 0.0% | 0.0% | 0.8% | 3.0% | 17.8% | 47.0% |
| 1982 | 5.8% | 1.2% | 0.2% | 8.2% | 0.1% | 0.4% | 0.1% | 5.1% | 0.0% | 0.0% | 0.0% | 1.3% | 7.0% | 24.8% | 45.9% |
| 1983 | 18.9% | 0.1% | 0.0% | 13.2% | 0.3% | 0.0% | 0.0% | 2.0% | 0.8% | 0.0% | 0.0% | 2.1% | 5.9% | 19.9% | 36.6% |
| 1984 | 4.6% | 0.3% | 0.4% | 2.5% | 0.1% | 0.1% | 0.1% | 2.1% | 0.1% | 0.0% | 0.0% | 1.2% | 6.3% | 24.7% | 57.6% |
| 1985 | 7.9% | 0.3% | 0.0% | 0.5% | 0.2% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 0.3% | 17.7% | 20.8% | 51.8% |
| 1986 | 4.9% | 1.2% | 0.0% | 7.5% | 0.7% | 2.6% | 0.0% | 6.2% | 0.0% | 0.0% | 0.7% | 0.0% | 8.8% | 17.1% | 50.3% |
| 1987 | 18.8% | 0.0% | 1.0% | 15.4% | 1.2% | 1.0% | 0.0% | 1.5% | 0.0% | 0.0% | 1.2% | 3.1% | 5.3% | 23.1% | 28.4% |
| 1988 | 11.5% | 0.4% | 0.6% | 7.8% | 0.8% | 0.0% | 0.0% | 3.7% | 0.0% | 0.0% | 0.0% | 2.4% | 6.5% | 30.3% | 36.0% |
| 1989 | 5.7% | 0.0% | 0.2% | 2.2% | 0.0% | 0.1% | 0.0% | 1.6% | 0.6% | 0.1% | 0.6% | 1.7% | 12.2% | 22.1% | 52.8% |
| 1990 | 10.3% | 0.8% | 0.3% | 2.0% | 0.2% | 0.5% | 0.2% | 2.7% | 0.0% | 0.1% | 0.7% | 1.5% | 15.6% | 28.0% | 37.3% |
| 1991 | 4.1% | 2.9% | 0.7% | 2.1% | 0.0% | 0.2% | 0.0% | 0.4% | 0.2% | 0.0% | 0.2% | 0.7% | 5.7% | 44.3% | 38.5% |
| 1992 | 7.7% | 3.2% | 0.2% | 2.0% | 0.0% | 0.1% | 0.2% | 3.1% | 0.0% | 0.1% | 0.2% | 2.8% | 5.3% | 31.5% | 43.5% |
| 1993 | 13.4% | 0.0% | 0.0% | 1.5% | 0.0% | 0.0% | 0.1% | 1.6% | 0.0% | 0.0% | 0.2% | 1.6% | 0.7% | 43.9% | 36.9% |
| 1994 | 5.8% | 0.7% | 1.1% | 0.9% | 0.3% | 0.2% | 0.1% | 0.8% | 0.0% | 0.0% | 0.0% | 0.2% | 4.8% | 40.8% | 44.4% |
| 1995 | 5.3% | 0.1% | 0.4% | 1.4% | 0.0% | 0.4% | 0.0% | 0.5% | 0.0% | 0.0% | 0.1% | 0.0% | 0.3% | 46.0% | 45.5% |
| 1996 | 3.4% | 0.0% | 0.0% | 0.2% | 0.0% | 0.3% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.1% | 1.2% | 8.9% | 85.9% |
| 1997 | 4.5% | 0.0% | 0.0% | 0.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.8% | 17.2% | 76.4% |
| 1998 | 5.7% | 0.4% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 0.0% | 0.0% | 0.2% | 0.4% | 18.5% | 74.4% |
| 1999 | 9.2% | 0.0% | 1.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.0% | 0.8% | 16.0% | 72.2% |
| 2000 | 13.8% | 0.2% | 1.0% | 0.1% | 0.0% | 0.0% | 0.4% | 0.3% | 0.0% | 0.0% | 0.4% | 0.3% | 2.2% | 31.7% | 49.7% |
| 2001 | 1.6% | 0.1% | 0.1% | 0.1% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.1% | 0.3% | 3.7% | 27.0% | 66.6% |
| 2002 | 2.2% | 0.3% | 0.1% | 0.7% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 0.0% | 0.0% | 0.9% | 15.4% | 22.5% | 57.0% |
| 2003 | 6.0% | 0.0% | 0.1% | 0.5% | 0.0% | 0.0% | 0.4% | 2.5% | 0.0% | 0.0% | 0.8% | 0.3% | 1.5% | 17.2% | 70.8% |
| 2004 | 4.1% | 1.3% | 0.1% | 0.7% | 0.0% | 0.0% | 0.0% | 6.0% | 0.0% | 0.0% | 0.0% | 1.9% | 6.6% | 20.5% | 58.7% |
| (80-84) | 9.7% | 0.7% | 0.2% | 10.6% | 0.4% | 0.3% | 0.1% | 3.7% | 0.2% | 0.0% | 0.0% | 1.3% | 4.6% | 20.5% | 47.7% |
| (85-98) | 7.8% | 0.7% | 0.3% | 3.2% | 0.2% | 0.4% | 0.0% | 1.6% | 0.1% | 0.0% | 0.3% | 1.1% | 6.1% | 28.0% | 50.2% |
| (99-04) | 6.2% | 0.3% | 0.4% | 0.3% | 0.0% | 0.0% | 0.1% | 1.7% | 0.0% | 0.0% | 0.3% | 0.6% | 5.0% | 22.5% | 62.5% |

Appendix E.51. Percent distribution of Columbia Summer Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 11.4% | 0.0% | 1.2% | 7.2% | 2.4% | 9.6% | 0.0% | 16.3% | 7.8% | 1.8% | 0.0% | 0.0% | 4.8% | 4.8% | 32.5% |
| 1980 | 33.1% | 0.0% | 0.9% | 8.8% | 4.0% | 1.2% | 0.0% | 16.7% | 0.0% | 0.0% | 0.0% | 1.5% | 0.6% | 0.0% | 33.1% |
| 1987 | 13.6% | 0.0% | 0.0% | 5.6% | 4.8% | 4.0% | 3.2% | 0.0% | 0.0% | 0.0% | 0.0% | 20.0% | 15.2% | 0.0% | 33.6% |
| 1988 | 1.1% | 0.8% | 0.0% | 7.6% | 0.0% | 7.6% | 1.9% | 15.9% | 0.0% | 1.5% | 4.2% | 3.4% | 15.2% | 3.0% | 37.9% |
| 1989 | 4.8% | 0.5% | 0.6% | 5.1% | 0.6% | 0.3% | 0.6% | 14.8% | 1.4% | 2.2% | 2.4% | 14.4% | 8.5% | 2.6% | 41.1% |
| 1990 | 9.7% | 0.0% | 0.0% | 6.6% | 1.1% | 1.3% | 0.0% | 19.5% | 0.6% | 0.4% | 0.0% | 5.7% | 10.8% | 2.5% | 41.8% |
| 1991 | 3.9% | 0.0% | 0.0% | 2.2% | 0.5% | 1.6% | 0.0% | 5.7% | 0.0% | 1.1% | 0.7% | 3.4% | 3.9% | 2.2% | 74.8% |
| 1992 | 14.1% | 0.0% | 0.0% | 3.4% | 2.1% | 1.0% | 0.0% | 14.8% | 0.7% | 0.0% | 0.0% | 6.5% | 1.4% | 1.4% | 54.6% |
| 1993 | 7.1% | 0.0% | 0.0% | 1.4% | 0.0% | 2.4% | 0.0% | 14.3% | 0.0% | 0.0% | 1.9% | 5.2% | 3.3% | 1.4% | 62.9% |
| 1994 | 13.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 13.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 10.8% | 0.0% | 62.2% |
| 1995 | 2.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 5.1% | 0.0% | 0.0% | 0.0% | 2.2% | 1.4% | 0.0% | 88.4% |
| 1996 | 13.3% | 0.6% | 0.0% | 0.0% | 0.0% | 3.0% | 0.0% | 0.0% | 2.2% | 0.0% | 0.0% | 2.8% | 3.9% | 4.1% | 70.2% |
| 1997 | 7.8% | 0.1% | 3.2% | 0.2% | 0.0% | 0.4% | 0.9% | 1.6% | 0.0% | 0.0% | 0.0% | 2.9% | 1.2% | 0.8% | 80.9% |
| 1998 | 8.6% | 0.1% | 0.9% | 0.5% | 0.0% | 0.1% | 0.5% | 0.0% | 0.0% | 0.0% | 0.6% | 1.9% | 5.0% | 1.0% | 80.9% |
| 1999 | 10.1% | 2.5% | 1.8% | 0.4% | 0.0% | 0.6% | 2.7% | 0.6% | 0.0% | 0.0% | 5.0% | 8.5% | 1.2% | 3.4% | 63.2% |
| 2000 | 21.7% | 1.4% | 2.6% | 0.4% | 0.0% | 0.0% | 1.4% | 4.5% | 0.6% | 0.0% | 5.0% | 3.1% | 1.1% | 3.9% | 54.3% |
| 2001 | 14.1% | 2.8% | 1.5% | 0.5% | 0.0% | 0.0% | 1.4% | 12.3% | 0.2% | 0.0% | 4.4% | 17.6% | 0.8% | 6.2% | 38.1% |
| 2002 | 22.4% | 0.0% | 1.4% | 10.5% | 0.0% | 0.0% | 2.1% | 15.5% | 0.1% | 0.0% | 0.8% | 8.7% | 1.1% | 5.9% | 31.4% |
| 2003 | 25.4% | 0.4% | 1.0% | 10.7% | 0.0% | 0.0% | 5.1% | 12.0% | 0.0% | 0.0% | 0.8% | 6.3% | 2.9% | 6.7% | 28.4% |
| 2004 | 13.8% | 0.3% | 1.1% | 4.9% | 0.0% | 0.0% | 1.4% | 11.6% | 0.2% | 0.0% | 1.4% | 10.7% | 8.0% | 15.9% | 30.6% |
| (79-80) | 22.3% | 0.0% | 1.1% | 8.0% | 3.2% | 5.4% | 0.0% | 16.5% | 3.9% | 0.9% | 0.0% | 0.8% | 2.7% | 2.4% | 32.8% |
| (87-98) | 8.4% | 0.2% | 0.4% | 2.7% | 0.8% | 1.8% | 1.7% | 7.6% | 0.4% | 0.4% | 0.8% | 5.7% | 6.7% | 1.6% | 60.8% |
| (99-04) | 17.9% | 1.2% | 1.6% | 4.6% | 0.0% | 0.1% | 2.4% | 9.4% | 0.2% | 0.0% | 2.9% | 9.2% | 2.5% | 7.0% | 41.0% |

Appendix E.52. Percent distribution of Columbia Summer Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 14.4% | 0.0% | 1.0% | 9.0% | 4.0% | 8.5% | 0.0% | 18.9% | 7.0% | 1.5% | 0.0% | 0.5% | 4.0% | 4.5% | 26.9% |
| 1980 | 32.8% | 0.0% | 0.9% | 9.2% | 4.3% | 1.1% | 0.0% | 18.1% | 0.0% | 0.0% | 0.0% | 1.7% | 0.6% | 0.0% | 31.3% |
| 1987 | 16.0% | 0.0% | 0.0% | 8.0% | 3.7% | 4.3% | 2.5% | 7.4% | 0.0% | 0.0% | 0.0% | 19.8% | 11.7% | 0.6% | 25.9% |
| 1988 | 1.9% | 2.2% | 0.0% | 10.0% | 0.0% | 7.5% | 1.9% | 20.9% | 0.0% | 1.2% | 4.0% | 3.4% | 13.1% | 2.8% | 31.2% |
| 1989 | 7.1% | 2.1% | 0.7% | 5.6% | 0.7% | 0.3% | 0.6% | 16.4% | 1.4% | 1.9% | 2.4% | 14.9% | 7.5% | 2.5% | 35.9% |
| 1990 | 10.6% | 0.0% | 0.0% | 7.6% | 1.1% | 1.3% | 0.0% | 20.3% | 0.6% | 0.3% | 0.0% | 5.7% | 10.3% | 2.6% | 39.5% |
| 1991 | 4.1% | 0.0% | 0.0% | 2.3% | 0.5% | 1.7% | 0.0% | 6.3% | 0.0% | 1.1% | 0.7% | 3.6% | 4.0% | 2.3% | 73.5% |
| 1992 | 18.5% | 0.0% | 0.0% | 3.4% | 1.9% | 0.9% | 0.0% | 15.4% | 0.6% | 0.0% | 0.0% | 6.6% | 1.3% | 1.6% | 49.8% |
| 1993 | 7.8% | 0.0% | 0.0% | 1.4% | 0.0% | 2.8% | 0.0% | 15.6% | 0.0% | 0.0% | 1.8% | 5.5% | 3.2% | 1.4% | 60.6% |
| 1994 | 17.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 15.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 10.0% | 0.0% | 57.5% |
| 1995 | 4.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.4% | 0.0% | 1.4% | 0.0% | 2.0% | 2.7% | 0.0% | 82.4% |
| 1996 | 21.3% | 0.7% | 0.0% | 1.8% | 0.0% | 3.0% | 0.0% | 2.5% | 2.5% | 0.2% | 0.0% | 2.5% | 3.2% | 3.9% | 58.3% |
| 1997 | 9.0% | 0.1% | 3.7% | 0.2% | 0.0% | 0.4% | 1.2% | 1.8% | 0.0% | 0.0% | 0.0% | 3.3% | 1.1% | 0.9% | 78.3% |
| 1998 | 10.2% | 0.5% | 1.2% | 0.5% | 0.0% | 0.1% | 0.7% | 0.0% | 0.0% | 0.0% | 0.6% | 2.1% | 4.9% | 1.0% | 78.2% |
| 1999 | 13.6% | 5.0% | 3.0% | 0.3% | 0.0% | 0.6% | 3.8% | 0.5% | 0.0% | 0.0% | 5.2% | 9.1% | 1.0% | 3.3% | 54.4% |
| 2000 | 25.7% | 2.3% | 3.5% | 0.4% | 0.0% | 0.0% | 1.9% | 4.2% | 0.7% | 0.1% | 5.3% | 3.3% | 1.0% | 3.9% | 47.8% |
| 2001 | 16.4% | 5.9% | 1.5% | 0.5% | 0.0% | 0.0% | 1.6% | 11.2% | 0.2% | 0.0% | 4.4% | 17.6% | 0.7% | 6.5% | 33.6% |
| 2002 | 23.5% | 0.1% | 1.5% | 10.7% | 0.0% | 0.0% | 2.6% | 15.2% | 0.1% | 0.0% | 0.9% | 9.0% | 1.0% | 6.0% | 29.2% |
| 2003 | 26.2% | 1.8% | 1.1% | 11.1% | 0.0% | 0.0% | 5.9% | 11.3% | 0.0% | 0.0% | 0.9% | 6.5% | 2.7% | 6.7% | 25.9% |
| 2004 | 14.6% | 0.7% | 1.1% | 4.9% | 0.0% | 0.0% | 1.9% | 11.3% | 0.2% | 0.0% | 1.6% | 10.6% | 7.7% | 16.3% | 29.1% |
| (79-80) | 23.6% | 0.0% | 0.9% | 9.1% | 4.1% | 4.8% | 0.0% | 18.5% | 3.5% | 0.7% | 0.0% | 1.1% | 2.3% | 2.2% | 29.1% |
| (87-98) | 10.7% | 0.5% | 0.5% | 3.4% | 0.7% | 1.9% | 1.8% | 9.5% | 0.4% | 0.5% | 0.8% | 5.8% | 6.1% | 1.6% | 55.9% |
| (99-04) | 20.0% | 2.6% | 1.9% | 4.7% | 0.0% | 0.1% | 2.9% | 9.0% | 0.2% | 0.0% | 3.1% | 9.3% | 2.4% | 7.1% | 36.7% |

Appendix E.53. Percent distribution of Cowlitz Tule Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1981 | 5.6% | 0.0% | 0.0% | 2.4% | 0.0% | 1.3% | 6.5% | 16.1% | 0.0% | 2.4% | 0.0% | 9.7% | 15.1% | 12.9% | 28.0% |
| 1982 | 3.7% | 0.0% | 0.2% | 1.4% | 0.5% | 2.1% | 0.0% | 14.5% | 0.0% | 1.2% | 0.9% | 18.5% | 9.7% | 12.5% | 34.9% |
| 1983 | 3.7% | 0.0% | 0.0% | 6.7% | 3.7% | 0.5% | 0.0% | 17.8% | 0.4% | 0.5% | 0.0% | 6.9% | 4.8% | 18.7% | 36.2% |
| 1984 | 4.4% | 0.0% | 0.0% | 7.2% | 2.1% | 0.1% | 0.8% | 24.5% | 0.0% | 1.7% | 0.0% | 4.4% | 15.1% | 3.6% | 36.0% |
| 1985 | 3.7% | 0.3% | 0.0% | 4.0% | 0.0% | 4.4% | 0.0% | 11.4% | 0.4% | 1.2% | 0.0% | 4.4% | 6.5% | 13.7% | 49.9% |
| 1986 | 0.4% | 0.1% | 0.0% | 0.2% | 0.6% | 0.8% | 0.0% | 12.6% | 0.4% | 1.1% | 0.0% | 13.0% | 31.0% | 12.4% | 27.4% |
| 1987 | 3.7% | 0.3% | 0.0% | 3.9% | 1.2% | 0.0% | 0.0% | 9.7% | 0.0% | 0.8% | 1.0% | 11.4% | 22.9% | 16.1% | 29.0% |
| 1988 | 1.7% | 0.3% | 0.0% | 1.9% | 0.0% | 0.1% | 0.0% | 15.9% | 0.0% | 0.6% | 0.0% | 15.5% | 24.0% | 12.3% | 27.7% |
| 1989 | 3.3% | 0.0% | 0.7% | 4.5% | 0.0% | 0.3% | 0.0% | 6.6% | 0.0% | 1.0% | 0.0% | 17.9% | 7.1% | 10.6% | 47.7% |
| 1990 | 4.4% | 0.0% | 0.0% | 1.8% | 2.9% | 2.6% | 0.0% | 14.2% | 0.0% | 0.7% | 0.0% | 9.5% | 0.0% | 12.0% | 51.8% |
| 1991 | 9.7% | 0.0% | 0.0% | 3.2% | 1.6% | 0.0% | 0.0% | 5.6% | 0.0% | 0.0% | 3.2% | 10.5% | 11.3% | 9.7% | 45.2% |
| 1992 | 2.2% | 0.0% | 0.0% | 0.0% | 2.2% | 0.0% | 1.6% | 17.7% | 0.0% | 0.0% | 0.0% | 7.0% | 5.4% | 4.8% | 59.1% |
| 1993 | 3.4% | 0.0% | 0.0% | 2.5% | 0.0% | 0.9% | 0.0% | 6.7% | 0.0% | 0.0% | 0.0% | 17.5% | 3.1% | 22.4% | 43.6% |
| 1994 | 4.2% | 0.0% | 0.0% | 1.9% | 0.0% | 0.0% | 0.0% | 1.9% | 0.0% | 0.0% | 0.0% | 3.3% | 0.0% | 0.0% | 88.7% |
| 1995 | 0.6% | 0.0% | 0.0% | 1.8% | 0.0% | 1.2% | 0.0% | 1.8% | 0.0% | 0.0% | 2.4% | 4.7% | 2.4% | 1.8% | 83.4% |
| 1996 | 4.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.2% | 0.0% | 0.0% | 5.9% | 1.1% | 3.7% | 83.0% |
| 1997 | 4.9% | 0.0% | 9.8% | 3.0% | 0.0% | 0.0% | 0.0% | 4.9% | 2.4% | 0.0% | 0.0% | 5.5% | 0.0% | 1.2% | 68.3% |
| 1998 | 3.7% | 0.0% | 0.0% | 7.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.9% | 0.0% | 2.5% | 76.5% |
| 1999 | 4.5% | 0.0% | 3.8% | 0.0% | 0.0% | 0.0% | 2.3% | 3.8% | 0.0% | 0.0% | 0.0% | 9.0% | 0.0% | 18.0% | 58.6% |
| 2000 | 3.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.2% | 0.0% | 0.0% | 12.4% | 13.4% | 5.2% | 7.2% | 51.5% |
| 2001 | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 0.0% | 2.9% | 10.5% | 1.5% | 11.9% | 71.0% |
| 2002 | 6.4% | 0.0% | 0.0% | 0.8% | 0.0% | 0.0% | 0.0% | 7.2% | 0.0% | 0.0% | 1.9% | 26.2% | 3.5% | 25.8% | 28.3% |
| 2003 | 5.0% | 0.0% | 0.0% | 1.3% | 0.0% | 0.0% | 0.0% | 9.6% | 0.7% | 0.0% | 6.7% | 16.5% | 8.5% | 10.9% | 40.9% |
| 2004 | 4.5% | 0.0% | 0.0% | 1.0% | 0.0% | 0.0% | 0.0% | 6.4% | 0.0% | 0.0% | 0.0% | 17.8% | 9.4% | 13.9% | 47.0% |
| (81-84) | 4.4% | 0.0% | 0.1% | 4.4% | 1.6% | 1.0% | 1.8% | 18.2% | 0.1% | 1.5% | 0.2% | 9.9% | 11.2% | 11.9% | 33.8% |
| (85-98) | 3.6% | 0.1% | 0.7% | 2.6% | 0.6% | 0.7% | 0.1% | 7.8% | 0.4% | 0.4% | 0.5% | 9.7% | 8.2% | 8.8% | 55.8% |
| (99-04) | 4.1% | 0.0% | 0.6% | 0.5% | 0.0% | 0.0% | 0.4% | 5.9% | 0.1% | 0.0% | 4.0% | 15.6% | 4.7% | 14.6% | 49.6% |

Appendix E.54. Percent distribution of Cowlitz Tule Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1981 | 6.0% | 0.0% | 0.0% | 2.4% | 0.0% | 1.2% | 6.3% | 18.8% | 0.0% | 2.2% | 0.0% | 11.3% | 14.2% | 12.7% | 25.0% |
| 1982 | 4.3% | 0.0% | 0.4% | 1.6% | 0.4% | 2.2% | 0.0% | 16.8% | 0.0% | 1.2% | 1.0% | 20.2% | 9.5% | 12.6% | 29.8% |
| 1983 | 4.4% | 0.0% | 0.0% | 7.2% | 3.9% | 0.5% | 0.0% | 18.9% | 0.3% | 0.5% | 0.0% | 7.8% | 4.7% | 18.7% | 33.2% |
| 1984 | 4.5% | 0.0% | 0.0% | 7.5% | 2.3% | 0.1% | 0.9% | 25.6% | 0.0% | 1.8% | 0.0% | 4.7% | 14.8% | 3.7% | 34.2% |
| 1985 | 4.0% | 1.1% | 0.0% | 4.4% | 0.0% | 4.4% | 0.0% | 12.6% | 0.4% | 1.2% | 0.0% | 5.1% | 6.3% | 14.9% | 45.4% |
| 1986 | 0.5% | 0.2% | 0.0% | 0.2% | 0.7% | 0.8% | 0.0% | 14.0% | 0.3% | 1.0% | 0.0% | 14.6% | 30.1% | 12.7% | 24.9% |
| 1987 | 6.0% | 0.7% | 0.0% | 4.6% | 1.4% | 0.0% | 0.0% | 11.2% | 0.0% | 0.7% | 0.9% | 12.1% | 21.2% | 15.5% | 25.6% |
| 1988 | 1.8% | 0.8% | 0.0% | 2.1% | 0.0% | 0.1% | 0.0% | 17.8% | 0.0% | 0.6% | 0.0% | 16.0% | 22.7% | 12.5% | 25.7% |
| 1989 | 4.6% | 0.0% | 0.7% | 4.7% | 0.0% | 0.3% | 0.0% | 7.2% | 0.0% | 1.0% | 0.0% | 18.8% | 6.9% | 11.0% | 44.8% |
| 1990 | 4.4% | 0.0% | 0.0% | 2.4% | 3.4% | 2.7% | 0.0% | 15.5% | 0.0% | 1.0% | 0.0% | 10.1% | 0.0% | 12.8% | 47.8% |
| 1991 | 12.4% | 0.0% | 0.0% | 3.6% | 1.5% | 0.0% | 0.0% | 6.6% | 0.0% | 0.0% | 2.9% | 11.7% | 10.9% | 9.5% | 40.9% |
| 1992 | 2.5% | 0.0% | 0.0% | 0.0% | 2.5% | 0.0% | 2.0% | 20.2% | 0.0% | 0.0% | 0.0% | 7.9% | 5.4% | 5.4% | 54.2% |
| 1993 | 4.3% | 0.0% | 0.0% | 3.0% | 0.0% | 1.1% | 0.0% | 7.6% | 0.0% | 0.0% | 0.0% | 18.7% | 3.0% | 23.8% | 38.5% |
| 1994 | 5.1% | 0.0% | 0.0% | 2.3% | 0.0% | 0.0% | 0.0% | 2.3% | 0.0% | 0.0% | 0.0% | 3.2% | 0.0% | 0.0% | 87.1% |
| 1995 | 1.1% | 0.0% | 0.0% | 2.8% | 0.0% | 1.1% | 0.0% | 2.3% | 0.0% | 2.3% | 2.3% | 4.5% | 2.3% | 1.7% | 79.7% |
| 1996 | 5.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.5% | 0.0% | 0.0% | 6.1% | 1.1% | 4.0% | 80.9% |
| 1997 | 5.7% | 0.0% | 10.8% | 3.4% | 0.0% | 0.0% | 0.0% | 5.7% | 2.8% | 1.1% | 0.0% | 5.7% | 0.0% | 1.1% | 63.6% |
| 1998 | 4.8% | 0.0% | 0.0% | 8.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 10.7% | 0.0% | 2.4% | 73.8% |
| 1999 | 6.9% | 0.0% | 4.1% | 0.0% | 0.0% | 0.0% | 2.8% | 3.4% | 0.0% | 0.0% | 0.0% | 9.7% | 0.0% | 19.3% | 53.8% |
| 2000 | 3.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.4% | 0.0% | 0.0% | 13.9% | 16.7% | 4.6% | 7.4% | 46.3% |
| 2001 | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 0.0% | 3.4% | 12.0% | 1.5% | 13.0% | 67.9% |
| 2002 | 7.1% | 0.0% | 0.0% | 0.9% | 0.0% | 0.0% | 0.0% | 6.7% | 0.0% | 0.0% | 2.1% | 28.0% | 3.4% | 26.0% | 25.8% |
| 2003 | 5.1% | 0.0% | 0.0% | 1.4% | 0.0% | 0.0% | 0.0% | 9.4% | 0.9% | 0.0% | 7.7% | 17.5% | 8.2% | 11.3% | 38.6% |
| 2004 | 5.5% | 0.0% | 0.0% | 0.9% | 0.0% | 0.0% | 0.0% | 6.0% | 0.0% | 0.0% | 0.0% | 19.4% | 9.2% | 15.2% | 43.8% |
| (81-84) | 4.8% | 0.0% | 0.1% | 4.7% | 1.6% | 1.0% | 1.8% | 20.0% | 0.1% | 1.4% | 0.2% | 11.0% | 10.8% | 11.9% | 30.6% |
| (85-98) | 4.5% | 0.2% | 0.8% | 3.0% | 0.7% | 0.8% | 0.1% | 8.8% | 0.4% | 0.6% | 0.4% | 10.4% | 7.8% | 9.1% | 52.3% |
| (99-04) | 4.9% | 0.0% | 0.7% | 0.5% | 0.0% | 0.0% | 0.5% | 5.7% | 0.1% | 0.0% | 4.5% | 17.2% | 4.5% | 15.4% | 46.0% |

Appendix E.55. Percent distribution of Spring Creek Tule Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 0.0% | 0.0% | 0.0% | 0.1% | 0.7% | 0.3% | 0.0% | 24.0% | 1.5% | 2.4% | 0.1% | 16.6% | 23.5% | 12.8% | 18.3% |
| 1980 | 0.1% | 0.0% | 0.0% | 0.1% | 0.5% | 0.1% | 0.0% | 25.4% | 2.8% | 1.0% | 0.1% | 23.6% | 23.7% | 10.1% | 12.6% |
| 1981 | 0.0% | 0.0% | 0.0% | 0.1% | 0.2% | 0.1% | 0.0% | 21.0% | 1.5% | 1.9% | 0.1% | 23.5% | 20.7% | 12.6% | 18.3% |
| 1982 | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 22.0% | 1.0% | 0.2% | 0.0% | 19.6% | 35.6% | 8.3% | 12.7% |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 29.8% | 1.1% | 0.0% | 0.5% | 8.4% | 20.2% | 9.8% | 29.7% |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 2.4% | 0.0% | 0.0% | 27.5% | 0.0% | 1.3% | 0.4% | 6.0% | 25.9% | 7.4% | 29.1% |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 14.2% | 0.0% | 0.2% | 0.7% | 13.8% | 27.2% | 4.0% | 39.7% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 2.9% | 0.0% | 0.0% | 20.6% | 1.9% | 1.6% | 2.5% | 2.5% | 36.2% | 7.9% | 23.8% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 7.9% | 0.0% | 0.0% | 0.0% | 14.0% | 38.6% | 20.2% | 19.3% |
| 1988 | 0.0% | 0.0% | 0.0% | 0.5% | 0.3% | 0.2% | 0.0% | 23.2% | 0.9% | 1.9% | 2.2% | 18.3% | 31.0% | 10.3% | 11.3% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 14.4% | 0.4% | 0.4% | 3.3% | 24.8% | 34.5% | 8.3% | 13.8% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.2% | 0.3% | 0.1% | 0.0% | 17.6% | 0.7% | 0.8% | 4.5% | 14.3% | 23.0% | 13.1% | 25.3% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.1% | 0.0% | 13.1% | 0.2% | 0.4% | 1.3% | 16.9% | 34.2% | 11.0% | 22.5% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 11.9% | 0.6% | 0.5% | 2.5% | 26.5% | 14.7% | 11.8% | 31.3% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 17.7% | 0.0% | 0.4% | 4.2% | 17.7% | 21.4% | 10.5% | 28.2% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 18.6% | 0.0% | 0.8% | 3.9% | 3.5% | 28.9% | 0.8% | 43.4% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.7% | 0.0% | 0.2% | 2.7% | 1.8% | 37.9% | 0.0% | 50.7% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.1% | 6.1% | 57.8% | 3.3% | 29.7% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 11.9% | 0.0% | 0.0% | 2.7% | 5.4% | 24.3% | 11.7% | 44.0% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.5% | 2.8% | 15.0% | 12.8% | 68.5% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.3% | 0.0% | 3.8% | 16.9% | 36.5% | 9.3% | 33.0% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.7% | 0.0% | 0.0% | 6.2% | 5.4% | 21.9% | 9.7% | 53.1% |
| 2001 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.2% | 0.4% | 0.0% | 1.1% | 18.7% | 30.8% | 7.2% | 37.4% |
| 2002 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.3% | 0.2% | 0.0% | 0.7% | 14.0% | 21.5% | 9.2% | 45.1% |
| 2003 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.8% | 0.0% | 0.0% | 7.0% | 10.2% | 21.1% | 5.4% | 46.6% |
| 2004 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 11.5% | 0.0% | 0.0% | 3.1% | 8.7% | 14.9% | 5.3% | 56.6% |
| (79-84) | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 0.1% | 0.0% | 25.0% | 1.3% | 1.1% | 0.2% | 16.3% | 24.9% | 10.1% | 20.1% |
| (85-98) | 0.0% | 0.0% | 0.0% | 0.1% | 0.3% | 0.0% | 0.0% | 12.7% | 0.3% | 0.5% | 2.4% | 12.0% | 30.3% | 9.0% | 32.2% |
| (99-04) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.5% | 0.1% | 0.0% | 3.6% | 12.3% | 24.5% | 7.7% | 45.3% |

Appendix E.56. Percent distribution of Spring Creek Tule Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 0.0% | 0.0% | 0.0% | 0.1% | 0.8% | 0.2% | 0.0% | 27.3% | 1.3% | 2.2% | 0.1% | 18.0% | 21.5% | 13.3% | 15.2% |
| 1980 | 0.1% | 0.0% | 0.0% | 0.1% | 0.6% | 0.1% | 0.0% | 27.8% | 2.5% | 0.9% | 0.1% | 24.7% | 21.9% | 10.7% | 10.6% |
| 1981 | 0.0% | 0.0% | 0.0% | 0.1% | 0.2% | 0.1% | 0.0% | 22.9% | 1.4% | 1.8% | 0.1% | 24.7% | 19.7% | 12.9% | 16.1% |
| 1982 | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 25.0% | 1.0% | 0.2% | 0.0% | 21.4% | 32.9% | 8.0% | 11.1% |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 31.5% | 1.1% | 0.0% | 0.5% | 9.1% | 18.9% | 12.1% | 26.4% |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 2.4% | 0.0% | 0.0% | 27.2% | 0.0% | 1.2% | 0.3% | 6.1% | 24.6% | 12.7% | 25.5% |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 15.3% | 0.0% | 0.2% | 0.6% | 16.0% | 27.0% | 4.1% | 36.6% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 2.9% | 0.0% | 0.0% | 21.8% | 1.8% | 1.8% | 2.7% | 2.7% | 35.4% | 8.8% | 22.1% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.9% | 0.0% | 0.0% | 0.0% | 15.2% | 40.4% | 19.9% | 14.6% |
| 1988 | 0.0% | 0.0% | 0.0% | 0.5% | 0.2% | 0.2% | 0.0% | 26.8% | 1.0% | 1.5% | 2.2% | 18.8% | 27.3% | 12.6% | 8.9% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 16.5% | 0.5% | 0.4% | 3.2% | 26.7% | 31.9% | 8.8% | 11.8% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.2% | 0.4% | 0.1% | 0.0% | 19.9% | 0.7% | 0.8% | 4.5% | 15.5% | 21.1% | 14.9% | 21.7% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.1% | 0.0% | 15.2% | 0.3% | 0.4% | 1.3% | 18.6% | 32.0% | 12.2% | 19.6% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 14.0% | 0.7% | 0.5% | 2.4% | 28.7% | 13.8% | 12.3% | 27.5% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 19.7% | 0.0% | 0.3% | 4.2% | 19.2% | 19.8% | 11.7% | 25.0% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 22.0% | 0.0% | 0.9% | 4.0% | 3.5% | 28.6% | 1.1% | 39.9% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 10.1% | 0.0% | 0.4% | 2.8% | 1.8% | 37.8% | 0.0% | 47.1% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.4% | 0.0% | 0.0% | 3.2% | 6.0% | 57.9% | 3.9% | 27.7% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 14.7% | 0.0% | 0.0% | 2.6% | 5.8% | 23.5% | 13.2% | 40.2% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 0.6% | 3.3% | 15.3% | 16.8% | 63.7% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.3% | 0.0% | 3.8% | 19.2% | 35.8% | 10.7% | 29.9% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.9% | 0.0% | 0.0% | 7.3% | 6.1% | 21.0% | 15.1% | 46.7% |
| 2001 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.3% | 0.5% | 0.0% | 1.2% | 21.1% | 30.0% | 9.3% | 33.7% |
| 2002 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.4% | 0.2% | 0.0% | 0.8% | 16.4% | 21.3% | 10.2% | 41.7% |
| 2003 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.9% | 0.0% | 0.0% | 8.2% | 11.3% | 20.9% | 5.9% | 43.8% |
| 2004 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 11.7% | 0.0% | 0.0% | 3.5% | 10.3% | 14.9% | 5.9% | 53.7% |
| (79-84) | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 0.1% | 0.0% | 26.9% | 1.2% | 1.1% | 0.2% | 17.3% | 23.2% | 11.6% | 17.5% |
| (85-98) | 0.0% | 0.0% | 0.0% | 0.1% | 0.3% | 0.0% | 0.0% | 14.8% | 0.4% | 0.5% | 2.4% | 13.0% | 29.4% | 10.0% | 29.0% |
| (99-04) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.6% | 0.2% | 0.0% | 4.1% | 14.1% | 24.0% | 9.5% | 41.6% |

Appendix E.57. Percent distribution of Columbia Lower River Hatchery Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1980 | 0.8% | 0.0% | 0.0% | 0.0% | 0.5% | 1.3% | 0.0% | 16.0% | 3.4% | 6.4% | 1.3% | 18.3% | 9.8% | 22.4% | 19.8% |
| 1981 | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.1% | 0.0% | 30.6% | 1.8% | 2.4% | 0.3% | 22.6% | 1.9% | 11.6% | 28.2% |
| 1982 | 0.0% | 0.0% | 0.0% | 0.3% | 1.8% | 0.0% | 0.0% | 26.0% | 0.8% | 0.3% | 0.5% | 18.6% | 16.4% | 9.0% | 26.5% |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 2.3% | 0.3% | 0.1% | 35.0% | 1.4% | 0.6% | 0.4% | 11.2% | 6.8% | 8.5% | 33.4% |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 3.2% | 0.0% | 0.0% | 49.9% | 1.3% | 1.6% | 0.3% | 5.9% | 11.3% | 3.7% | 22.7% |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.4% | 0.0% | 28.2% | 1.1% | 1.2% | 0.7% | 15.6% | 4.1% | 5.8% | 41.9% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 0.1% | 9.1% | 2.5% | 7.5% | 2.7% | 6.9% | 11.2% | 11.5% | 47.9% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.2% | 1.6% | 0.0% | 0.0% | 26.9% | 0.5% | 0.2% | 2.5% | 16.6% | 20.7% | 9.5% | 21.3% |
| 1988 | 0.3% | 0.0% | 0.0% | 0.3% | 0.6% | 0.0% | 0.0% | 28.8% | 1.0% | 0.0% | 2.4% | 11.5% | 24.3% | 3.2% | 27.6% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 15.4% | 0.0% | 2.0% | 0.0% | 22.4% | 5.9% | 5.1% | 49.2% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 19.8% | 0.0% | 1.7% | 0.0% | 16.3% | 0.3% | 11.1% | 50.3% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 10.2% | 0.7% | 2.5% | 2.0% | 9.3% | 2.3% | 14.9% | 57.9% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 16.3% | 0.0% | 1.0% | 1.9% | 28.0% | 0.8% | 11.0% | 40.5% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 18.4% | 0.0% | 0.0% | 4.5% | 19.7% | 2.0% | 11.1% | 43.6% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 27.6% | 10.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 62.1% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.3% | 10.0% | 86.7% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.1% | 6.5% | 0.0% | 85.5% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.4% | 2.9% | 0.0% | 3.9% | 8.7% | 1.0% | 11.6% | 55.6% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.0% | 1.0% | 0.0% | 0.0% | 5.1% | 1.0% | 2.0% | 23.2% | 63.6% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.3% | 0.0% | 0.0% | 9.1% | 6.8% | 3.6% | 9.4% | 68.7% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 15.9% | 2.2% | 0.0% | 16.4% | 2.2% | 2.6% | 4.3% | 56.5% |
| 2001 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.2% | 0.2% | 0.0% | 3.4% | 19.2% | 1.5% | 8.6% | 58.9% |
| 2002 | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 10.6% | 0.0% | 0.0% | 1.8% | 20.9% | 8.6% | 11.0% | 46.7% |
| 2003 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 14.9% | 0.3% | 0.0% | 13.4% | 14.6% | 6.5% | 9.4% | 40.9% |
| 2004 | 0.5% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.8% | 20.9% | 0.5% | 0.0% | 9.6% | 8.5% | 18.5% | 5.0% | 35.4% |
| (80-84) | 0.2% | 0.0% | 0.0% | 0.1% | 1.7% | 0.3% | 0.0% | 31.5% | 1.7% | 2.3% | 0.6% | 15.3% | 9.2% | 11.0% | 26.1% |
| (85-98) | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.1% | 0.3% | 15.6% | 1.4% | 1.1% | 1.8% | 11.7% | 6.0% | 9.1% | 52.4% |
| (99-04) | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 12.1% | 0.5% | 0.0% | 9.0% | 12.0% | 6.9% | 8.0% | 51.2% |

Appendix E.58. Percent distribution of Columbia Lower River Hatchery Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1980 | 0.4% | 0.0% | 0.0% | 0.1% | 0.8% | 0.8% | 0.0% | 32.4% | 2.0% | 4.2% | 0.7% | 23.1% | 6.7% | 17.7% | 10.9% |
| 1981 | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.1% | 0.0% | 33.4% | 1.6% | 2.2% | 0.3% | 25.0% | 1.8% | 11.5% | 23.6% |
| 1982 | 0.0% | 0.0% | 0.0% | 0.3% | 2.0% | 0.0% | 0.0% | 29.2% | 0.8% | 0.3% | 0.5% | 20.0% | 15.2% | 8.9% | 22.9% |
| 1983 | 0.0% | 0.0% | 0.0% | 0.0% | 2.4% | 0.3% | 0.1% | 37.0% | 1.3% | 0.5% | 0.4% | 12.3% | 6.7% | 9.6% | 29.4% |
| 1984 | 0.0% | 0.0% | 0.0% | 0.0% | 3.4% | 0.0% | 0.0% | 51.6% | 1.3% | 1.6% | 0.2% | 6.3% | 11.1% | 4.1% | 20.4% |
| 1985 | 0.0% | 0.0% | 0.0% | 0.0% | 0.9% | 0.4% | 0.0% | 30.3% | 1.1% | 1.2% | 0.7% | 17.7% | 4.1% | 5.9% | 37.7% |
| 1986 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.2% | 8.5% | 1.9% | 6.3% | 2.5% | 6.3% | 9.5% | 30.0% | 34.1% |
| 1987 | 0.0% | 0.0% | 0.0% | 0.2% | 1.9% | 0.0% | 0.0% | 33.0% | 0.5% | 0.2% | 2.2% | 17.3% | 18.4% | 8.6% | 17.6% |
| 1988 | 0.3% | 0.0% | 0.0% | 0.3% | 0.6% | 0.0% | 0.0% | 31.6% | 1.0% | 0.0% | 2.4% | 11.7% | 23.1% | 3.3% | 25.8% |
| 1989 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 17.0% | 0.0% | 1.8% | 0.0% | 25.3% | 5.4% | 5.4% | 45.1% |
| 1990 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.0% | 22.8% | 0.0% | 1.5% | 0.0% | 18.2% | 0.3% | 12.0% | 44.8% |
| 1991 | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.0% | 12.3% | 1.0% | 2.4% | 2.2% | 10.9% | 2.4% | 18.2% | 50.6% |
| 1992 | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 19.5% | 0.0% | 0.8% | 1.8% | 30.3% | 0.7% | 11.3% | 34.9% |
| 1993 | 0.0% | 0.0% | 0.0% | 0.0% | 0.7% | 0.0% | 0.0% | 20.8% | 0.0% | 0.0% | 4.3% | 20.8% | 1.9% | 11.6% | 39.9% |
| 1994 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 31.3% | 12.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 56.3% |
| 1995 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.1% | 0.0% | 0.0% | 3.1% | 12.5% | 81.3% |
| 1996 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.1% | 6.5% | 0.0% | 85.5% |
| 1997 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 20.2% | 3.1% | 0.4% | 3.5% | 9.2% | 0.9% | 12.3% | 50.4% |
| 1998 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 6.5% | 0.9% | 0.0% | 0.0% | 5.6% | 0.9% | 1.9% | 25.9% | 58.3% |
| 1999 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.2% | 0.0% | 0.0% | 9.6% | 8.0% | 3.7% | 11.1% | 65.3% |
| 2000 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.3% | 2.7% | 0.0% | 19.0% | 2.3% | 2.3% | 7.6% | 49.8% |
| 2001 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.3% | 0.2% | 0.0% | 3.8% | 22.0% | 1.5% | 10.6% | 53.7% |
| 2002 | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 10.6% | 0.0% | 0.0% | 2.1% | 24.0% | 8.3% | 11.8% | 42.6% |
| 2003 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 14.2% | 0.4% | 0.0% | 15.2% | 16.2% | 6.4% | 9.8% | 37.7% |
| 2004 | 0.5% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 1.1% | 20.6% | 0.6% | 0.0% | 10.8% | 8.9% | 18.1% | 5.3% | 33.8% |
| (80-84) | 0.1% | 0.0% | 0.0% | 0.1% | 1.8% | 0.2% | 0.0% | 36.7% | 1.4% | 1.8% | 0.4% | 17.4% | 8.3% | 10.3% | 21.4% |
| (85-98) | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.1% | 0.5% | 17.7% | 1.5% | 1.3% | 1.8% | 12.6% | 5.6% | 11.2% | 47.3% |
| (99-04) | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 12.0% | 0.6% | 0.0% | 10.1% | 13.6% | 6.7% | 9.4% | 47.2% |

Appendix E.59. Percent distribution of Upriver Bright Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 18.0% | 0.3% | 0.6% | 7.6% | 4.0% | 3.7% | 0.1% | 11.8% | 0.5% | 0.7% | 0.0% | 1.3% | 23.0% | 1.8% | 26.7% |
| 1980 | 19.9% | 0.6% | 0.5% | 6.5% | 1.6% | 1.7% | 0.1% | 7.3% | 1.0% | 0.2% | 0.0% | 1.1% | 6.3% | 1.8% | 51.4% |
| 1981 | 16.1% | 0.0% | 0.4% | 5.6% | 1.1% | 1.3% | 0.0% | 3.8% | 0.4% | 0.5% | 0.2% | 0.5% | 3.6% | 1.0% | 65.8% |
| 1982 | 6.4% | 0.4% | 0.2% | 3.5% | 0.2% | 1.1% | 0.1% | 4.6% | 0.0% | 0.4% | 0.0% | 0.6% | 2.5% | 0.7% | 79.2% |
| 1983 | 15.5% | 0.2% | 0.0% | 10.7% | 1.8% | 3.4% | 0.2% | 3.7% | 0.2% | 0.1% | 0.0% | 0.4% | 8.1% | 0.0% | 55.6% |
| 1984 | 14.5% | 1.1% | 0.1% | 8.6% | 2.0% | 1.5% | 0.2% | 7.2% | 0.2% | 0.8% | 0.2% | 0.2% | 15.3% | 1.9% | 46.3% |
| 1985 | 9.2% | 1.2% | 0.2% | 8.8% | 0.8% | 1.3% | 0.0% | 7.9% | 0.1% | 1.2% | 0.1% | 0.4% | 32.8% | 4.5% | 31.5% |
| 1986 | 10.3% | 0.7% | 0.1% | 7.9% | 1.2% | 1.0% | 0.0% | 6.3% | 0.1% | 0.2% | 0.1% | 0.7% | 33.1% | 2.4% | 35.8% |
| 1987 | 14.6% | 0.4% | 0.4% | 12.4% | 1.8% | 0.6% | 0.1% | 7.8% | 0.0% | 0.1% | 0.3% | 1.5% | 35.2% | 3.7% | 21.2% |
| 1988 | 10.2% | 0.8% | 0.5% | 7.4% | 0.6% | 0.6% | 0.0% | 11.2% | 0.0% | 0.1% | 0.0% | 2.1% | 47.0% | 2.6% | 16.9% |
| 1989 | 11.9% | 0.0% | 0.2% | 14.9% | 0.2% | 0.7% | 0.6% | 7.7% | 0.0% | 0.7% | 0.0% | 1.2% | 42.5% | 2.0% | 17.3% |
| 1990 | 13.6% | 0.0% | 1.0% | 9.9% | 0.7% | 0.7% | 0.0% | 8.1% | 0.0% | 0.0% | 0.0% | 1.2% | 33.8% | 2.4% | 28.6% |
| 1991 | 6.3% | 0.4% | 2.6% | 5.9% | 0.0% | 0.0% | 0.0% | 8.9% | 0.0% | 0.0% | 0.0% | 0.7% | 19.6% | 4.4% | 51.1% |
| 1992 | 3.0% | 0.0% | 0.0% | 3.0% | 0.0% | 2.3% | 0.0% | 11.5% | 0.0% | 0.7% | 1.0% | 0.0% | 17.0% | 6.6% | 55.1% |
| 1993 | 10.9% | 0.0% | 0.0% | 6.7% | 0.0% | 0.4% | 0.6% | 17.0% | 0.0% | 0.0% | 0.0% | 1.7% | 15.7% | 6.5% | 40.4% |
| 1994 | 9.8% | 0.9% | 0.0% | 8.0% | 0.2% | 0.9% | 1.7% | 6.9% | 0.0% | 0.0% | 0.7% | 0.0% | 14.2% | 3.5% | 53.1% |
| 1995 | 8.1% | 0.1% | 1.7% | 2.0% | 0.0% | 0.4% | 0.0% | 5.3% | 0.0% | 0.0% | 0.0% | 0.7% | 9.9% | 4.3% | 67.3% |
| 1996 | 2.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.3% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.8% | 22.4% | 5.4% | 68.0% |
| 1997 | 11.1% | 0.3% | 2.5% | 4.5% | 0.2% | 0.0% | 0.6% | 0.5% | 0.0% | 0.0% | 0.1% | 1.0% | 20.6% | 11.4% | 47.2% |
| 1998 | 8.1% | 1.5% | 2.2% | 2.6% | 0.0% | 0.0% | 0.5% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 13.7% | 6.4% | 64.9% |
| 1999 | 10.4% | 0.6% | 2.6% | 3.8% | 0.0% | 0.0% | 0.8% | 0.0% | 0.4% | 0.0% | 0.3% | 0.6% | 13.5% | 9.7% | 57.4% |
| 2000 | 16.8% | 0.1% | 2.4% | 0.0% | 0.0% | 0.0% | 0.4% | 0.9% | 0.0% | 0.0% | 2.7% | 0.3% | 21.1% | 4.6% | 50.7% |
| 2001 | 3.8% | 0.0% | 0.7% | 0.0% | 0.0% | 0.0% | 0.7% | 0.7% | 0.0% | 0.0% | 0.4% | 1.7% | 12.9% | 7.8% | 71.5% |
| 2002 | 14.4% | 0.0% | 2.3% | 0.8% | 0.0% | 0.0% | 1.0% | 1.4% | 0.3% | 0.1% | 0.3% | 1.7% | 18.2% | 8.5% | 51.1% |
| 2003 | 13.0% | 0.9% | 0.5% | 4.3% | 0.0% | 0.0% | 3.1% | 1.0% | 0.0% | 0.0% | 1.3% | 0.7% | 13.7% | 7.0% | 54.6% |
| 2004 | 8.6% | 1.2% | 0.5% | 2.6% | 0.0% | 0.0% | 2.0% | 2.4% | 0.0% | 0.0% | 0.4% | 0.8% | 16.8% | 7.0% | 57.6% |
| (79-84) | 15.1% | 0.5% | 0.3% | 7.1% | 1.8% | 2.1% | 0.1% | 6.4% | 0.4% | 0.5% | 0.1% | 0.7% | 9.8% | 1.2% | 54.2% |
| (85-98) | 9.3% | 0.4% | 0.8% | 6.7% | 0.4% | 0.7% | 0.3% | 7.1% | 0.0% | 0.2% | 0.2% | 0.9% | 25.5% | 4.7% | 42.7% |
| (99-04) | 11.2% | 0.5% | 1.5% | 1.9% | 0.0% | 0.0% | 1.3% | 1.1% | 0.1% | 0.0% | 0.9% | 0.9% | 16.0% | 7.4% | 57.1% |

Appendix E.60. Percent distribution of Upriver Bright Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1979 | 18.4% | 0.3% | 0.6% | 7.9% | 4.1% | 3.7% | 0.1% | 12.5% | 0.5% | 0.7% | 0.0% | 1.3% | 22.3% | 2.0% | 25.5% |
| 1980 | 20.8% | 0.6% | 0.6% | 7.0% | 1.7% | 1.7% | 0.1% | 7.8% | 1.0% | 0.2% | 0.0% | 1.1% | 6.2% | 1.9% | 49.2% |
| 1981 | 17.1% | 0.0% | 0.4% | 5.9% | 1.1% | 1.3% | 0.0% | 4.1% | 0.3% | 0.5% | 0.2% | 0.6% | 3.6% | 1.1% | 63.9% |
| 1982 | 8.9% | 0.4% | 0.3% | 4.4% | 0.3% | 1.1% | 0.2% | 5.5% | 0.0% | 0.5% | 0.0% | 0.8% | 2.5% | 0.7% | 74.5% |
| 1983 | 22.1% | 0.3% | 0.0% | 11.7% | 2.0% | 3.3% | 0.2% | 3.8% | 0.2% | 0.1% | 0.0% | 0.4% | 7.4% | 0.0% | 48.5% |
| 1984 | 17.6% | 1.2% | 0.2% | 9.8% | 2.2% | 1.4% | 0.2% | 8.2% | 0.2% | 0.8% | 0.2% | 0.2% | 14.4% | 2.3% | 41.0% |
| 1985 | 12.9% | 2.3% | 0.3% | 9.0% | 0.8% | 1.3% | 0.0% | 8.1% | 0.1% | 1.1% | 0.1% | 0.5% | 30.9% | 4.6% | 28.2% |
| 1986 | 12.2% | 1.5% | 0.1% | 8.1% | 1.3% | 1.0% | 0.0% | 6.7% | 0.1% | 0.2% | 0.1% | 0.8% | 31.9% | 2.7% | 33.4% |
| 1987 | 19.4% | 1.0% | 0.4% | 13.1% | 2.0% | 0.6% | 0.1% | 8.5% | 0.0% | 0.1% | 0.3% | 1.5% | 31.4% | 3.5% | 18.3% |
| 1988 | 11.5% | 2.1% | 0.5% | 7.9% | 0.6% | 0.6% | 0.0% | 12.4% | 0.0% | 0.1% | 0.0% | 2.2% | 44.0% | 2.7% | 15.5% |
| 1989 | 14.5% | 0.0% | 0.2% | 15.2% | 0.2% | 0.7% | 0.5% | 8.1% | 0.0% | 0.7% | 0.0% | 1.2% | 40.4% | 2.0% | 16.1% |
| 1990 | 14.2% | 0.0% | 1.1% | 10.8% | 0.8% | 0.7% | 0.0% | 8.7% | 0.0% | 0.0% | 0.0% | 1.3% | 32.6% | 2.5% | 27.2% |
| 1991 | 8.1% | 1.3% | 3.4% | 6.7% | 0.0% | 0.0% | 0.0% | 10.1% | 0.0% | 0.0% | 0.0% | 1.0% | 18.5% | 4.7% | 46.3% |
| 1992 | 3.6% | 0.0% | 0.0% | 3.6% | 0.0% | 2.4% | 0.0% | 13.4% | 0.0% | 0.6% | 1.2% | 0.0% | 16.7% | 7.3% | 51.1% |
| 1993 | 16.6% | 0.0% | 0.0% | 7.6% | 0.0% | 0.3% | 0.5% | 18.6% | 0.0% | 0.0% | 0.0% | 1.6% | 14.0% | 6.1% | 34.7% |
| 1994 | 11.8% | 1.8% | 0.0% | 8.5% | 0.2% | 1.0% | 1.7% | 7.3% | 0.0% | 0.0% | 0.6% | 0.0% | 13.6% | 3.6% | 49.9% |
| 1995 | 10.2% | 0.1% | 2.4% | 2.7% | 0.0% | 0.5% | 0.0% | 7.0% | 0.0% | 0.1% | 0.0% | 0.7% | 9.6% | 4.5% | 62.1% |
| 1996 | 4.4% | 0.0% | 0.0% | 1.4% | 0.0% | 0.2% | 0.5% | 0.7% | 0.0% | 0.0% | 0.0% | 0.7% | 22.1% | 6.2% | 63.7% |
| 1997 | 12.8% | 0.5% | 3.2% | 4.9% | 0.2% | 0.0% | 0.9% | 0.6% | 0.0% | 0.0% | 0.1% | 1.0% | 19.7% | 11.8% | 44.4% |
| 1998 | 10.0% | 4.6% | 2.8% | 3.0% | 0.0% | 0.0% | 0.6% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 13.2% | 7.0% | 58.8% |
| 1999 | 13.4% | 1.5% | 2.8% | 4.0% | 0.0% | 0.0% | 0.8% | 0.0% | 0.4% | 0.0% | 0.3% | 0.6% | 12.9% | 10.1% | 53.1% |
| 2000 | 22.4% | 0.1% | 3.3% | 0.0% | 0.0% | 0.0% | 0.7% | 1.1% | 0.0% | 0.0% | 3.3% | 0.3% | 19.1% | 4.5% | 45.2% |
| 2001 | 5.4% | 0.0% | 1.0% | 0.0% | 0.0% | 0.0% | 1.1% | 0.7% | 0.0% | 0.0% | 0.4% | 1.9% | 13.0% | 8.8% | 67.8% |
| 2002 | 16.3% | 0.0% | 2.6% | 0.9% | 0.0% | 0.0% | 1.2% | 1.3% | 0.4% | 1.1% | 0.3% | 1.9% | 17.5% | 8.9% | 47.5% |
| 2003 | 14.4% | 2.5% | 0.6% | 4.7% | 0.0% | 0.0% | 4.4% | 1.0% | 0.0% | 0.0% | 1.5% | 0.7% | 12.9% | 7.3% | 50.0% |
| 2004 | 10.7% | 3.8% | 0.6% | 3.1% | 0.0% | 0.0% | 2.6% | 2.3% | 0.0% | 0.0% | 0.4% | 0.9% | 16.0% | 7.4% | 52.4% |
| (79-84) | 17.5% | 0.5% | 0.3% | 7.8% | 1.9% | 2.1% | 0.1% | 7.0% | 0.4% | 0.5% | 0.1% | 0.7% | 9.4% | 1.3% | 50.4% |
| (85-98) | 11.6% | 1.1% | 1.0% | 7.3% | 0.4% | 0.7% | 0.3% | 7.9% | 0.0% | 0.2% | 0.2% | 0.9% | 24.2% | 4.9% | 39.3% |
| (99-04) | 13.8% | 1.3% | 1.8% | 2.1% | 0.0% | 0.0% | 1.8% | 1.1% | 0.1% | 0.2% | 1.0% | 1.1% | 15.2% | 7.8% | 52.7% |

Appendix E.61. Percent distribution of Hanford Wild Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1990 | 8.4% | 0.5% | 0.0% | 4.3% | 0.5% | 0.5% | 0.0% | 8.4% | 0.0% | 0.2% | 3.6% | 0.5% | 22.5% | 7.0% | 43.6% |
| 1991 | 8.6% | 0.0% | 1.3% | 9.4% | 0.2% | 0.0% | 0.5% | 4.7% | 0.8% | 0.0% | 0.0% | 1.0% | 23.3% | 4.4% | 45.7% |
| 1992 | 16.4% | 1.7% | 1.4% | 5.9% | 0.0% | 0.0% | 0.0% | 16.0% | 0.0% | 0.0% | 0.0% | 1.0% | 18.5% | 2.8% | 36.2% |
| 1993 | 14.0% | 0.0% | 2.1% | 2.9% | 0.0% | 0.5% | 1.3% | 5.3% | 0.0% | 1.9% | 1.9% | 3.7% | 16.1% | 8.2% | 42.1% |
| 1994 | 14.4% | 0.8% | 0.0% | 4.8% | 0.3% | 1.1% | 0.0% | 4.4% | 0.0% | 0.3% | 0.0% | 0.7% | 12.4% | 5.4% | 55.3% |
| 1995 | 11.0% | 0.0% | 3.7% | 4.3% | 0.0% | 0.0% | 0.0% | 2.3% | 0.0% | 0.0% | 0.0% | 0.0% | 9.8% | 7.0% | 62.0% |
| 1996 | 9.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 28.4% | 7.8% | 53.5% |
| 1997 | 16.3% | 0.6% | 1.0% | 3.6% | 0.0% | 0.0% | 1.9% | 0.8% | 0.0% | 0.0% | 0.0% | 1.0% | 13.9% | 7.4% | 53.4% |
| 1998 | 12.8% | 0.0% | 0.0% | 8.5% | 0.0% | 0.0% | 1.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 17.3% | 6.4% | 53.5% |
| 1999 | 10.4% | 0.4% | 2.1% | 7.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 12.9% | 6.6% | 60.6% |
| 2000 | 16.4% | 0.5% | 1.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 29.1% | 5.5% | 46.8% |
| 2001 | 4.3% | 1.1% | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.1% | 20.4% | 14.4% | 57.8% |
| 2002 | 13.9% | 0.0% | 1.3% | 0.1% | 0.0% | 0.0% | 1.0% | 3.0% | 0.0% | 0.0% | 0.0% | 1.4% | 9.8% | 11.0% | 58.5% |
| 2003 | 10.5% | 0.0% | 0.7% | 3.2% | 0.0% | 0.0% | 0.4% | 0.0% | 0.0% | 0.0% | 0.9% | 0.5% | 12.7% | 7.7% | 63.2% |
| 2004 | 16.4% | 0.0% | 2.8% | 4.8% | 0.0% | 0.0% | 2.5% | 2.5% | 0.0% | 0.0% | 0.0% | 0.4% | 20.9% | 3.8% | 46.0% |
| (90-98) | 12.4% | 0.4% | 1.1% | 4.9% | 0.1% | 0.3% | 0.6% | 4.7% | 0.1% | 0.3% | 0.6% | 0.9% | 18.0% | 6.3% | 49.5% |
| (99-04) | 12.0% | 0.3% | 1.6% | 2.5% | 0.0% | 0.0% | 0.6% | 0.9% | 0.0% | 0.0% | 0.2% | 0.6% | 17.6% | 8.1% | 55.5% |

Appendix E.62. Percent distribution of Hanford Wild Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1990 | 9.3% | 1.1% | 0.4% | 5.1% | 0.4% | 0.4% | 0.0% | 8.9% | 0.0% | 0.2% | 3.6% | 0.6% | 21.7% | 7.4% | 40.8% |
| 1991 | 10.7% | 0.0% | 1.4% | 10.4% | 0.2% | 0.0% | 0.5% | 5.1% | 1.0% | 0.0% | 0.0% | 1.1% | 22.1% | 4.5% | 43.2% |
| 1992 | 18.1% | 5.4% | 1.5% | 6.9% | 0.0% | 0.0% | 0.0% | 16.9% | 0.0% | 0.0% | 0.0% | 0.9% | 16.3% | 2.4% | 31.4% |
| 1993 | 20.6% | 0.0% | 2.1% | 3.0% | 0.0% | 0.5% | 1.2% | 6.0% | 0.0% | 1.6% | 1.9% | 3.7% | 14.4% | 8.1% | 36.9% |
| 1994 | 17.5% | 1.9% | 0.0% | 5.2% | 0.3% | 1.0% | 0.0% | 4.7% | 0.0% | 0.3% | 0.0% | 0.6% | 11.7% | 5.5% | 51.2% |
| 1995 | 13.1% | 0.0% | 4.1% | 5.4% | 0.0% | 0.0% | 0.0% | 2.8% | 0.0% | 0.3% | 0.0% | 0.0% | 9.2% | 7.1% | 57.9% |
| 1996 | 13.0% | 0.0% | 0.0% | 0.2% | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 27.4% | 8.0% | 50.7% |
| 1997 | 17.9% | 1.2% | 1.1% | 3.6% | 0.0% | 0.0% | 2.4% | 0.9% | 0.0% | 0.2% | 0.0% | 0.9% | 13.4% | 7.7% | 50.8% |
| 1998 | 14.7% | 0.0% | 0.0% | 9.5% | 0.0% | 0.0% | 2.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.7% | 6.6% | 50.6% |
| 1999 | 13.7% | 1.5% | 2.3% | 7.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 12.2% | 6.9% | 55.7% |
| 2000 | 19.7% | 0.4% | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 27.8% | 5.6% | 44.0% |
| 2001 | 5.9% | 2.7% | 1.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.3% | 19.7% | 15.1% | 54.2% |
| 2002 | 17.8% | 0.0% | 1.4% | 0.1% | 0.0% | 0.0% | 1.1% | 2.9% | 0.0% | 0.0% | 0.0% | 1.6% | 9.3% | 11.3% | 54.5% |
| 2003 | 11.3% | 0.0% | 0.8% | 3.4% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 1.1% | 0.5% | 12.5% | 8.1% | 61.6% |
| 2004 | 17.9% | 0.0% | 2.9% | 5.2% | 0.0% | 0.0% | 3.3% | 2.5% | 0.0% | 0.0% | 0.0% | 0.4% | 20.1% | 3.9% | 44.0% |
| (90-98) | 15.0% | 1.1% | 1.2% | 5.5% | 0.1% | 0.3% | 0.7% | 5.0% | 0.1% | 0.3% | 0.6% | 0.9% | 17.0% | 6.4% | 45.9% |
| (99-04) | 14.4% | 0.8% | 1.8% | 2.7% | 0.0% | 0.0% | 0.8% | 0.9% | 0.0% | 0.0% | 0.2% | 0.6% | 16.9% | 8.5% | 52.3% |

Appendix E.63. Percent distribution of Lyons Ferry Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1988 | 2.8% | 0.0% | 0.0% | 3.3% | 0.6% | 0.8% | 0.0% | 18.6% | 0.0% | 0.3% | 0.0% | 10.8% | 29.7% | 3.9% | 29.4% |
| 1989 | 2.8% | 0.0% | 0.0% | 6.3% | 0.0% | 0.4% | 0.0% | 16.0% | 0.0% | 1.2% | 0.9% | 12.3% | 27.3% | 6.6% | 26.2% |
| 1990 | 5.3% | 0.0% | 0.0% | 3.5% | 0.0% | 0.5% | 0.0% | 16.1% | 0.0% | 0.0% | 0.0% | 9.6% | 26.4% | 5.8% | 32.8% |
| 1991 | 2.7% | 0.0% | 1.8% | 4.9% | 0.0% | 0.4% | 0.0% | 8.8% | 0.0% | 0.9% | 0.0% | 4.0% | 12.8% | 2.7% | 61.1% |
| 1992 | 1.2% | 1.2% | 0.0% | 3.6% | 0.0% | 1.2% | 0.0% | 10.7% | 0.0% | 1.2% | 3.0% | 5.9% | 8.3% | 1.8% | 62.1% |
| 1993 | 3.6% | 0.0% | 0.0% | 4.7% | 0.8% | 0.8% | 0.0% | 10.3% | 0.0% | 1.2% | 0.0% | 7.9% | 13.8% | 1.6% | 55.3% |
| 1994 | 6.2% | 0.5% | 1.4% | 6.0% | 0.7% | 0.5% | 0.0% | 7.1% | 0.7% | 2.2% | 0.0% | 0.0% | 7.3% | 0.5% | 66.8% |
| 2003 | 8.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 1.2% | 0.0% | 0.0% | 0.0% | 6.0% | 13.4% | 5.7% | 65.8% |
| 2004 | 3.6% | 0.0% | 0.0% | 2.1% | 0.0% | 0.0% | 2.3% | 2.5% | 0.0% | 0.0% | 1.9% | 7.9% | 8.7% | 6.2% | 64.9% |
| (88-94) | 3.5% | 0.2% | 0.5% | 4.6% | 0.3% | 0.7% | 0.0% | 12.5% | 0.1% | 1.0% | 0.5% | 7.2% | 17.9% | 3.3% | 47.7% |
| (03-04) | 5.8% | 0.0% | 0.0% | 1.0% | 0.0% | 0.0% | 1.1% | 1.8% | 0.0% | 0.0% | 0.9% | 6.9% | 11.0% | 5.9% | 65.3% |

Appendix E.64. Percent distribution of Lyons Ferry Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1988 | 3.2% | 0.0% | 0.1% | 4.0% | 0.6% | 0.7% | 0.0% | 21.1% | 0.0% | 0.2% | 0.0% | 11.6% | 27.7% | 4.0% | 26.8% |
| 1989 | 4.2% | 0.0% | 0.0% | 7.0% | 0.0% | 0.4% | 0.0% | 17.9% | 0.0% | 1.1% | 0.9% | 12.9% | 25.4% | 6.6% | 23.6% |
| 1990 | 5.5% | 0.0% | 0.0% | 3.7% | 0.0% | 0.5% | 0.0% | 17.1% | 0.0% | 0.0% | 0.0% | 10.0% | 25.7% | 6.3% | 31.3% |
| 1991 | 3.4% | 0.0% | 2.1% | 5.5% | 0.0% | 0.4% | 0.0% | 10.1% | 0.0% | 0.8% | 0.0% | 4.2% | 12.6% | 2.9% | 58.0% |
| 1992 | 1.6% | 5.2% | 0.0% | 4.2% | 0.0% | 1.6% | 0.0% | 12.0% | 0.0% | 1.0% | 3.1% | 6.3% | 7.9% | 2.1% | 55.0% |
| 1993 | 5.4% | 0.7% | 0.4% | 5.8% | 1.1% | 0.7% | 0.0% | 11.6% | 0.0% | 1.1% | 0.0% | 8.0% | 13.0% | 1.4% | 50.7% |
| 1994 | 7.1% | 1.2% | 1.3% | 5.9% | 0.7% | 0.7% | 0.0% | 7.4% | 0.7% | 2.8% | 0.0% | 0.5% | 7.4% | 0.8% | 63.7% |
| 2003 | 8.3% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 1.1% | 0.0% | 0.0% | 0.0% | 6.9% | 13.4% | 6.9% | 63.1% |
| 2004 | 3.5% | 0.0% | 0.0% | 2.0% | 0.0% | 0.0% | 3.0% | 2.4% | 0.0% | 0.0% | 2.0% | 8.1% | 8.7% | 6.7% | 63.6% |
| (88-94) | 4.3% | 1.0% | 0.6% | 5.1% | 0.3% | 0.7% | 0.0% | 13.9% | 0.1% | 1.0% | 0.6% | 7.6% | 17.1% | 3.5% | 44.1% |
| (03-04) | 5.9% | 0.0% | 0.0% | 1.2% | 0.0% | 0.0% | 1.5% | 1.8% | 0.0% | 0.0% | 1.0% | 7.5% | 11.1% | 6.8% | 63.4% |

Appendix E.65. Percent distribution of Lewis River Wild Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1981 | 6.4% | 0.0% | 0.0% | 3.3% | 1.4% | 0.2% | 2.1% | 6.0% | 0.0% | 0.7% | 0.0% | 2.0% | 4.2% | 15.9% | 57.8% |
| 1982 | 6.0% | 1.3% | 0.2% | 3.0% | 1.4% | 0.8% | 0.0% | 10.7% | 0.4% | 0.8% | 0.0% | 4.1% | 6.2% | 23.5% | 41.7% |
| 1986 | 4.9% | 0.0% | 0.0% | 1.6% | 2.2% | 0.9% | 0.0% | 6.8% | 0.0% | 0.0% | 2.5% | 3.3% | 26.6% | 12.3% | 39.0% |
| 1987 | 4.1% | 0.0% | 0.0% | 4.7% | 1.3% | 0.0% | 0.0% | 8.4% | 0.0% | 0.0% | 0.9% | 2.7% | 25.7% | 6.3% | 46.0% |
| 1988 | 4.4% | 0.0% | 0.0% | 2.9% | 0.0% | 0.5% | 0.0% | 8.9% | 0.0% | 0.1% | 0.0% | 4.7% | 23.1% | 16.7% | 38.7% |
| 1989 | 1.8% | 0.2% | 0.2% | 4.5% | 0.2% | 0.7% | 0.5% | 5.1% | 0.0% | 0.8% | 0.5% | 4.9% | 9.5% | 7.3% | 63.9% |
| 1990 | 5.4% | 0.0% | 0.0% | 1.7% | 0.4% | 0.6% | 0.6% | 12.1% | 0.0% | 0.0% | 0.8% | 4.0% | 3.3% | 5.2% | 65.8% |
| 1991 | 6.0% | 0.1% | 0.0% | 3.8% | 0.5% | 0.0% | 1.1% | 5.9% | 0.0% | 0.7% | 0.0% | 2.4% | 15.8% | 7.1% | 56.6% |
| 1992 | 1.6% | 0.0% | 0.0% | 3.8% | 1.8% | 0.0% | 0.7% | 6.2% | 0.0% | 0.0% | 0.0% | 2.9% | 4.5% | 23.4% | 55.1% |
| 1993 | 3.6% | 0.0% | 1.0% | 4.9% | 0.0% | 0.3% | 0.0% | 7.6% | 0.0% | 1.6% | 0.0% | 0.8% | 6.8% | 9.1% | 64.3% |
| 1994 | 6.4% | 0.0% | 0.0% | 3.2% | 0.0% | 0.0% | 0.0% | 3.2% | 0.0% | 1.6% | 0.0% | 0.8% | 1.6% | 0.0% | 83.2% |
| 1995 | 6.6% | 0.0% | 2.3% | 3.2% | 0.0% | 0.4% | 0.0% | 5.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 24.6% | 57.6% |
| 1996 | 7.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.8% | 0.9% | 4.6% | 84.0% |
| 1997 | 12.6% | 0.0% | 0.0% | 3.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.6% | 80.7% |
| 1998 | 8.1% | 0.0% | 0.0% | 3.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 2.0% | 84.8% |
| 1999 | 11.8% | 0.0% | 0.0% | 5.9% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 82.4% |
| 2000 | 3.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.4% | 3.0% | 77.6% |
| 2001 | 5.0% | 0.0% | 1.4% | 0.0% | 0.0% | 0.0% | 0.0% | 8.6% | 0.0% | 0.0% | 2.3% | 5.9% | 2.3% | 5.4% | 69.4% |
| 2002 | 11.2% | 0.0% | 1.6% | 0.0% | 0.0% | 0.0% | 0.0% | 6.0% | 0.0% | 0.0% | 6.3% | 5.2% | 4.9% | 4.6% | 60.1% |
| 2003 | 9.1% | 0.0% | 0.0% | 1.5% | 0.0% | 0.0% | 2.1% | 4.9% | 0.0% | 0.0% | 3.4% | 9.1% | 6.5% | 6.5% | 57.0% |
| 2004 | 6.1% | 0.0% | 0.5% | 2.7% | 0.0% | 0.0% | 1.0% | 2.2% | 0.0% | 0.0% | 0.0% | 0.7% | 2.4% | 1.8% | 82.5% |
| (81-82) | 6.2% | 0.6% | 0.1% | 3.2% | 1.4% | 0.5% | 1.1% | 8.3% | 0.2% | 0.7% | 0.0% | 3.1% | 5.2% | 19.7% | 49.7% |
| (86-98) | 5.6% | 0.0% | 0.3% | 3.1% | 0.5% | 0.3% | 0.2% | 5.3% | 0.0% | 0.4% | 0.4% | 2.2% | 9.2% | 9.4% | 63.1% |
| (99-04) | 7.7% | 0.0% | 0.6% | 1.7% | 0.0% | 0.0% | 0.5% | 3.6% | 0.0% | 0.0% | 2.0% | 3.5% | 5.4% | 3.6% | 71.5% |

Appendix E.66. Percent distribution of Lewis River Wild Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1981 | 7.4% | 0.0% | 0.0% | 3.8% | 1.6% | 0.2% | 2.1% | 7.5% | 0.0% | 0.7% | 0.0% | 2.5% | 4.2% | 16.8% | 53.1% |
| 1982 | 7.4% | 1.2% | 0.2% | 3.5% | 1.6% | 0.7% | 0.0% | 11.7% | 0.4% | 0.7% | 0.0% | 4.2% | 6.0% | 23.5% | 38.8% |
| 1986 | 6.4% | 0.0% | 0.0% | 2.2% | 2.2% | 1.0% | 0.0% | 8.0% | 0.0% | 0.0% | 2.6% | 3.8% | 25.5% | 12.3% | 36.0% |
| 1987 | 5.7% | 0.0% | 0.0% | 5.3% | 1.4% | 0.0% | 0.0% | 9.5% | 0.0% | 0.0% | 0.9% | 2.9% | 24.9% | 6.6% | 42.7% |
| 1988 | 5.2% | 0.0% | 0.0% | 3.5% | 0.0% | 0.5% | 0.0% | 10.7% | 0.0% | 0.1% | 0.0% | 5.0% | 21.9% | 17.7% | 35.4% |
| 1989 | 2.4% | 0.6% | 0.3% | 5.1% | 0.2% | 0.7% | 0.4% | 5.8% | 0.0% | 0.8% | 0.5% | 5.4% | 9.3% | 7.8% | 60.5% |
| 1990 | 7.8% | 0.0% | 0.0% | 1.9% | 0.5% | 0.7% | 0.6% | 13.3% | 0.0% | 0.0% | 0.8% | 4.2% | 3.2% | 5.5% | 61.5% |
| 1991 | 7.0% | 0.3% | 0.0% | 4.1% | 0.4% | 0.0% | 1.2% | 6.4% | 0.0% | 0.7% | 0.0% | 2.5% | 15.4% | 7.7% | 54.2% |
| 1992 | 1.7% | 0.0% | 0.0% | 4.3% | 1.9% | 0.0% | 0.7% | 6.7% | 0.0% | 0.0% | 0.0% | 3.1% | 4.5% | 24.9% | 52.2% |
| 1993 | 4.4% | 0.0% | 1.2% | 5.7% | 0.0% | 0.2% | 0.0% | 8.4% | 0.0% | 1.5% | 0.0% | 1.5% | 6.7% | 9.4% | 61.0% |
| 1994 | 9.4% | 0.0% | 0.0% | 4.9% | 0.0% | 0.0% | 0.0% | 3.8% | 0.0% | 1.5% | 0.0% | 0.8% | 1.5% | 0.0% | 78.2% |
| 1995 | 7.8% | 0.0% | 2.3% | 3.9% | 0.0% | 0.5% | 0.0% | 6.4% | 0.0% | 0.2% | 0.0% | 0.0% | 0.0% | 25.3% | 53.7% |
| 1996 | 9.1% | 0.0% | 0.0% | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.7% | 0.9% | 4.8% | 82.2% |
| 1997 | 14.0% | 0.0% | 0.0% | 3.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 3.9% | 78.9% |
| 1998 | 8.1% | 0.0% | 0.0% | 3.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.0% | 2.0% | 84.8% |
| 1999 | 18.3% | 0.0% | 1.7% | 5.0% | 0.0% | 0.0% | 0.0% | 1.7% | 0.0% | 0.0% | 1.7% | 1.7% | 0.0% | 0.0% | 70.0% |
| 2000 | 6.8% | 0.0% | 1.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.7% | 15.1% | 2.7% | 71.2% |
| 2001 | 5.9% | 0.0% | 1.7% | 0.0% | 0.0% | 0.0% | 0.0% | 8.9% | 0.0% | 0.0% | 3.4% | 6.4% | 2.1% | 6.4% | 65.3% |
| 2002 | 14.3% | 0.0% | 1.8% | 0.0% | 0.0% | 0.0% | 0.0% | 6.0% | 0.0% | 0.0% | 6.8% | 6.5% | 4.8% | 4.8% | 55.1% |
| 2003 | 10.2% | 0.0% | 0.0% | 1.6% | 0.0% | 0.0% | 2.6% | 4.8% | 0.0% | 0.0% | 4.0% | 10.0% | 6.4% | 6.8% | 53.8% |
| 2004 | 6.9% | 0.0% | 0.6% | 2.9% | 0.0% | 0.0% | 1.4% | 2.2% | 0.0% | 0.0% | 0.0% | 0.7% | 2.4% | 1.9% | 81.0% |
| (81-82) | 7.4% | 0.6% | 0.1% | 3.6% | 1.6% | 0.5% | 1.1% | 9.6% | 0.2% | 0.7% | 0.0% | 3.3% | 5.1% | 20.1% | 46.0% |
| (86-98) | 6.9% | 0.1% | 0.3% | 3.6% | 0.5% | 0.3% | 0.2% | 6.1% | 0.0% | 0.4% | 0.4% | 2.4% | 8.9% | 9.8% | 60.1% |
| (99-04) | 10.4% | 0.0% | 1.2% | 1.6% | 0.0% | 0.0% | 0.7% | 3.9% | 0.0% | 0.0% | 2.6% | 4.7% | 5.1% | 3.8% | 66.1% |

Appendix E.67. Percent distribution of Salmon River Chinook reported catch among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1981 | 13.9% | 0.0% | 0.4% | 28.2% | 0.6% | 1.8% | 0.0% | 3.7% | 0.0% | 0.0% | 0.7% | 1.3% | 0.0% | 17.1% | 32.2% |
| 1982 | 10.4% | 1.5% | 0.9% | 14.4% | 1.1% | 0.8% | 0.0% | 7.0% | 0.0% | 0.0% | 0.0% | 2.6% | 0.0% | 21.4% | 39.9% |
| 1983 | 20.6% | 0.6% | 0.0% | 21.5% | 0.6% | 0.0% | 0.0% | 10.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 15.6% | 30.6% |
| 1984 | 10.5% | 0.0% | 0.0% | 16.9% | 3.5% | 0.4% | 0.0% | 3.4% | 0.0% | 0.8% | 0.0% | 0.3% | 0.4% | 21.5% | 42.4% |
| 1985 | 11.9% | 6.5% | 0.0% | 19.1% | 1.1% | 0.3% | 0.0% | 1.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 19.9% | 39.8% |
| 1986 | 15.2% | 0.0% | 0.0% | 9.0% | 4.7% | 0.6% | 0.0% | 2.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.2% | 52.1% |
| 1987 | 10.4% | 0.0% | 0.0% | 15.3% | 0.4% | 0.0% | 0.0% | 2.4% | 0.0% | 0.0% | 0.0% | 2.6% | 0.0% | 24.1% | 44.8% |
| 1988 | 9.6% | 0.0% | 0.0% | 6.4% | 0.6% | 0.0% | 0.0% | 3.9% | 0.0% | 0.0% | 0.0% | 0.8% | 0.0% | 16.0% | 62.7% |
| 1989 | 8.4% | 0.0% | 0.0% | 11.4% | 0.0% | 0.2% | 0.0% | 3.9% | 0.0% | 1.2% | 0.0% | 3.4% | 0.0% | 24.7% | 46.8% |
| 1990 | 11.9% | 0.7% | 0.0% | 10.6% | 0.3% | 0.7% | 1.3% | 7.8% | 0.0% | 0.3% | 0.0% | 3.0% | 0.0% | 25.6% | 37.9% |
| 1991 | 18.4% | 0.0% | 0.5% | 15.2% | 0.1% | 0.7% | 0.8% | 5.8% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 24.9% | 33.4% |
| 1992 | 2.6% | 0.6% | 0.0% | 6.6% | 0.8% | 0.4% | 1.8% | 15.4% | 0.0% | 0.0% | 0.0% | 1.8% | 0.0% | 15.9% | 54.1% |
| 1993 | 7.7% | 0.2% | 0.2% | 15.3% | 0.2% | 0.0% | 1.1% | 17.8% | 0.0% | 0.5% | 0.0% | 3.2% | 0.0% | 23.0% | 30.8% |
| 1994 | 8.8% | 0.2% | 1.0% | 14.8% | 0.2% | 0.1% | 2.1% | 4.6% | 0.0% | 0.0% | 0.0% | 1.5% | 0.0% | 17.7% | 49.0% |
| 1995 | 6.8% | 0.2% | 0.3% | 4.6% | 0.1% | 0.1% | 0.6% | 0.9% | 0.0% | 0.0% | 0.2% | 0.1% | 0.0% | 30.6% | 55.5% |
| 1996 | 11.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.7% | 0.0% | 52.6% | 31.5% |
| 1997 | 27.7% | 0.0% | 1.6% | 3.3% | 0.1% | 0.0% | 0.4% | 0.2% | 0.0% | 0.0% | 0.0% | 1.4% | 0.0% | 19.2% | 46.1% |
| 1998 | 10.5% | 0.4% | 0.4% | 11.1% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.1% | 32.5% | 44.4% |
| 1999 | 12.5% | 0.4% | 0.0% | 2.7% | 0.0% | 0.0% | 2.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 36.2% | 45.6% |
| 2000 | 14.9% | 0.0% | 0.6% | 2.6% | 0.0% | 0.0% | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 8.7% | 72.3% |
| 2001 | 12.5% | 0.0% | 0.7% | 2.7% | 0.0% | 0.0% | 0.3% | 0.3% | 0.0% | 0.0% | 0.2% | 2.5% | 0.1% | 27.2% | 53.5% |
| 2002 | 18.2% | 0.0% | 0.9% | 2.9% | 0.0% | 0.0% | 1.9% | 0.1% | 0.0% | 0.0% | 0.0% | 1.6% | 0.0% | 37.0% | 37.4% |
| 2003 | 12.6% | 0.6% | 0.6% | 5.7% | 0.0% | 0.0% | 4.4% | 0.0% | 0.0% | 0.0% | 0.4% | 1.4% | 0.0% | 34.3% | 40.2% |
| 2004 | 18.2% | 0.8% | 0.9% | 7.2% | 0.0% | 0.0% | 4.1% | 1.2% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 24.0% | 43.0% |
| (81-84) | 13.8% | 0.5% | 0.3% | 20.2% | 1.5% | 0.7% | 0.0% | 6.1% | 0.0% | 0.2% | 0.2% | 1.1% | 0.1% | 18.9% | 36.3% |
| (85-98) | 11.5% | 0.6% | 0.3% | 10.2% | 0.6% | 0.2% | 0.6% | 4.7% | 0.0% | 0.1% | 0.0% | 1.6% | 0.0% | 24.5% | 44.9% |
| (99-04) | 14.8% | 0.3% | 0.6% | 4.0% | 0.0% | 0.0% | 2.2% | 0.3% | 0.0% | 0.0% | 0.1% | 1.2% | 0.0% | 27.9% | 48.7% |

Appendix E.68. Percent distribution of Salmon River Chinook total fishing mortalities among fisheries and escapement.

| Catch Year | Alaska Troll | Alaska Net | Alaska Sport | North Troll | Central Troll | N/CBC Net | N/CBC Sport | WCVI Troll | GeoSt Tr&Sp | Other Fisheries | | | | | Escapement |
|------------|--------------|------------|--------------|-------------|---------------|-----------|-------------|------------|-------------|-----------------|--------------|------------|----------|------------|------------|
| | | | | | | | | | | Canada Net | Canada Sport | U.S. Troll | U.S. Net | U.S. Sport | |
| 1981 | 15.8% | 0.0% | 0.4% | 29.9% | 1.0% | 1.8% | 0.0% | 4.7% | 0.0% | 0.0% | 0.6% | 1.4% | 0.0% | 16.4% | 27.9% |
| 1982 | 14.2% | 1.8% | 0.9% | 17.7% | 1.4% | 0.6% | 0.0% | 7.4% | 0.0% | 0.0% | 0.0% | 2.3% | 0.0% | 20.2% | 33.4% |
| 1983 | 26.3% | 0.7% | 0.0% | 22.1% | 0.7% | 0.0% | 0.0% | 10.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 14.1% | 26.0% |
| 1984 | 11.8% | 0.0% | 0.0% | 17.9% | 3.4% | 0.4% | 0.0% | 3.5% | 0.0% | 0.7% | 0.0% | 0.2% | 0.4% | 22.3% | 39.4% |
| 1985 | 14.5% | 11.8% | 0.0% | 17.7% | 1.1% | 0.2% | 0.0% | 1.6% | 0.0% | 0.0% | 0.0% | 0.1% | 0.0% | 20.3% | 32.5% |
| 1986 | 22.0% | 0.0% | 0.0% | 11.1% | 4.3% | 0.5% | 0.0% | 3.0% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 15.7% | 42.9% |
| 1987 | 17.7% | 0.0% | 0.0% | 15.5% | 0.5% | 0.0% | 0.0% | 2.7% | 0.0% | 0.0% | 0.0% | 2.5% | 0.0% | 22.5% | 38.6% |
| 1988 | 15.0% | 0.0% | 0.0% | 8.7% | 0.9% | 0.0% | 0.0% | 5.3% | 0.0% | 0.0% | 0.0% | 0.9% | 0.0% | 15.5% | 53.6% |
| 1989 | 18.9% | 0.0% | 0.0% | 16.0% | 0.0% | 0.1% | 0.0% | 4.5% | 0.0% | 1.0% | 0.0% | 3.2% | 0.0% | 21.6% | 34.6% |
| 1990 | 18.8% | 2.0% | 0.0% | 12.8% | 0.3% | 0.6% | 1.2% | 7.9% | 0.0% | 0.2% | 0.0% | 2.9% | 0.0% | 23.2% | 30.2% |
| 1991 | 24.1% | 0.0% | 0.5% | 16.4% | 0.1% | 0.7% | 0.8% | 6.1% | 0.0% | 0.0% | 0.0% | 0.2% | 0.0% | 23.1% | 28.0% |
| 1992 | 5.0% | 1.8% | 0.0% | 8.3% | 0.9% | 0.3% | 2.1% | 17.6% | 0.0% | 0.0% | 0.0% | 2.0% | 0.0% | 15.7% | 46.3% |
| 1993 | 11.2% | 0.6% | 0.2% | 17.2% | 0.2% | 0.0% | 1.0% | 18.8% | 0.0% | 0.4% | 0.0% | 3.2% | 0.0% | 22.1% | 25.1% |
| 1994 | 16.3% | 0.4% | 1.0% | 15.0% | 0.2% | 0.1% | 2.1% | 4.7% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 16.8% | 42.2% |
| 1995 | 10.4% | 0.3% | 0.4% | 6.7% | 0.2% | 0.1% | 0.8% | 1.2% | 0.0% | 0.0% | 0.2% | 0.1% | 0.0% | 31.0% | 48.6% |
| 1996 | 20.6% | 0.0% | 0.0% | 2.7% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 3.9% | 0.0% | 47.7% | 24.6% |
| 1997 | 32.2% | 0.0% | 1.7% | 3.4% | 0.1% | 0.0% | 0.4% | 0.2% | 0.0% | 0.0% | 0.0% | 1.5% | 0.0% | 18.9% | 41.6% |
| 1998 | 11.8% | 1.2% | 0.5% | 11.8% | 0.0% | 0.0% | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.1% | 33.0% | 40.8% |
| 1999 | 18.0% | 0.8% | 0.0% | 3.0% | 0.0% | 0.0% | 3.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.6% | 0.0% | 35.4% | 39.3% |
| 2000 | 20.6% | 0.0% | 0.8% | 3.1% | 0.0% | 0.0% | 0.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.4% | 0.0% | 9.2% | 65.1% |
| 2001 | 17.4% | 0.0% | 1.0% | 3.0% | 0.0% | 0.0% | 0.4% | 0.2% | 0.0% | 0.0% | 0.2% | 2.8% | 0.1% | 27.7% | 47.1% |
| 2002 | 22.6% | 0.0% | 1.2% | 3.2% | 0.0% | 0.0% | 2.4% | 0.1% | 0.0% | 0.0% | 0.0% | 1.7% | 0.0% | 36.8% | 32.0% |
| 2003 | 14.5% | 2.2% | 0.6% | 6.3% | 0.0% | 0.0% | 5.2% | 0.0% | 0.0% | 0.0% | 0.4% | 1.5% | 0.0% | 34.1% | 35.2% |
| 2004 | 20.6% | 2.7% | 0.8% | 7.5% | 0.0% | 0.0% | 5.1% | 1.1% | 0.0% | 0.0% | 0.0% | 0.5% | 0.0% | 23.4% | 38.1% |
| (81-84) | 17.0% | 0.6% | 0.3% | 21.9% | 1.6% | 0.7% | 0.0% | 6.4% | 0.0% | 0.2% | 0.2% | 1.0% | 0.1% | 18.3% | 31.7% |
| (85-98) | 17.0% | 1.3% | 0.3% | 11.7% | 0.6% | 0.2% | 0.6% | 5.3% | 0.0% | 0.1% | 0.0% | 1.6% | 0.0% | 23.3% | 37.8% |
| (99-04) | 18.9% | 1.0% | 0.7% | 4.3% | 0.0% | 0.0% | 2.8% | 0.2% | 0.0% | 0.0% | 0.1% | 1.3% | 0.0% | 27.8% | 42.8% |

Appendix F. Time series of abundance indices from 1979 to 2005 for SEAK, NBC, and WCVI AABM fisheries as estimated by CTC Chinook Model calibration CLB0604.

This time series is NOT the first postseason AI and is for trend analysis only (Figures 3.4 to 3.6). For evaluation of overage and underage (Tables 3.4 and 3.5), use the first postseason AI in Table 3.3 instead.

| Year | SEAK | NBC | WCVI |
|------|------|------|------|
| 1979 | 0.97 | 1.04 | 1.10 |
| 1980 | 1.03 | 0.98 | 0.97 |
| 1981 | 0.92 | 0.94 | 0.93 |
| 1982 | 1.08 | 1.05 | 1.01 |
| 1983 | 1.28 | 1.23 | 0.93 |
| 1984 | 1.47 | 1.40 | 1.01 |
| 1985 | 1.35 | 1.33 | 0.99 |
| 1986 | 1.51 | 1.48 | 1.02 |
| 1987 | 1.77 | 1.76 | 1.18 |
| 1988 | 2.17 | 1.87 | 1.13 |
| 1989 | 1.88 | 1.70 | 0.98 |
| 1990 | 1.90 | 1.65 | 0.89 |
| 1991 | 1.81 | 1.53 | 0.75 |
| 1992 | 1.67 | 1.41 | 0.77 |
| 1993 | 1.68 | 1.43 | 0.69 |
| 1994 | 1.58 | 1.26 | 0.52 |
| 1995 | 1.07 | 0.98 | 0.41 |
| 1996 | 0.94 | 0.93 | 0.49 |
| 1997 | 1.24 | 1.12 | 0.58 |
| 1998 | 1.20 | 1.01 | 0.56 |
| 1999 | 1.09 | 0.95 | 0.49 |
| 2000 | 0.97 | 0.93 | 0.49 |
| 2001 | 1.18 | 1.22 | 0.77 |
| 2002 | 1.77 | 1.70 | 1.13 |
| 2003 | 2.28 | 1.93 | 1.19 |
| 2004 | 2.13 | 1.83 | 0.97 |
| 2005 | 1.90 | 1.65 | 0.84 |
| 2006 | 1.69 | 1.53 | 0.75 |

Appendix G. Model estimates of the stock composition of the AABM, and other troll and sport fisheries for 2005 and the average from 1985 to 2004.

“Catch as Percent of Fishery” represents the stock composition of a specific fishery; “Catch as Percent of All Fisheries” represents the proportion of the total catch of a stock that is caught in a specific fishery; “Percent of Total Return” represents the proportion of total return (catch + escapement) caught in a specific fishery.

LIST OF APPENDIX G TABLES

| | <u>Page</u> |
|---|--------------------|
| Appendix G.1. Southeast Alaska All Gear. | 184 |
| Appendix G.2. North B.C. Troll and Sport. | 185 |
| Appendix G.3. Central B.C. Troll. | 186 |
| Appendix G.4. WCVI Troll and Outside Sport. | 187 |
| Appendix G.5. Georgia Strait Sport and Troll. | 188 |
| Appendix G.6. Washington/Oregon Troll and Sport. | 189 |

Appendix G.1. Southeast Alaska All Gear.

| Model Stock | 2005 Catch as Percent of Fishery | Average (1985 - 2004) | | |
|--------------------------------|--|-----------------------------------|---|--|
| | | Catch as Percent of Fishery | Catch as Percent of All Fisheries | Catch as Percent of Total Return |
| WCVI Hatchery | 18.84% | 16.74% | 42.59% | 16.22% |
| Oregon Coastal North Migrating | 17.27% | 16.17% | 34.94% | 15.13% |
| Columbia Upriver Bright | 19.27% | 15.74% | 26.82% | 12.83% |
| North/Central BC | 9.12% | 14.94% | 27.12% | 10.15% |
| Fraser Early | 5.19% | 5.68% | 27.36% | 6.80% |
| Mid-Columbia Brights | 7.03% | 4.95% | 32.68% | 12.30% |
| Upper Georgia Strait | 5.28% | 4.11% | 36.08% | 19.75% |
| Alaska South SE | 3.65% | 3.97% | 96.64% | 35.00% |
| WCVI Wild | 1.33% | 3.61% | 42.63% | 16.36% |
| Washington Coastal Wild | 2.68% | 3.47% | 19.31% | 9.97% |
| WA Coastal Hatchery | 1.85% | 2.60% | 16.18% | 9.34% |
| Columbia Upriver Summer | 4.46% | 2.24% | 34.54% | 13.69% |
| Willamette River Hatchery | 1.06% | 2.09% | 13.26% | 4.60% |
| Fall Cowlitz Hatchery | 0.99% | 1.24% | 6.54% | 2.39% |
| Lewis River Wild | 0.63% | 0.89% | 17.55% | 7.19% |
| Lower GS Hatchery | 0.40% | 0.43% | 3.72% | 1.84% |
| Lower Georgia Strait | 0.08% | 0.26% | 4.35% | 2.10% |
| Fraser Late | 0.13% | 0.23% | 0.50% | 0.15% |
| PS Hatchery Fingerling | 0.13% | 0.15% | 0.47% | 0.25% |
| Skagit Summer/Fall | 0.12% | 0.11% | 4.22% | 1.10% |
| Spring Cowlitz Hatchery | 0.07% | 0.08% | 1.65% | 0.80% |
| Snake River Fall | 0.31% | 0.08% | 8.39% | 5.10% |
| Puget Sound Natural | 0.03% | 0.07% | 0.49% | 0.25% |
| Stillaguamish Summer/Fall | 0.04% | 0.06% | 15.15% | 5.51% |
| Nooksack Fall | 0.01% | 0.04% | 0.14% | 0.10% |
| Snohomish Summer/Fall | 0.03% | 0.04% | 3.40% | 0.90% |
| PS Yearling | 0.01% | 0.02% | 0.47% | 0.32% |
| Lower Bonneville Hatchery | 0.00% | 0.00% | 0.00% | 0.00% |
| Spring Creek Hatchery | 0.00% | 0.00% | 0.00% | 0.00% |
| Nooksack Spring | 0.00% | 0.00% | 0.00% | 0.00% |

Appendix G.2.North B.C. Troll and Sport.

| Model Stock | 2005 Catch as Percent of Fishery | Average (1985 - 2004) | | |
|--------------------------------|--|-----------------------------------|---|--|
| | | Catch as Percent of Fishery | Catch as Percent of All Fisheries | Catch as Percent of Total Return |
| North/Central BC | 45.97% | 44.14% | 61.10% | 24.75% |
| Oregon Coastal North Migrating | 11.85% | 14.51% | 26.05% | 12.36% |
| Columbia Upriver Bright | 8.13% | 7.30% | 10.68% | 5.39% |
| WCVI Hatchery | 5.59% | 6.67% | 12.86% | 5.52% |
| Upper Georgia Strait | 8.17% | 4.25% | 31.28% | 17.56% |
| Fraser Early | 2.85% | 3.36% | 13.89% | 4.13% |
| Washington Coastal Wild | 1.93% | 3.20% | 14.08% | 8.03% |
| Willamette River Hatchery | 1.35% | 3.05% | 14.51% | 5.90% |
| WA Coastal Hatchery | 1.35% | 2.41% | 12.44% | 7.54% |
| Mid-Columbia Brights | 2.91% | 2.09% | 12.56% | 5.12% |
| Columbia Upriver Summer | 4.05% | 1.75% | 22.21% | 9.31% |
| WCVI Wild | 0.40% | 1.48% | 12.69% | 5.49% |
| Lower GS Hatchery | 1.24% | 1.11% | 8.45% | 4.24% |
| Fall Cowlitz Hatchery | 0.92% | 1.03% | 4.33% | 1.71% |
| Fraser Late | 0.65% | 0.82% | 1.42% | 0.48% |
| Lower Georgia Strait | 0.26% | 0.58% | 8.38% | 4.23% |
| Skagit Summer/Fall | 0.58% | 0.41% | 14.34% | 3.80% |
| Nooksack Fall | 0.16% | 0.37% | 1.16% | 0.83% |
| Lewis River Wild | 0.32% | 0.36% | 5.25% | 2.49% |
| PS Hatchery Fingerling | 0.32% | 0.28% | 0.78% | 0.42% |
| Spring Cowlitz Hatchery | 0.22% | 0.24% | 4.00% | 2.12% |
| Snohomish Summer/Fall | 0.21% | 0.19% | 13.60% | 3.79% |
| Puget Sound Natural | 0.06% | 0.11% | 0.73% | 0.38% |
| Alaska South SE | 0.10% | 0.10% | 2.28% | 0.82% |
| PS Yearling | 0.10% | 0.09% | 1.73% | 1.15% |
| Snake River Fall | 0.21% | 0.05% | 5.73% | 3.76% |
| Stillaguamish Summer/Fall | 0.04% | 0.04% | 8.43% | 3.14% |
| Spring Creek Hatchery | 0.04% | 0.02% | 0.06% | 0.04% |
| Nooksack Spring | 0.00% | 0.00% | 1.22% | 0.45% |
| Lower Bonneville Hatchery | 0.00% | 0.00% | 0.00% | 0.00% |

Appendix G.3. Central B.C. Troll.

| Model Stock | 2005 Catch as Percent of Fishery | Average (1985 – 2004) | | |
|--------------------------------|--|-----------------------------------|---|--|
| | | Catch as Percent of Fishery | Catch as Percent of All Fisheries | Catch as Percent of Total Return |
| Fraser Late | 14.29% | 20.66% | 2.18% | 1.16% |
| WCVI Hatchery | 23.81% | 17.87% | 3.21% | 1.39% |
| Columbia Upriver Bright | 14.29% | 8.33% | 0.94% | 0.53% |
| North/Central BC | 4.76% | 6.59% | 1.14% | 0.41% |
| Upper Georgia Strait | 9.52% | 6.05% | 3.63% | 2.23% |
| WCVI Wild | 0.00% | 3.71% | 3.16% | 1.37% |
| Columbia Upriver Summer | 9.52% | 3.69% | 3.62% | 1.70% |
| Fraser Early | 4.76% | 3.55% | 1.06% | 0.35% |
| Washington Coastal Wild | 4.76% | 3.41% | 1.22% | 0.77% |
| Lower GS Hatchery | 4.76% | 2.98% | 1.55% | 1.01% |
| WA Coastal Hatchery | 0.00% | 2.59% | 1.13% | 0.72% |
| Mid-Columbia Brights | 4.76% | 2.55% | 1.11% | 0.54% |
| Oregon Coastal North Migrating | 4.76% | 2.24% | 0.35% | 0.17% |
| Lower Bonneville Hatchery | 0.00% | 1.99% | 0.83% | 0.43% |
| Nooksack Fall | 0.00% | 1.59% | 0.38% | 0.31% |
| Lower Georgia Strait | 0.00% | 1.55% | 1.47% | 1.00% |
| PS Hatchery Fingerling | 0.00% | 1.34% | 0.27% | 0.18% |
| Skagit Summer/Fall | 0.00% | 1.04% | 2.18% | 0.86% |
| Lewis River Wild | 0.00% | 0.66% | 0.69% | 0.36% |
| Puget Sound Natural | 0.00% | 0.60% | 0.27% | 0.18% |
| Snohomish Summer/Fall | 0.00% | 0.50% | 1.70% | 0.87% |
| Spring Creek Hatchery | 0.00% | 0.42% | 0.10% | 0.08% |
| PS Yearling | 0.00% | 0.29% | 0.39% | 0.30% |
| Willamette River Hatchery | 0.00% | 0.28% | 0.09% | 0.05% |
| Spring Cowlitz Hatchery | 0.00% | 0.16% | 0.19% | 0.14% |
| Fall Cowlitz Hatchery | 0.00% | 0.13% | 0.05% | 0.02% |
| Stillaguamish Summer/Fall | 0.00% | 0.12% | 1.77% | 0.86% |
| Snake River Fall | 0.00% | 0.10% | 0.67% | 0.49% |
| Nooksack Spring | 0.00% | 0.01% | 0.27% | 0.15% |
| Alaska South SE | 0.00% | 0.00% | 0.01% | 0.00% |

Appendix G.4. WCVI Troll and Outside Sport.

| Model Stock | 2005 Catch as Percent of Fishery | Average (1985–2004) | | |
|--------------------------------|--|-----------------------------------|---|--|
| | | Catch as Percent of Fishery | Catch as Percent of All Fisheries | Catch as Percent of Total Return |
| Fraser Late | 15.11% | 19.71% | 20.57% | 9.51% |
| Columbia Upriver Bright | 13.68% | 10.38% | 11.13% | 5.76% |
| PS Hatchery Fingerling | 8.93% | 8.35% | 15.08% | 9.29% |
| Fall Cowlitz Hatchery | 7.24% | 7.76% | 25.36% | 11.57% |
| Spring Creek Hatchery | 15.52% | 7.34% | 15.09% | 11.80% |
| Lower Bonneville Hatchery | 1.92% | 6.58% | 30.95% | 14.36% |
| WCVI Hatchery | 4.51% | 6.28% | 9.90% | 4.34% |
| Oregon Coastal North Migrating | 6.86% | 6.23% | 8.76% | 4.05% |
| Nooksack Fall | 1.30% | 4.87% | 10.75% | 8.30% |
| Puget Sound Natural | 1.88% | 3.72% | 15.16% | 9.37% |
| Mid-Columbia Brights | 5.07% | 3.26% | 13.17% | 5.72% |
| Columbia Upriver Summer | 5.50% | 2.39% | 23.43% | 10.44% |
| Washington Coastal Wild | 1.73% | 2.02% | 6.74% | 3.91% |
| Willamette River Hatchery | 1.20% | 1.85% | 6.21% | 2.72% |
| WA Coastal Hatchery | 1.30% | 1.55% | 6.15% | 3.75% |
| WCVI Wild | 0.34% | 1.39% | 9.74% | 4.31% |
| Fraser Early | 1.01% | 1.23% | 3.43% | 1.03% |
| Skagit Summer/Fall | 1.51% | 0.91% | 19.92% | 6.49% |
| Lewis River Wild | 0.84% | 0.82% | 10.53% | 4.96% |
| PS Yearling | 0.70% | 0.74% | 9.51% | 7.07% |
| Spring Cowlitz Hatchery | 0.76% | 0.61% | 7.06% | 4.54% |
| Snohomish Summer/Fall | 0.50% | 0.43% | 17.00% | 6.51% |
| Lower GS Hatchery | 0.50% | 0.41% | 1.99% | 1.14% |
| Snake River Fall | 1.47% | 0.37% | 23.49% | 15.99% |
| North/Central BC | 0.27% | 0.35% | 0.44% | 0.16% |
| Lower Georgia Strait | 0.11% | 0.23% | 1.99% | 1.16% |
| Stillaguamish Summer/Fall | 0.09% | 0.10% | 14.29% | 6.18% |
| Upper Georgia Strait | 0.13% | 0.09% | 0.55% | 0.32% |
| Nooksack Spring | 0.02% | 0.02% | 8.72% | 3.51% |
| Alaska South SE | 0.00% | 0.00% | 0.00% | 0.00% |

Appendix G.5. Georgia Strait Sport and Troll.

| Model Stock | 2005 Catch as Percent of Fishery | Average (1985–2004) | | |
|--------------------------------|--|-----------------------------------|---|--|
| | | Catch as Percent of Fishery | Catch as Percent of All Fisheries | Catch as Percent of Total Return |
| Fraser Late | 44.45% | 49.36% | 44.28% | 20.73% |
| Lower GS Hatchery | 14.82% | 10.61% | 45.95% | 26.66% |
| Nooksack Fall | 3.77% | 9.85% | 19.00% | 14.36% |
| Lower Georgia Strait | 3.13% | 6.11% | 46.92% | 28.20% |
| PS Hatchery Fingerling | 6.65% | 4.73% | 7.54% | 4.57% |
| Fraser Early | 5.08% | 3.94% | 9.34% | 2.60% |
| Upper Georgia Strait | 6.30% | 2.73% | 12.32% | 7.03% |
| Puget Sound Natural | 1.33% | 2.03% | 7.30% | 4.40% |
| PS Yearling | 2.39% | 1.78% | 19.00% | 13.95% |
| Skagit Summer/Fall | 2.13% | 1.17% | 22.38% | 7.21% |
| Columbia Upriver Bright | 1.68% | 1.14% | 1.02% | 0.52% |
| Washington Coastal Wild | 0.89% | 0.94% | 2.76% | 1.56% |
| WCVI Hatchery | 1.10% | 0.87% | 1.27% | 0.46% |
| Spring Creek Hatchery | 2.02% | 0.85% | 1.46% | 1.13% |
| Lower Bonneville Hatchery | 0.29% | 0.80% | 3.21% | 1.34% |
| WA Coastal Hatchery | 0.70% | 0.72% | 2.45% | 1.50% |
| Snohomish Summer/Fall | 0.80% | 0.55% | 19.82% | 7.11% |
| North/Central BC | 0.48% | 0.41% | 0.48% | 0.17% |
| Mid-Columbia Brights | 0.60% | 0.36% | 1.29% | 0.55% |
| Columbia Upriver Summer | 0.72% | 0.29% | 2.67% | 1.13% |
| Nooksack Spring | 0.21% | 0.18% | 57.45% | 24.30% |
| Stillaguamish Summer/Fall | 0.20% | 0.18% | 21.74% | 9.25% |
| WCVI Wild | 0.07% | 0.18% | 1.25% | 0.45% |
| Willamette River Hatchery | 0.13% | 0.12% | 0.36% | 0.15% |
| Spring Cowlitz Hatchery | 0.04% | 0.04% | 0.40% | 0.23% |
| Lewis River Wild | 0.00% | 0.02% | 0.20% | 0.11% |
| Fall Cowlitz Hatchery | 0.00% | 0.02% | 0.04% | 0.02% |
| Snake River Fall | 0.01% | 0.00% | 0.07% | 0.05% |
| Oregon Coastal North Migrating | 0.00% | 0.00% | 0.00% | 0.00% |
| Alaska South SE | 0.00% | 0.00% | 0.00% | 0.00% |

Appendix G.6. Washington/Oregon Troll and Sport.

| Model Stock | 2005 Catch as Percent of Fishery | Average (1985–2004) | | |
|--------------------------------|--|-----------------------------------|---|--|
| | | Catch as Percent of Fishery | Catch as Percent of All Fisheries | Catch as Percent of Total Return |
| Spring Creek Hatchery | 39.07% | 22.99% | 30.03% | 23.70% |
| Fraser Late | 13.33% | 19.79% | 12.96% | 5.43% |
| Fall Cowlitz Hatchery | 18.00% | 19.50% | 39.77% | 16.93% |
| Lower Bonneville Hatchery | 2.88% | 12.39% | 38.11% | 15.96% |
| Columbia Upriver Bright | 5.46% | 3.88% | 2.60% | 1.32% |
| Spring Cowlitz Hatchery | 4.50% | 3.87% | 30.92% | 17.13% |
| PS Hatchery Fingerling | 2.75% | 3.47% | 3.68% | 2.16% |
| Oregon Coastal North Migrating | 3.06% | 2.57% | 2.16% | 0.96% |
| Nooksack Fall | 0.39% | 2.03% | 2.50% | 1.88% |
| Willamette River Hatchery | 1.34% | 1.84% | 3.98% | 1.57% |
| Puget Sound Natural | 0.57% | 1.58% | 3.71% | 2.13% |
| Lewis River Wild | 1.45% | 1.37% | 11.69% | 4.79% |
| Mid-Columbia Brights | 2.00% | 1.21% | 3.05% | 1.26% |
| Washington Coastal Wild | 0.78% | 1.15% | 2.15% | 1.15% |
| WA Coastal Hatchery | 0.61% | 0.90% | 1.93% | 1.12% |
| Columbia Upriver Summer | 1.36% | 0.52% | 3.21% | 1.37% |
| Snake River Fall | 1.88% | 0.51% | 20.73% | 13.63% |
| Fraser Early | 0.28% | 0.17% | 0.35% | 0.09% |
| PS Yearling | 0.11% | 0.12% | 0.93% | 0.66% |
| Alaska South SE | 0.08% | 0.07% | 0.71% | 0.25% |
| Lower GS Hatchery | 0.06% | 0.03% | 0.11% | 0.06% |
| WCVI Hatchery | 0.01% | 0.02% | 0.03% | 0.01% |
| Lower Georgia Strait | 0.01% | 0.01% | 0.12% | 0.06% |
| WCVI Wild | 0.00% | 0.01% | 0.03% | 0.01% |
| Skagit Summer/Fall | 0.01% | 0.00% | 0.03% | 0.01% |
| Snohomish Summer/Fall | 0.00% | 0.00% | 0.03% | 0.01% |
| Upper Georgia Strait | 0.00% | 0.00% | 0.00% | 0.00% |
| Nooksack Spring | 0.00% | 0.00% | 0.00% | 0.00% |
| North/Central BC | 0.00% | 0.00% | 0.00% | 0.00% |
| Stillaguamish Summer/Fall | 0.00% | 0.00% | 0.00% | 0.00% |

Appendix H. Incidental mortality rates applied in the CTC model. Rates in original model were applied to all years. In the current model, rates in some fisheries vary in accordance to changes in management regulations.

| Fishery Number | Fishery | Rates in original Model | | | Rates applied in Model CLB0604 | | | Applicable Years |
|----------------|-----------|-------------------------|------------|---------|--------------------------------|------------|---------|------------------|
| | | Sublegal Rate | Legal Rate | Dropoff | Sublegal Rate | Legal Rate | Dropoff | |
| 1 | Alaska T | 0.3 | 0.3 | 0 | 0.255 | 0.211 | 0.008 | All |
| 2 | North T | 0.3 | 0.3 | 0 | 0.255 | 0.211 | 0.017 | 1979-1995 |
| 2 | North T | | | | 0.220 | 0.185 | 0.016 | 1996-2004 |
| 3 | Centr T | 0.3 | 0.3 | 0 | 0.255 | 0.211 | 0.017 | 1979-1995 |
| 3 | Centr T | | | | 0.220 | 0.185 | 0.016 | 1996-2004 |
| 4 | WCVI T | 0.3 | 0.3 | 0 | 0.255 | 0.211 | 0.017 | 1979-1997 |
| 4 | WCVI T | | | | 0.220 | 0.185 | 0.016 | 1998-2004 |
| 5 | WA/OR T | 0.3 | 0.3 | 0 | 0.255 | 0.211 | 0.017 | 1979-1983 |
| 5 | WA/OR T | | | | 0.220 | 0.185 | 0.016 | 1984-2004 |
| 6 | Geo St T | 0.3 | 0.3 | 0 | 0.255 | 0.211 | 0.017 | 1979-1985,1987 |
| 6 | Geo St T | | | | 0.220 | 0.185 | 0.016 | 1986,1988-2004 |
| 7 | Alaska N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 8 | North N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 9 | Centr N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 10 | WCVI N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 11 | J De F N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 12 | PgtNth N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 13 | PgtSth N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 14 | WashCst N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 15 | Col R N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 16 | JohnSt N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 17 | Fraser N | 0.9 | 0.9 | 0 | 0.9 | 0.9 | 0 | All |
| 18 | Alaska S | 0.3 | 0.3 | 0 | 0.123 | 0.123 | 0.036 | All |
| 19 | Nor/Cen S | 0.3 | 0.3 | 0 | 0.123 | 0.123 | 0.036 | All |
| 20 | WCVI S | 0.3 | 0.3 | 0 | 0.123 | 0.123 | 0.069 | All |
| 21 | WashOcn S | 0.3 | 0.3 | 0 | 0.123 | 0.123 | 0.069 | All |
| 22 | PgtNth S | 0.3 | 0.3 | 0 | 0.123 | 0.123 | 0.145 | All |
| 23 | PgtSth S | 0.3 | 0.3 | 0 | 0.123 | 0.123 | 0.145 | All |
| 24 | Geo St S | 0.3 | 0.3 | 0 | 0.322 | 0.322 | 0.069 | 1979-1981 |
| 24 | Geo St S | | | | 0.123 | 0.123 | 0.069 | 1982-2004 |
| 25 | Col R S | 0.3 | 0.3 | 0 | 0.123 | 0.123 | 0.069 | All |

Appendix I. Components of the February 2005 agreement on harvest sharing for Stikine and Taku Rivers Chinook.

The allowable catch (AC) and Base Terminal run (BTR) for both rivers are calculated as follows:

AC = Terminal run – Base Terminal run (BTR), where:

Terminal run = total Chinook run size of each river minus the US troll catch of Chinook salmon from each river outside Districts 108 and 111.

BTR = escapement target + test fishery Base Level Catch (BLC)

BLCs for the Stikine River include the following:

- U.S. Stikine BLC: 3,400 large Chinook¹
- Canadian Stikine BLC: 2,300 large Chinook²
- Test fishery: 1,400 large Chinook;

¹ Includes average combined US gillnet, troll and sport catches of Stikine Chinook salmon in District 108.

² Includes average combined Canadian Aboriginal, commercial and sport catches of Stikine Chinook salmon.

BLCs for the Taku River include the following:

- U.S. Taku BLC: 3,500 large Chinook³
- Canadian Taku BLC: 1,500 large Chinook⁴
- Test fishery: 1,400 large Chinook;

³ Includes average combined US gillnet and sport catches of Stikine Chinook salmon in District 111. No troll fisheries occurred in District 111 from 1985 – 2003..

⁴ Includes average combined Canadian Aboriginal, commercial and sport catches of Taku Chinook salmon.

Harvest sharing and accounting of the Stikine River AC shall be as follows:

| Allowable Catch Range | | Allowable Catch Share | | | |
|-----------------------|---------|-----------------------|--------|--------|--------|
| | | U.S. | | Canada | |
| Lower | Upper | Lower | Upper | Lower | Upper |
| 0 | 5,000 | 0 | 500 | 0 | 4,500 |
| 5,001 | 20,000 | 501 | 11,000 | 4,500 | 9,000 |
| 20,001 | 30,000 | 11,001 | 17,500 | 9,000 | 12,500 |
| 30,001 | 50,000 | 17,501 | 30,500 | 12,500 | 19,500 |
| 50,001 | 100,000 | 30,501 | 63,000 | 19,500 | 37,000 |

Harvest sharing and accounting of the Taku River AC shall be as follows:

| Allowable Catch Range | | Allowable Catch Share | | | |
|-----------------------|---------|-----------------------|--------|--------|--------|
| | | U.S. | | Canada | |
| Lower | Upper | Lower | Upper | Lower | Upper |
| 0 | 5,000 | 0 | 0 | 0 | 5,000 |
| 5,001 | 20,000 | 1 | 11,000 | 5,000 | 9,000 |
| 20,001 | 30,000 | 11,001 | 17,500 | 9,000 | 12,500 |
| 30,001 | 50,000 | 17,501 | 30,500 | 12,500 | 19,500 |
| 50,001 | 100,000 | 30,501 | 63,000 | 19,500 | 37,000 |

Appendix J. Evaluation of mark-selective fisheries.

Double index tags (DIT) were intended to be used to evaluate mark selective fisheries and seven Puget Sound fall Chinook stocks have DIT groups. The DIT is used as a monitoring tool to test the hypothesis that there are differences between the marked and unmarked tagged groups that would be due to MSFs and also for estimation of mortalities of unmarked fish in MSFs.

MONITORING FUNCTION OF DIT

DIT stocks provide an opportunity to evaluate whether there are any differences between the marked and unmarked tagged pair in the return rate to escapement. Whether there has been a significant change in the ratio of unmarked to marked fish from release to return can be tested comparing the return rates of the unmarked and marked group. A z-test was used to test the null hypothesis of no difference between return rates for marked and unmarked DIT groups (Joint Coho DIT Analysis Workgroup, 2003). The significance level was set at $\alpha = 0.05$, so a test with a P-value less than 0.05 is indicated as significant in Table 3.8. The *P*-values did not account for multiple comparisons and therefore should be interpreted with caution, as in multiple comparisons some tests are expected to be significant due to random chance. Out of 52 tests for brood-ages with marked and unmarked returns to the hatchery, only 6 were significant (Table J.1), and the actual differences were small.

Table J.1. Marked and unmarked returns to hatchery escapement for Puget Sound DIT stocks, % of release returning to hatchery and test of hypothesis of no difference between unmarked and marked % return.

| Stock | Brood | Age | Marked | | Unmarked | | % of Release | | Ho: $p_u = p_m$ | |
|-------|-------|-----|-------------|----------|-------------|----------|--------------|----------|-----------------|-------|
| | | | Est. Tagged | Variance | Est. Tagged | Variance | Marked | Unmarked | Z | Sign. |
| GAD | 1998 | 5 | 15.0 | - | 7.0 | - | 0.01% | 0.00% | (1.71) | ns |
| | 1999 | 4 | 219.0 | - | 264.0 | - | 0.11% | 0.12% | 1.52 | ns |
| | | 5 | 13.0 | - | 12.0 | - | 0.01% | 0.01% | (0.32) | ns |
| | 2000 | 3 | 188.0 | - | 181.3 | 0.3 | 0.08% | 0.08% | (0.44) | ns |
| | | 4 | 235.0 | - | 260.6 | 4.1 | 0.11% | 0.12% | 1.05 | ns |
| | 2001 | 2 | 33.0 | - | 35.3 | 0.5 | 0.02% | 0.02% | 0.55 | ns |
| | | 3 | 285.0 | - | 265.0 | - | 0.13% | 0.13% | (0.10) | ns |
| | 2002 | 2 | 172.8 | 0.8 | 193.8 | 0.8 | 0.08% | 0.09% | 1.14 | ns |
| GRO | 1999 | 4 | 471.0 | - | 513.2 | 9.9 | 0.26% | 0.28% | 1.40 | ns |
| | | 5 | 52.0 | - | 52.0 | - | 0.03% | 0.03% | 0.02 | ns |
| | 2000 | 3 | 272.4 | 3.3 | 280.0 | - | 0.13% | 0.14% | 0.16 | ns |
| | | 4 | 298.0 | 0.0 | 333.0 | 0.0 | 0.15% | 0.16% | 1.22 | ns |
| | 2001 | 2 | 13.0 | - | 12.0 | - | 0.01% | 0.01% | (0.20) | ns |
| | | 3 | 165.2 | 0.2 | 156.1 | 0.1 | 0.08% | 0.08% | (0.52) | ns |
| | 2002 | 2 | 69.8 | 1.9 | 93.4 | 2.5 | 0.04% | 0.05% | 2.00 | * |
| GRN | 1998 | 5 | 15.0 | - | 7.0 | - | 0.01% | 0.00% | (1.79) | ns |
| | 1999 | 4 | 219.0 | - | 264.0 | - | 0.11% | 0.13% | 1.80 | ns |
| | | 5 | 13.0 | - | 12.0 | - | 0.01% | 0.01% | (0.26) | ns |
| | 2000 | 3 | 188.0 | - | 181.3 | 0.3 | 0.10% | 0.09% | (0.76) | ns |
| | | 4 | 235.0 | - | 260.6 | 4.1 | 0.12% | 0.13% | 0.68 | ns |
| | 2001 | 2 | 33.0 | - | 35.3 | 0.5 | 0.02% | 0.02% | 0.67 | ns |
| | | 3 | 285.0 | - | 265.0 | - | 0.16% | 0.16% | 0.25 | ns |
| | 2002 | 2 | 172.8 | 0.8 | 193.8 | 0.8 | 0.09% | 0.10% | 0.81 | ns |
| NIS | 1998 | 5 | 11.2 | 0.2 | 8.1 | 0.1 | 0.01% | 0.00% | (0.59) | ns |

| Stock | Brood | Age | Marked | | Unmarked | | % of Release | | Ho: $p_u=p_m$ | |
|-------|-------|-----|-------------|----------|-------------|----------|--------------|----------|---------------|-------|
| | | | Est. Tagged | Variance | Est. Tagged | Variance | Marked | Unmarked | Z | Sign. |
| | 1999 | 4 | 202.2 | 2.2 | 205.2 | 2.2 | 0.10% | 0.11% | 0.36 | ns |
| | | 5 | 14.0 | - | 8.0 | - | 0.01% | 0.00% | (1.23) | ns |
| | 2000 | 3 | 196.8 | 1.9 | 204.9 | 1.9 | 0.12% | 0.12% | 0.02 | ns |
| | | 4 | 259.1 | 0.1 | 249.0 | - | 0.15% | 0.14% | (0.88) | ns |
| | 2001 | 2 | 65.1 | 0.1 | 68.0 | - | 0.03% | 0.03% | 0.27 | ns |
| | | 3 | 149.0 | - | 177.0 | - | 0.07% | 0.08% | 1.57 | ns |
| | 2002 | 2 | 302.1 | 0.1 | 414.1 | 0.1 | 0.17% | 0.22% | 3.34 | * |
| NSF | 1998 | 5 | 42.1 | 119.2 | 49.8 | 140.9 | 0.03% | 0.03% | 0.76 | ns |
| | 1999 | 4 | 252.8 | 715.4 | 169.6 | 477.0 | 0.13% | 0.08% | (4.26) | * |
| | | 5 | 15.0 | - | 12.0 | - | 0.01% | 0.01% | (0.63) | ns |
| | 2000 | 3 | 118.8 | 326.5 | 108.2 | 303.5 | 0.06% | 0.05% | (0.78) | ns |
| | | 4 | 76.0 | - | 73.2 | 0.2 | 0.04% | 0.04% | (0.30) | ns |
| | 2001 | 2 | 30.3 | 23.3 | 39.1 | 27.4 | 0.02% | 0.02% | 1.02 | ns |
| | | 3 | 179.0 | - | 169.0 | - | 0.09% | 0.09% | (0.61) | ns |
| SAM | 2002 | 2 | 7.4 | 6.3 | 11.6 | 26.6 | 0.00% | 0.01% | 0.94 | ns |
| | 1998 | 5 | 31.0 | 65.1 | 27.9 | 58.6 | 0.02% | 0.01% | (0.45) | ns |
| | 1999 | 4 | 173.6 | 364.6 | 176.7 | 371.1 | 0.10% | 0.10% | (0.06) | ns |
| | | 5 | 7.4 | 6.3 | 5.6 | 4.7 | 0.00% | 0.00% | (0.56) | ns |
| | 2000 | 3 | 12.4 | 26.0 | 9.3 | 19.5 | 0.01% | 0.01% | (0.71) | ns |
| | | 4 | 88.8 | 75.5 | 55.5 | 47.2 | 0.06% | 0.04% | (2.89) | * |
| | 2001 | 3 | 57.4 | 48.7 | 81.4 | 69.2 | 0.03% | 0.04% | 1.98 | * |
| SKS | 2002 | 2 | 7.4 | 6.3 | 11.6 | 26.6 | 0.00% | 0.01% | 1.02 | ns |
| | 1998 | 5 | 2.0 | - | 2.0 | - | 0.00% | 0.00% | (0.02) | ns |
| | 1999 | 4 | 398.0 | - | 414.0 | - | 0.56% | 0.57% | 0.29 | ns |
| | | 5 | 336.0 | - | 342.2 | 0.2 | 0.47% | 0.47% | (0.01) | ns |
| | 2000 | 3 | 105.0 | - | 150.0 | - | 0.14% | 0.20% | 2.90 | * |
| | | 4 | 575.0 | - | 528.0 | - | 0.78% | 0.72% | (1.25) | ns |
| | 2001 | 3 | 214.2 | 0.2 | 206.2 | 0.2 | 0.28% | 0.28% | (0.02) | ns |

These results indicate that the mark-selective fishery to which these stocks are vulnerable (the Washington JDF sport fishery in Areas 5 and 6) does not have sufficient impact that a difference can be detected given the number of tagged fish returning to the hatcheries.

ESTIMATION OF EXPLOITATION RATES FOR UNMARKED FISH.

The estimation of mortalities of unmarked fish in MSFs and exploitation rates was carried out using different approaches for the stocks with DIT and those that do not have DIT, or single index tag group (SIT).

Methods adopted for DIT stocks.

The paired ratio method (SFEC 2002) was used to estimate the mortalities of unmarked tagged fish due to mark-selective release.

$$L_{u,f,a} = L_{m,f,a} \lambda_{f,a} p_{f,a} \quad (1)$$

where,

| | |
|-----------------|---|
| $L_{x,f,a}$ | = landed mortality of fish in mark selective fishery f at age a for group x , where x is marked (m) or unmarked (u) tagged fish |
| $\lambda_{f,a}$ | = unmarked to marked ratio of DIT group in mark selective fishery |
| $p_{f,a}$ | = release mortality in fishery f and at age a . |

The ratio of unmarked to marked fish at release was used to estimate the encounters of unmarked fish from landed marked and tagged fish for the DIT stocks.

The incidental mortalities among the unmarked fish can be related to the marked landed catch using $\lambda_{f,a}$ because the encounter and release mortality rates are assumed to be the same for marked and unmarked fish. For example, unmarked shakers would be

$$shaker_{u,f,a} = \lambda_f (L_{m,f,a} \times EncRate_{f,a} \times ShakerMortRate_{f,a}) \quad (2)$$

The exploitation rate was calculated for unmarked fish from the beginning cohort size

$$ER_u = \sum_{f=1}^n C_{u,f} / N_{u,1} \quad (3)$$

where $C_{u,f}$ is the landed plus incidental mortality for unmarked tagged fish in fishery f and $N_{u,1}$ is the beginning cohort size for unmarked fish.

Methods adopted for SIT stocks

A change-in-ratio model was used to estimate fishing induced mortality and exploitation rates for unmarked Chinook in mark-selective fisheries, where a DIT group was not available. This method, also called a proportional migration model, can be summarized as follows. A time series of $N_{m,f}$ was established, that represents the abundance of marked Chinook available to fishery f , where f represents one in a series of n fisheries. This time series, obtained from COHSHK, also accounts for removals due to escapement and natural mortality. Assuming that the youngest Chinook is age 2 and the oldest age is age 5, the abundance, $N_{m,n,A}$, of age A marked Chinook available to fishery n would be

$$N_{m,n,A} = N_{m,0,2} - \sum_{a=2}^A (M_{0,a} \times N_{m,0,a}) - \sum_{a=2}^A \sum_{f=1}^{n-1} C_{m,f,a} - \sum_{a=2}^A E_{m,n,a} \quad (4)$$

where $N_{m,0,2}$ = initial abundance of marked Chinook at age 2 at the start of the year before any fisheries,

| | |
|---------------|---|
| $M_{0,a}$ | = the natural mortality rate at age a , |
| $N_{m,0,a}$ | = ocean abundance of age a marked Chinook at the start of the year prior to any fisheries, |
| $C_{m,f,a}$ | = mortality (landed and incidental) of marked fish (m) of age a in fishery f |
| $E_{m,n+1,a}$ | = spawning escapement of age a marked Chinook at the end of the year after all fisheries have been completed, and |
| n | = time step and number of fisheries prosecuted. |

The initial lambda is assumed to be equal to 1. After each fishery, escapement, and natural mortality event, the ratio between the unmarked and marked abundance was recalculated from the age a specific abundances

$$\lambda_{n,a} = \frac{N_{u,n-1,a}}{N_{m,n-1,a}} \quad (5)$$

to estimate a parallel time series of unmarked abundance.

$$N_{u,n,A} = N_{u,n-1,a} - \sum_{a=2}^A (M_{0,a} \times \lambda_{0,a} N_{m,0,a}) - \sum_{a=2}^A \sum_{f=1}^{n-1} \lambda_{f,a} C_{m,f,a} r_{f,a} - \sum_{a=2}^A \lambda_{n,a} E_{m,n,a} \quad (6)$$

The mortality of unmarked fish due to mark-selective release was calculated from the landed mortality of marked fish, $L_{m,f,a}$,

$$L_{u,f,a} = L_{m,f,a} \lambda_{f,a} r_{f,a} p_{f,a} \quad (7)$$

where,

$r_{f,a}$ = represent the proportion of the fishery that is selective,

u = unmarked

m = marked

For most stocks, only the Strait of Juan de Fuca component of the North Puget Sound sport fishery was selective, i.e. the July and August harvest in Areas 5 and 6 in the North Puget Sound sport fishery. For the Willamette springs stock, the tangle net component of the Columbia River net and the Columbia River sport fisheries were also selective.

$$P_{NPugetSoundSport,a} = \frac{Area5July \& Aug_a + Area6July \& Aug_a}{Area5 \& 6 \& 7_a} \quad (7)$$

$$P_{ColRnet,a} = \frac{TangleNet_a}{TangleNet_a + otherNet_a} \quad (9)$$

$$P_{ColRSport,a} = \frac{Willamette_a + Clackamas_a + MainstemSport_a + tributary_a}{Willamette_a + Clackamas_a + MainstemSport_a + tributary_a + JohnDay_a + SherarsFalls_a} \quad (10)$$

The incidental mortalities among the unmarked fish and the exploitation rate for unmarked fish were calculated from equations (2) and (3) and are shown in Table J.2.

Example of estimated release mortalities using brood year 2000 age-3 CHI in Table J.2.

- number of CWT recoveries in NPS sport = 7

- proportion of NPS sport that was selective for brood year 2000 age 3 CHI = 0.323741 (area 5 & 6, Jul & Aug only).
- CWt recoveries from area 5 & 6, Jul & Aug only = 2.27 (7×0.323741)
- release mortality rate = 0.14
- release mortalities in selective fishery = 0.32 (2.27×0.14 , this is the release mortalities from area 5-6, Jul-Aug only)
- non-selective landings = 4.73 ($7 - 2.27$, this is the landed catch outside of Jul-Aug area 5-6 + all of the catch from area 7)
- total impacts = 5.1 ($0.32 + 4.73 = 5.05$, which rounds up to 5.1)
- ER = total impacts, selective + non-selective / cohort. Thus, $(0.32 \text{ release mort} + 4.73 \text{ landed} + 1.8 \text{ IM})/6190.4 = 0.0011$ for the unmarked group and $(7 \text{ landed} + 1.8 \text{ IM})/6190 = 0.0014$ for the marked group.

Table J.2. Estimation of MSF mortalities (landed and due to MSF release) and incidental mortalities (IM) for marked and unmarked stocks. The cohort size is after natural mortality.

| Stock | Brood | Age | Marked cohort for age | CWTs landed | IM | ER in MSF | Unmarked cohort for age | MSF Mortality | IM | ER in MSF |
|-----------------------------------|-------|-----|-----------------------|-------------|------|-----------|-------------------------|---------------|------|-----------|
| Area 5 and 6 of North Puget Sound | | | | | | | | | | |
| CHI | 2000 | 3 | 6,190.4 | 7.0 | 1.8 | 0.14% | 6,190.4 | 5.1 | 1.8 | 0.11% |
| COW | 2000 | 3 | 282.4 | 7.0 | 2.0 | 3.19% | 282.4 | 5.4 | 2.0 | 2.62% |
| GAD | 2000 | 3 | 1,205.3 | 15.0 | 4.0 | 1.58% | 901.7 | 1.9 | 4.0 | 0.65% |
| | 2000 | 4 | 712.3 | 5.0 | 1.0 | 0.84% | 488.5 | 2.6 | 2.0 | 0.95% |
| | 2001 | 3 | 1,688.3 | 35.0 | 10.0 | 2.67% | 1,207.8 | 4.0 | 9.4 | 1.11% |
| GRN | 1999 | 4 | 661.1 | 18.5 | 5.9 | 3.70% | 595.9 | 2.3 | 4.6 | 1.16% |
| | 2000 | 3 | 812.3 | 14.0 | 4.0 | 2.22% | 727.9 | 1.8 | 2.3 | 0.56% |
| | 2000 | 4 | 510.4 | 10.0 | 3.0 | 2.55% | 492.3 | 1.3 | 1.7 | 0.60% |
| | 2002 | 2 | 3,102.5 | 4.0 | 21.0 | 0.81% | 2,045.5 | 1.3 | 8.3 | 0.47% |
| GRO | 1999 | 4 | 1,029.3 | - | 6.0 | 0.58% | 10.1 | - | - | 0.00% |
| | 2000 | 3 | 1,791.2 | 27.0 | 7.0 | 1.90% | 1,233.0 | 10.3 | 16.0 | 2.14% |
| | 2000 | 4 | 848.8 | 20.9 | 5.2 | 3.07% | 576.2 | 2.6 | 6.0 | 1.49% |
| | 2001 | 3 | 750.6 | 9.0 | 3.0 | 1.60% | 639.5 | 1.1 | 2.9 | 0.63% |
| | 2002 | 2 | 1,866.4 | 5.0 | 19.0 | 1.29% | 2,409.5 | 1.5 | 17.6 | 0.79% |
| HOK | 1999 | 5 | 769.8 | 5.0 | 1.3 | 0.82% | 769.8 | 0.7 | 1.3 | 0.26% |
| LRH | 2000 | 3 | 2,923.0 | 16.0 | 4.1 | 0.69% | 2,923.0 | 2.2 | 4.1 | 0.22% |
| | 2001 | 3 | 428.1 | 2.0 | 0.6 | 0.61% | 428.1 | 0.3 | 0.6 | 0.21% |
| NIS | 1999 | 4 | 977.9 | 8.0 | 2.0 | 1.02% | 748.9 | 3.2 | 2.8 | 0.80% |
| | 2000 | 3 | 2,032.3 | 24.5 | 6.3 | 1.52% | 1,874.7 | 3.1 | 6.5 | 0.51% |
| | 2000 | 4 | 994.9 | 15.0 | 4.0 | 1.91% | 956.6 | 2.9 | 5.4 | 0.87% |
| NSF | 1999 | 4 | 533.1 | 2.0 | 0.6 | 0.49% | 301.2 | 0.3 | 0.6 | 0.28% |
| | 2000 | 4 | 164.7 | 6.0 | 2.0 | 4.86% | 206.3 | 0.7 | 2.0 | 1.34% |
| | 2001 | 3 | 1,034.4 | 4.0 | 1.0 | 0.48% | 847.0 | 4.5 | 6.0 | 1.24% |
| SAM | 1999 | 4 | 678.1 | 3.9 | 0.8 | 0.70% | 464.4 | 6.7 | 2.3 | 1.94% |
| | | 5 | 36.3 | 4.0 | 1.0 | 13.78% | 28.8 | 3.5 | 2.0 | 19.20% |
| | 2002 | 2 | 842.7 | 4.0 | 12.0 | 1.90% | 867.9 | 1.3 | 12.1 | 1.55% |
| SKF | 1999 | 4 | 929.6 | 8.0 | 2.0 | 1.08% | 929.6 | 5.7 | 2.0 | 0.83% |
| | 2001 | 3 | 788.7 | 4.0 | 1.1 | 0.65% | 788.7 | 2.5 | 1.1 | 0.46% |
| SKS | 1999 | 4 | 1,384.5 | 7.9 | 1.9 | 0.71% | 1,215.3 | 1.0 | 1.9 | 0.24% |
| | | 5 | 429.2 | 3.8 | 1.0 | 1.12% | 410.7 | 0.5 | 1.0 | 0.37% |
| | 2000 | 4 | 1,157.2 | 12.1 | 3.6 | 1.36% | 902.9 | 9.0 | 11.4 | 2.25% |
| SOO | 1999 | 4 | 655.9 | 5.0 | 1.2 | 0.95% | 655.9 | 0.7 | 1.2 | 0.29% |
| | 2001 | 3 | 954.9 | 5.0 | 1.4 | 0.67% | 954.9 | 0.7 | 1.4 | 0.22% |
| SPR | 1999 | 4 | 2,424.3 | 3.0 | 0.7 | 0.15% | 2,424.3 | 0.4 | 0.7 | 0.05% |
| | 2000 | 3 | 6,194.0 | 5.0 | 1.3 | 0.10% | 6,194.0 | 0.7 | 1.3 | 0.03% |
| | 2001 | 3 | 4,996.1 | 17.0 | 4.9 | 0.44% | 4,996.1 | 5.0 | 4.9 | 0.20% |
| SSF | 1998 | 5 | 416.3 | 3.0 | 0.7 | 0.89% | 416.3 | 0.4 | 0.7 | 0.27% |
| | 2001 | 3 | 951.1 | 5.0 | 1.4 | 0.67% | 951.1 | 0.7 | 1.4 | 0.22% |
| STL | 2002 | 2 | 3,011.4 | 2.0 | 1.4 | 0.11% | 3,011.4 | 0.3 | 1.4 | 0.06% |
| SUM | 2000 | 3 | 8,518.0 | 5.0 | 1.3 | 0.07% | 8,518.0 | 0.7 | 1.3 | 0.02% |
| URB | 2000 | 3 | 1,360.1 | 5.0 | 1.3 | 0.46% | 1,360.1 | 0.7 | 1.3 | 0.15% |
| Columbia River net | | | | | | | | | | |
| WSH | 1997 | 3 | 449.1 | - | 3.6 | 0.80% | 449.1 | - | 3.6 | 0.80% |

| Stock | Brood | Age | Marked cohort for age | CWTs landed | IM | ER in MSF | Unmarked cohort for age | MSF Mortality | IM | ER in MSF |
|----------------------|-------|-----|-----------------------|-------------|---------|-----------|-------------------------|---------------|---------|-----------|
| WSH | 1997 | 4 | 27,450.7 | 358.0 | 172.3 | 1.93% | 27,450.7 | 358.0 | 172.3 | 1.93% |
| WSH | 1997 | 5 | 12,574.1 | 2,856.0 | 14.3 | 22.83% | 12,574.1 | 523.4 | 14.3 | 4.28% |
| WSH | 1997 | 6 | 78.2 | 7.0 | - | 8.95% | 78.2 | 1.0 | - | 1.25% |
| WSH | 1998 | 3 | 178.9 | - | 0.5 | 0.28% | 178.9 | - | 0.5 | 0.28% |
| WSH | 1998 | 4 | 6,288.0 | 205.0 | 12.4 | 3.46% | 6,288.0 | 72.8 | 12.4 | 1.35% |
| WSH | 1998 | 5 | 3,982.7 | 49.0 | 1.4 | 1.27% | 3,982.7 | 24.6 | 1.4 | 0.65% |
| WSH | 1998 | 6 | 63.4 | 10.0 | - | 15.77% | 63.4 | 1.4 | - | 2.21% |
| WSH | 1999 | 3 | 134.2 | - | 0.5 | 0.37% | 134.2 | - | 0.5 | 0.37% |
| WSH | 1999 | 4 | 2,113.5 | 44.0 | 11.0 | 2.60% | 2,113.5 | 40.5 | 11.0 | 2.44% |
| WSH | 1999 | 5 | 2,059.4 | 246.0 | 2.0 | 12.04% | 2,059.4 | 50.0 | 2.0 | 2.53% |
| WSH | 2000 | 3 | 171.2 | - | 0.3 | 0.18% | 171.2 | - | 0.3 | 0.18% |
| WSH | 2000 | 4 | 3,487.1 | 160.0 | 6.4 | 4.77% | 3,487.1 | 31.2 | 6.4 | 1.08% |
| WSH | 2001 | 3 | 64.4 | - | - | 0.00% | 64.4 | - | - | 0.00% |
| Columbia River Sport | | | | | | | | | | |
| WSH | 1997 | 3 | 445.5 | 103.0 | 27.5 | 29.29% | 445.5 | 92.6 | 27.5 | 26.97% |
| WSH | 1997 | 4 | 26,920.4 | 5,361.0 | 1,468.4 | 25.37% | 26,920.4 | 750.5 | 1,468.4 | 8.24% |
| WSH | 1997 | 5 | 9,703.8 | 2,741.0 | 253.8 | 30.86% | 12,036.4 | 476.0 | 314.8 | 6.57% |
| WSH | 1997 | 6 | 71.2 | - | 0.2 | 0.28% | 77.2 | - | 0.2 | 0.28% |
| WSH | 1998 | 3 | 178.4 | - | 6.4 | 3.59% | 178.4 | - | 6.4 | 3.59% |
| WSH | 1998 | 4 | 6,070.6 | 1,121.0 | 276.6 | 23.02% | 6,202.8 | 160.4 | 282.6 | 7.14% |
| WSH | 1998 | 5 | 3,932.3 | 653.0 | 61.3 | 18.16% | 3,956.7 | 92.0 | 61.7 | 3.88% |
| WSH | 1998 | 6 | 53.4 | 19.0 | 1.4 | 38.20% | 62.0 | 3.1 | 1.6 | 7.60% |
| WSH | 1999 | 3 | 133.7 | 29.0 | 4.7 | 25.21% | 133.7 | 4.1 | 4.7 | 6.55% |
| WSH | 1999 | 4 | 2,058.5 | 344.0 | 87.5 | 20.96% | 2,062.0 | 48.2 | 87.6 | 6.59% |
| WSH | 1999 | 5 | 1,811.4 | 396.0 | 35.4 | 23.82% | 2,007.4 | 61.4 | 39.2 | 5.01% |
| WSH | 2000 | 3 | 170.9 | 57.0 | 6.9 | 37.39% | 170.9 | 8.0 | 6.9 | 8.71% |
| WSH | 2000 | 4 | 3,320.7 | 735.0 | 120.7 | 25.77% | 3,449.5 | 106.9 | 125.4 | 6.73% |
| WSH | 2001 | 3 | 64.4 | 6.0 | 0.4 | 9.94% | 64.4 | 0.8 | 0.4 | 1.93% |