

**PACIFIC SALMON COMMISSION
JOINT CHUM TECHNICAL COMMITTEE
2015 POST SEASON SUMMARY REPORT
REPORT TCCHUM (18) – 02**

May 2018

MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2015

Canadian Members

Mr. Pieter Van Will, Co-Chair, CDFO
Mr. John Candy, CDFO
Ms. Kim Charlie, Sts'ailes
Ms. Marla Maxwell, CDFO
Mr. Joe Tadey, CDFO
Ms. Louise de Mestral Bezanson, CDFO

United States Members

Mr. Jay Zischke, Co-Chair, Suquamish Indian Tribe
Mr. Scott Bass, PNPTC
Mr. Bill Patton, NWIFC
Dr. Maureen Small, WDFW
Dr. Gary Winans, NOAA

MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2016

Canadian Members

Mr. Pieter Van Will, Co-Chair, CDFO
Mr. John Candy, CDFO
Ms. Kim Charlie, Sts'ailes
Ms. Marla Maxwell, CDFO
Mr. Joe Tadey, CDFO

United States Members

Mr. Jay Zischke, Co-Chair, Suquamish Indian Tribe
Mr. Scott Bass, PNPTC
Mr. Bill Patton, NWIFC
Dr. Maureen Small, WDFW
Dr. Gary Winans, NOAA

MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2017

Canadian Members

Mr. Pieter Van Will, Co-Chair, CDFO
Mr. John Candy, CDFO
Ms. Kim Charlie, Sts'ailes
Ms. Marla Maxwell, CDFO
Mr. Joe Tadey, CDFO

United States Members

Mr. Bill Patton, Co-Chair, NWIFC
Mr. Scott Bass, PNPTC
Dr. Maureen Small, WDFW
Dr. Gary Winans, NOAA
Mr. Ben Starkhouse, Lummi Nation

MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2018

Canadian Members

Mr. Pieter Van Will, Co-Chair, CDFO
Mr. John Candy, CDFO
Ms. Kim Charlie, Sts'ailes
Ms. Brittany Jenewein, CDFO
Ms. Marla Maxwell, CDFO
Mr. Joe Tadey, CDFO

United States Members

Mr. Bill Patton, Co-Chair, NWIFC
Mr. Scott Bass, PNPTC
Dr. Maureen Small, WDFW
Mr. Ben Starkhouse, Lummi Nation
Dr. Gary Winans, NOAA

LIST OF ACRONYMS WITH DEFINITIONS

| | |
|---------------|--|
| B.C. | British Columbia |
| CDFO | Canadian Department of Fisheries and Oceans or Fisheries and Oceans Canada |
| CPUE | Catch per Unit Effort |
| CU | Conservation Unit (Canada) |
| CWT | Coded-Wire Tag |
| ECVI | East Coast Vancouver Island |
| EO | Economic Opportunity |
| ESSR | Excess Salmon to Spawning Requirements |
| ESU | Evolutionarily Significant Unit |
| FN | First Nation |
| FN FSC | First Nations Food, Social and Ceremonial |
| FSC | Food, Social, and Ceremonial (Canada) |
| GN | Gill net |
| GSI | Genetic Stock Identification |
| ITQ | Individual Transferable Quota |
| mtDNA | Mitochondrial DNA |
| mSAT | Microsatellite DNA |
| NOAA | National Oceanic and Atmospheric Administration |
| NWIFC | Northwest Indian Fisheries Commission |
| PFMA | Pacific Fishery Management Area |
| PNPTC | Point No Point Treaty Council |
| PSC | Pacific Salmon Commission |
| PST | Pacific Salmon Treaty |
| SJF | Strait of Juan de Fuca |
| SJI/PR | San Juan Islands/Point Roberts |
| SN | Seine net |
| SNPs | Single-Nucleotide Polymorphisms |
| TCCHUM | Chum Technical Committee |
| TR | Troll |
| U.S. | United States of America |
| WA | Washington State |
| WCVI | West Coast Vancouver Island |
| WDFW | Washington Department of Fish and Wildlife |
| WSP | Wild Salmon Policy (Canada's Policy for Conservation of Wild Pacific Salmon) |

TABLE OF CONTENTS

| | |
|---|-----|
| MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2015 | i |
| MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2016 | i |
| MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2017 | i |
| MEMBERSHIP OF THE PSC JOINT CHUM TECHNICAL COMMITTEE, 2018 | i |
| LIST OF ACRONYMS WITH DEFINITIONS | ii |
| TABLE OF CONTENTS | iii |
| LIST OF TABLES | v |
| LIST OF FIGURES | vi |
| LIST OF APPENDICES | vi |
| 1 Executive Summary | 1 |
| 2 Status of Treaty Requirements | 1 |
| 3 Southern British Columbia Chum salmon | 6 |
| 3.2 Status of Treaty Requirements | 6 |
| 3.3 Conservation and Harvest Management Strategies | 6 |
| 3.3.1 Johnstone Strait Chum Salmon Management Strategy | 7 |
| 3.3.2 Strait of Georgia Chum Salmon Management Strategy | 9 |
| 3.3.3 Fraser River Chum Salmon Management Strategy | 16 |
| 3.3.4 West Coast Vancouver Island Chum Salmon Management Strategy | 18 |
| 3.4 Annual Fishery Descriptions | 20 |
| 3.4.1 Johnstone Strait | 21 |
| 3.4.2 Strait of Georgia | 22 |
| 3.4.3 Fraser River | 23 |
| 3.4.4 West Coast Vancouver Island | 27 |
| 3.5 Escapement | 28 |
| 3.5.1 Inside Southern | 29 |
| 3.5.2 West Coast Vancouver Island | 29 |
| 4 United States Chum Salmon | 30 |
| 4.1 Washington Run Sizes, Spawning Escapements, and Catches | 30 |

| | |
|---|----|
| 4.2 United States Strait of Juan de Fuca Fisheries (Areas 4B, 5, 6C) | 37 |
| 4.2.1 Management Intent..... | 37 |
| 4.2.2 Fishery Description | 37 |
| 4.3 San Juan Islands / Point Roberts Fisheries (Areas 7 and 7A)..... | 38 |
| 4.3.1 Management Intent..... | 38 |
| 4.3.2 Fishery Description | 38 |
| 5 Stock Identification | 39 |
| 5.1 Tagging of Adult Chum Salmon..... | 39 |
| 5.2 Otolith, Fin Marking and Coded Wire Tagging (CWT)..... | 39 |
| 5.2.1 Canada..... | 39 |
| 5.2.2 United States | 41 |
| 5.3 Genetic Stock Identification | 42 |
| 5.3.1 Fishery Sample Collection for Genetic Stock Identification | 43 |
| 5.3.2 Baseline Collection for Genetic Stock Identification | 44 |
| REFERENCES CITED..... | 47 |
| APPENDICES | 48 |
| Appendix D. Concept proposals (see work plans) submitted to Pacific Salmon Commission Southern Endowment Fund..... | 55 |

LIST OF TABLES

| | |
|---|----|
| Table 3-1. Aggregate Inner South Coast Chum total return, harvest and spawning escapements. Comparison of fixed target and actual harvest rates in the Johnstone Strait fisheries. 2006–2015 ¹ | 8 |
| Table 3-2. Current spawning escapement goal, hatchery requirements and overall escapement goal for Chum salmon returning to the Strait of Georgia by PFMA's and system. | 10 |
| Table 3-3. Key decision points for Fraser River Chum salmon management. | 17 |
| Table 3-4. Estimated harvest of summer Chum salmon by PFMA for commercial, research and test fishing vessels ¹ (July through the second week of September), 2006–2015. | 20 |
| Table 3-5. Estimated Chum salmon harvest from Chum salmon-directed fisheries occurring in Johnstone Strait by PFMA, 2015..... | 22 |
| Table 3-6. Estimated Chum salmon harvest from Chum salmon-directed fisheries occurring in the Strait of Georgia by PFMA, 2015. | 23 |
| Table 3-7. Estimated harvest of Chum salmon from salmon fisheries occurring in Area 29 (both marine and in-river) and in Region 2 (non-tidal) of the Fraser River, 2015. | 24 |
| Table 3-8. Estimated harvest of Chum salmon from First Nations and commercial salmon fisheries occurring in Area 29 and Fraser River, 2006-2015. | 25 |
| Table 3-9. Estimated harvest of Chum salmon in assessed (i.e. surveyed) lower Fraser River recreational fisheries, 2006–2015..... | 27 |
| Table 3-10. Nitinat area harvest, hatchery broodstock collection and spawning escapement estimates by PFMA, 2006–2015. | 28 |
| Table 3-11. Inside Southern Chum salmon net escapement estimates (spawning escapement plus hatchery broodstock) for Fraser and Non-Fraser stock aggregates, 2006–2015 ¹ | 29 |
| Table 3-12. WCVI Chum salmon spawning escapement estimates ¹ by PFMA, 2006–2015. | 30 |
| Table 4-1. Puget Sound summer Chum salmon pre-and post-season estimates of run size and spawning escapements, 2006–2015..... | 31 |
| Table 4-2. Washington fall Chum salmon pre-and post-season estimates of run size and spawning escapements, 2006–2015..... | 34 |
| Table 4-3. Washington winter Chum salmon pre- and post-season estimates of run size and spawning escapements, 2006–2015..... | 35 |
| Table 4-4. Harvest of Chum salmon in the Strait of Juan de Fuca (SJF) and the San Juan Islands (SJI) commercial fisheries during the HC/JDF summer Chum salmon accounting period (July 1–September 15) in years 2006–2015. | 36 |
| Table 4-5. Harvest of summer, fall, and winter Chum salmon in SJF, SJI/PR, Puget Sound and Washington coastal areas, 2006–2015 ¹ | 37 |

| | |
|---|----|
| Table 4-6. Harvest of Chum Salmon in U.S. Areas 6, 7 and 7A Chum fisheries during 2015. ... | 38 |
| Table 4-7. Bycatch of Chinook salmon, Coho salmon and Steelhead during U.S. commercial fisheries in Areas 6, 7 and 7A Chum salmon fisheries during 2015. | 39 |
| Table 5-1. Releases of Chum salmon with thermally-marked otoliths from WCVI based Canadian hatchery facilities, 2006–2015 | 40 |
| Table 5-2. Releases of marked Chum salmon from southern B.C. based Canadian hatchery facilities, 2006 to 2015 | 41 |
| Table 5-3. Numbers of Hood Canal summer Chum salmon released with thermally-marked otoliths or an adipose fin clip by broodyear, 2006–2015 (Point No Point Treaty Tribes and Washington Department of Fish and Wildlife 2015). | 42 |
| Table 5-5. Chum salmon tissue samples collected in Johnstone Strait fisheries and analyzed for GSI, 2006–2015..... | 43 |
| Table 5-6. Chum salmon tissue samples from Areas 7 and 7a analyzed for GSI in 2007-2015. . | 44 |
| Table 5-7. Fall Chum salmon GSI tissue collections from Southern B.C. by Conservation Unit, collection site, life stage, and influence ¹ of enhancement, 2015. | 45 |
| Table 5-8. Chum salmon GSI tissue collections from WA State, 2015. | 46 |

LIST OF FIGURES

| | |
|---|----|
| Figure 2-1. Southern Chum Strategic Plan developed by the joint Chum Technical Committee in 2012. | 2 |
| Figure 4-1. Puget Sound summer Chum salmon abundance, 2006-2015, by region of origin (JDF: Strait of Juan de Fuca, HOOD: Hood Canal, SPS: South Puget Sound)..... | 31 |
| Figure 4-2. Puget Sound fall Chum salmon abundance, 2006-2015, by region of origin (JDF: Strait of Juan de Fuca, NOOK: Nooksack-Samish, SKAG: Skagit, STSNO: Stillaguamish/Snohomish, SPS: South Puget Sound, HOOD: Hood Canal). | 32 |
| Figure 4- 3. South Puget Sound winter Chum salmon abundance, 2006-2015. | 33 |

LIST OF APPENDICES

| | |
|---|----|
| Appendix A. ANNEX IV, CHAPTER 6, OF THE PACIFIC SALMON TREATY | 48 |
| Appendix B-1. United States commercial salmon catch areas for the Strait of Juan de Fuca and Northern Puget Sound. | 51 |
| Appendix B-2. Canadian Pacific Fishery Management Areas (PFMA) for Southern B.C. - Marine areas. | 52 |

| | | |
|---------------|--|----|
| Appendix B-3. | Canadian Pacific Fishery Management Areas (PFMA) for Southern B.C. - Lower Fraser River and surrounding areas. | 53 |
| Appendix C. | PSC Work Plan for the Joint Chum Technical Committee, 2015. | 54 |
| Appendix D. | Concept proposals (see work plans) submitted to Pacific Salmon Commission Southern Endowment Fund | 55 |

1 Executive Summary

This Pacific Salmon Commission (PSC) Joint Chum Technical Committee report presents the information on Chum salmon stocks and fisheries in southern British Columbia (B.C.) and Washington (WA) for the year 2015 to address the specific provisions and requirements of Chapter 6, Annex IV (Chum Annex) of the Pacific Salmon Treaty (PST or Treaty) (Appendix A). The Treaty between the governments of Canada and the United States of America (U.S.) concerning Pacific salmon is designed to facilitate co-operation in the management, research and enhancement of Pacific salmon stocks. The Chum Annex requires that Canada and the U.S. maintain a Joint Chum Technical Committee reporting to the Southern Panel and the Commission and that certain fisheries for Chum salmon in southern B.C. and WA be managed in a specified manner (Appendix A). Certain fisheries of each country, while not specifically mentioned in the PST, are known to harvest Chum salmon originating in the other country.

This report presents various aspects of Chum salmon found in B.C. waters between Vancouver Island and the mainland, off the west coast of Vancouver Island, and in WA waters. This report also discusses the management actions of Canada and the U.S. in relation to the PST requirements for Chum salmon and provides a summary of the last 10 years of catch and escapement information for Chum salmon of concern to the Treaty. Returns in 2015 were below the recent 9-yr average in B.C. and slightly below in WA. Catch overage was noted in US 7/7A fisheries 2014 and a portion of the overage was paid back in 2015 consistent with the Treaty. The Chum Technical Committee continued work on components of the strategic plan outlined in the 2010 report, which included collecting and exchanging tissue samples from mixed-stock fisheries and spawning escapements and run reconstruction model development.

2 Status of Treaty Requirements

Chum salmon stocks and fisheries in southern B.C. and in U.S. Areas 4B, 5, 6C, 7, and 7A (Appendix B) were managed under the terms set out in Chapter 6, Annex IV (Appendix A). The following provides a brief synopsis of the Canadian and U.S. management actions taken to meet its provisions in 2015.

Paragraph 1:

The Parties were to maintain a Joint Chum Technical Committee (“the Committee”) reporting, unless otherwise agreed, to the Southern Panel and the Commission. [Specific tasks for the Committee are listed in Annex IV Chapter 6, Paragraph 1. Some of these include review stock status, develop new methods for stock management and report on management and research findings.]

The Committee convened three times in 2015: during the PSC Post-Season Meeting in January, during the PSC Annual Meeting in February, and during the Joint Chum Technical Meeting in May. In 2010 the committee developed a strategic plan to guide development and application of a jointly agreed genetic baseline for Chum salmon stocks in southern BC and Washington and eventual development of the Chum Genetic and Environmental Model (ChumGEM) (Figure 2-1). Four proposals were submitted to the PST Southern Boundary Restoration and Enhancement Fund (Southern Fund) to carry forward priorities identified by the TCCHUM. Additional detail

of the work performed by the Committee in 2015 can be found in the Committee's 2015 Work Plan (Appendix C).




Figure 2-1. Southern Chum Strategic Plan developed by the joint Chum Technical Committee in 2012.

Paragraph 2:

When the Parties provide stock composition information for fisheries, the Committee shall evaluate and report its conclusions using bilaterally agreed upon methods.

For this report, the Committee continued to use historical stock composition information to estimate Chum salmon stock composition for 2015 fisheries. These historical stock composition estimates may no longer be representative of current stock composition, and the Committee is continuing to evaluate and implement new methods following the Committee's Strategic Plan (Figure 2-1) of which improved stock identification methodologies play a significant role. The initial stages of this Strategic Plan have been implemented and work on the development and implementation of the final stage of the plan is ongoing.

Paragraph 3:

Canada and the United States shall assess catch levels and make attempts to collect additional genetic samples from any Chum salmon caught during the July 1 through September 15 time period in the boundary area fisheries (U.S. Areas 4B, 5, 6C, 7 and 7A; Canadian Areas 18, 19, 20, 21, and 29).

Table 3-4 and Table 4-4 provide the harvest of Chum salmon during the period of summer Chum salmon migration in boundary areas. Due to the low numbers of Chum salmon encountered, neither party collected samples for genetic stock identification (GSI) during this time period.

Paragraph 4:

During the period from July 1 through September 15, Canada will require the live release of Chum salmon from all purse seine gear fishing in the Strait of Juan de Fuca (Canadian Area 20) and the United States will require the same for the non-Indian seine fisheries in Areas 7 and 7A. Note: By U.S. regulation, purse seine fisheries are not permitted in U.S. Areas 4B, 5 and 6C.

Regulations were implemented by both countries to require the live release of Chum salmon in these areas during this time period.

Paragraph 5:

Canada will manage its Johnstone Strait, Strait of Georgia, and Fraser River Chum salmon fisheries to provide continued rebuilding of depressed naturally spawning Chum salmon stocks, and, to the extent practicable, not increase interceptions of U.S. origin Chum salmon. Terminal fisheries conducted on specific stocks with identified surpluses will be managed to minimize interception of non-targeted stocks.

Table 3-1 provides an evaluation of the performance of the current Johnstone Strait management strategy (2006–2015). Historical GSI results and genetic estimates from 2006 through 2015 continued to indicate a low contribution of U.S. stocks in Johnstone Strait, Strait of Georgia and Fraser River Chum salmon fisheries.

Management strategies (see section 3.3) and annual fishery descriptions (see section 3.4) indicate that Canada's Johnstone Strait, Strait of Georgia, and Fraser River Chum salmon fisheries in addition to the terminal area fisheries, were managed to meet the provisions of Paragraph 5 of Annex IV, Chapter 6, of the PST.

Paragraph 6:

Canada will manage its Johnstone Strait mixed stock fishery as follows:

- a) Inside Southern Chum salmon levels of less than 1.0 million as estimated by Canada are defined, for purposes of this chapter, as critical.*
- b) For run sizes above the critical threshold, Canada will conduct fisheries with an exploitation rate of up to 20% in Johnstone Strait of Inside Southern Chum salmon; and*

- c) *When run sizes are expected to be below the critical threshold, Canada will notify the United States and will only conduct assessment fisheries and non-commercial fisheries. Commercial fisheries targeting Chum salmon will be suspended.*

While no specific estimates of Inside Southern Chum salmon abundance were provided, Canada notified the U.S. in early October, on the basis of inside area fisheries, that the return abundance of Inside Southern Chum salmon was likely above the critical threshold. Canada continued to manage the Johnstone Strait fisheries to target to 20% of Inside Southern Chum.

Paragraph 7:

Canada will manage its Fraser River fisheries for Chum salmon as follows:

- a) *For Fraser River terminal area run sizes, identified in-season, at abundance levels lower than 900,000 Chum salmon, the Canadian commercial Chum salmon fisheries within the Fraser River and in associated marine areas (Area 29), will be suspended; and*
- b) *For Fraser River terminal area run sizes, identified in-season at levels greater than 900,000 Chum salmon, Canadian commercial Chum salmon fisheries within the Fraser River, shall be guided by the limits of the in-river Total Allowable Catch set by Canada.*

In 2015, initial in-season abundance estimate indicated a terminal return above the specified Fraser River gross escapement threshold (see section 3.4.3).

Paragraph 8:

Canada will manage the Nitinat gill net and purse seine fisheries for Chum salmon to minimize the harvest of non-targeted stocks.

Due to above goal abundance of Nitinat Chum in 2015, both commercial gill net and purse seine fisheries occurred (see section 3.4.4.1; Table 3-10). To minimize the harvest of non-targeted stocks a variety of management actions were applied to this fishery (see section 3.3.4.1).

Paragraph 9:

Canada shall conduct a genetic sampling program of Chum salmon taken in the West Coast Vancouver Island troll fishery if early-season catch information indicates that catch totals for the July 1 through September 15 season may reach levels similar to 1985 and 1986. Sampling, should it occur, will include catches taken from the southern areas (Canadian Areas 121–124).

Chum salmon catch levels in the 2015 West Coast Vancouver Island troll fishery were significantly below the 1985 and 1986 levels. Therefore, no GSI sampling occurred.

Paragraph 10:

The United States will manage its Chum salmon fishery in Areas 7 and 7A as follows:

- a) *Inside Southern Chum salmon levels of less than 1.0 million as estimated by Canada are defined, for purposes of this chapter, as critical;*
- b) *For run sizes below the critical threshold, the U.S. catch of Chum salmon in Areas 7 and 7A shall be limited to Chum salmon taken incidentally to other species and in other minor fisheries, but shall not exceed 20,000, provided that catches for the purpose of genetic stock identification sampling shall not be included in the aforementioned limit;*
- c) *For run sizes above the critical threshold, the base catch ceiling for the U.S. Chum salmon fisheries in Areas 7 and 7A will be 130,000 Chum salmon;*
- d) *Canada will provide a run size estimate of Chum salmon entering the Fraser River no later than October 22. If the estimate is less than 900,000, the U.S. will limit its fishery impacts on Fraser River Chum salmon by restricting catch in Areas 7 and 7A to not exceed 20,000 additional Chum salmon from the day following the date the U.S. is notified. The total catch is not to exceed the catch ceiling of 130,000 Chum salmon;*
- e) *U.S. commercial fisheries for fall Chum salmon in Areas 7 and 7A will not occur prior to October 10;*
- f) *The U.S. will manage the Areas 7 and 7A fisheries for Chum salmon with the intent to minimize the harvest of non-targeted species;*
- g) *No U.S. catch shortfalls may be accrued; however any overages shall be carried forward as indicated in (h) and (i);*
- h) *Due to management imprecision, a catch in the U.S. of up to 135,000 Chum salmon will not result in an overage calculation. Catches in excess of 135,000 Chum salmon shall result in an overage being calculated by subtracting 130,000 from the total Chum catch. Overages will be accounted for by reducing the U.S. annual catch ceiling in up to two subsequent non-critical Inside Southern Chum salmon years; and*
- i) *From the day following the date the U.S. is notified of a run size below the critical threshold as defined in 10(b) or (d), any catches in excess of 20,000 Chum salmon will result in an overage. Overages will be accounted for by reducing the U.S. annual catch ceilings in up to two subsequent non-critical Inside Southern Chum salmon years.*

In 2015, U.S. fisheries in Areas 7 and 7A were managed in accordance with the above provisions. U.S. commercial fisheries were initiated as scheduled on October 10. An estimated Fraser River Chum salmon run size was provided by Canada prior to October 22, consistent with the Treaty obligations. The fishery continued through November 7. A portion of the U.S. 2014 catch ceiling overage was paid back during 2015 (see section 4.3.2).

Paragraph 11:

The United States shall conduct its Chum salmon fishery in the Strait of Juan de Fuca (United States Areas 4B, 5, and 6C) so as to maintain the limited effort nature of this fishery, and, to the extent practicable, not increase interceptions of Canadian origin Chum

salmon. The United States shall continue to monitor this fishery to determine if recent catch levels indicate an increasing level of interception.

This fishery is restricted to Treaty Indian fishers from four tribes and to gill net gear only. Due to the limited effort of the Juan de Fuca fishery, the Committee has not sampled the fishery for contribution of Canadian origin Chum salmon since 1996. Genetic stock identification (GSI) samples collected from this fishery in prior years indicated the majority of the catch is Chum salmon of U.S. origin (Beattie et al. 1996).

Paragraph 12:

All information concerning by-catch of other salmon species from the Chum salmon fisheries covered by this chapter will be shared between the Parties in the annual Post Season Report.

By-catch information for other salmon species was shared in accordance with the Treaty through each Party's post-season reporting process.

Paragraph 13:

Should circumstances arise that are inconsistent with either Parties understanding of the intent of the chapter, the Southern Panel will discuss the matter postseason and explore options for taking the appropriate corrective actions.

No such circumstances arose in 2015.

3 Southern British Columbia Chum salmon

Southern B.C. Chum salmon stocks and fishing areas (Pacific Fishery Management Areas or PFMAs; Appendix B) are, for the purposes of management, analysis and reporting, grouped into four major units: Johnstone Strait, Strait of Georgia, Fraser River, and the West Coast Vancouver Island (WCVI). Additionally, Chum salmon originating from Johnstone Strait, the Strait of Georgia, and the Fraser River are collectively described as Inside Southern Chum salmon.

3.2 Status of Treaty Requirements

During 2015, the Southern B.C. Chum salmon fisheries were managed according to the requirements of Annex IV, Chapter 6, as amended in 2009 (Appendix A).

3.3 Conservation and Harvest Management Strategies

The general management approach adopted by Fisheries and Oceans Canada (CDFO) for Southern B.C. Chum salmon is to achieve wild escapement targets, while augmenting production through enhancement of selected stocks. In practice, this approach is achieved through the application, in mixed stock fisheries, of harvest rates which are compatible with wild or natural stock productivity. If there are Chum salmon stocks that return to their area of origin in numbers above that area's escapement goal, they may be subject to additional harvesting within their terminal areas.

The following describes the management strategies and fishing plans for the four units of Southern B.C. Chum salmon stocks.

3.3.1 Johnstone Strait Chum Salmon Management Strategy

Since 2002, the mixed stock Chum directed fisheries in Johnstone Strait have been managed to approximate a fixed exploitation rate (~20%). Tagging studies conducted from 2000 through 2002 helped in the development of this fixed exploitation rate strategy by assessing the migration timing and harvest rate by fishing gear on an available abundance of Chum salmon in Johnstone Strait. This fixed exploitation strategy continued to be employed through 2015 (CDFO 2015).

Some of the key objectives of this strategy are to ensure sufficient escapement levels while providing more stable fishing opportunities (CDFO 2015). The exploitation rate is set at 20% across all harvesters, when abundance is estimated to be above a critical level. The critical level, or threshold, was determined to be 1.0 million Chum salmon (Appendix A). The Johnstone Strait Chum salmon test fishery, as well as the information gained from any commercial openings, is used in-season to determine the likelihood of achieving the critical level for Inside Southern Chum salmon. When expected levels are less than 1.0 million, only assessment fisheries and non-commercial fisheries will be conducted and any planned commercial fisheries targeting Chum salmon will be suspended (Appendix A). When the critical level is expected to be achieved, fisheries with an exploitation rate of up to 20% in Johnstone Strait can be conducted (Appendix A).

Of this 20%, the commercial sector is allocated 15%. The remaining 5% is set aside to provide for First Nations' food, social and ceremonial needs (FSC) satisfy recreational and test fishing requirements, and to provide a buffer to the commercial exploitation (CDFO 2015). The impact of the Johnstone Strait fisheries during fixed exploitation approach (2006–2015), on Inside Southern Chum salmon stocks are detailed in Table 3-1.

The specific objectives of the fixed exploitation rate strategy are to:

- continue to rebuild/maintain stocks to the optimum wild escapement objective (defined as 2.5 million wild Inside Southern Chum salmon);
- establish a pre-season fishing plan to achieve an exploitation rate of 20% on Inside Southern Chum salmon in the Johnstone Strait Chum salmon fishery; and,
- stabilize commercial catch rates to provide opportunities at both low and high abundance levels

Table 3-1. Aggregate Inner South Coast Chum total return, harvest and spawning escapements. Comparison of fixed target and actual harvest rates in the Johnstone Strait fisheries. 2006–2015¹

| | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 |
|--|-----------|-----------|--------------------------|---------------|---------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| PRE-SEASON | | | | | | | | | | |
| Inside Southern Abundance Forecast (or Outlook) | Average | Average | Average to Above Average | Below Average | Below Average | Average to Below Average | Average to Below Average | Average to Below Average | Average to Below Average | Average to Below Average |
| POST-SEASON | | | | | | | | | | |
| Inside Southern Abundance² | 4,301,780 | 2,728,082 | 2,228,121 | 2,227,113 | 1,181,534 | 3,449,783 | 3,501,098 | 3,156,129 | 2,354,404 | 2,751,791 |
| Inside Southern Harvest | 1,527,453 | 796,301 | 580,247 | 747,046 | 127,973 | 1,441,680 | 911,427 | 1,079,567 | 732,429 | 1,240,133 |
| Est. Inside Southern Harvest Rate | 35.51% | 29.19% | 26.04% | 33.54% | 10.83% | 41.79% | 26.03% | 34.21% | 31.11% | 45.07% |
| <u>Johnstone Strait Harvest</u> | | | | | | | | | | |
| Commercial, Sport Area 11-13³ | 859,201 | 480,620 | 299,775 | 513,884 | 55,520 | 750,436 | 392,608 | 635,814 | 332,140 | 529,541 |
| First Nations Area 11-13 | 16,453 | 14,870 | 11,630 | 13,576 | 4,060 | 9,880 | 13,106 | 13,444 | 113 | 22,472 |
| Johnstone Strait Harvest Total | 875,654 | 495,490 | 311,405 | 527,460 | 59,580 | 760,316 | 405,714 | 649,258 | 332,253 | 552,013 |
| Target Johnstone Strait Harvest Rate | 20% | 20% | 20% | 20% | 20% | 20% | 20% | 20% | 20% | 20% |
| Est. Johnstone Strait Harvest Rate | 20.36% | 18.16% | 13.98% | 23.68% | 5.04% | 22.04% | 11.59% | 20.57% | 14.11% | 20.06% |
| ESCAPEMENT (Includes wild and enhanced)⁴ | | | | | | | | | | |
| Inside Southern Escapement | 2,774,327 | 1,931,781 | 1,647,874 | 1,480,067 | 1,053,561 | 2,008,103 | 2,589,671 | 2,076,562 | 1,621,975 | 1,511,658 |

(1) Historic data in this table have been updated with most recent estimates; values may deviate from past reports.

(2) Total Inside Southern abundance includes total Inside Southern harvest plus escapement. Harvest composition based on historic GSI for all fisheries.

(3) Includes commercial, sport and test fishery harvest.

(4) Escapement estimates do not include any removals associated with hatchery rack (e.g. ESSR); those values are included in harvest. Escapement estimates have not been expanded for populations not monitored for escapement

3.3.2 Strait of Georgia Chum Salmon Management Strategy

Chum salmon stocks return to all areas of the Strait of Georgia (mid-Vancouver Island PFMA's 14-19) and are of enhanced and wild (natural) origin. Fisheries can target enhanced returns (e.g. Area 14: Puntledge, Big Qualicum and Little Qualicum rivers), wild returns (e.g. Area 16: main river systems of Tzoonie, Deserter and Skwawka rivers in the Jervis inlet area) and mixed returns (e.g. Area 19: directed primarily at enhanced Goldstream River population although some natural Cowichan River Chum salmon are also harvested).

With one exception, pre-season forecasts of the return of Strait of Georgia Chum salmon for population aggregates corresponding to PFMA's have limited use due to their inability to reasonably predict returns (CDFO 2015). The one exception is Chum salmon returning to Area 14. As a result, unlike some other units of Southern BC Chum Salmon (e.g. WCVI see section 3.3.4), pre-season fishing plans based on a pre-season forecast are not typically developed for Strait of Georgia Chum salmon stocks.

Strait of Georgia Chum salmon are managed as a component of a "mixed-stock harvest strategy" for Johnstone Strait and the northern Strait of Georgia. In-season management of Strait of Georgia Chum salmon is guided by advice from the South Coast Chum Advisory Committee (Chum Advisory Committee or Chum Working Group). The Chum Working Group represents interests for Strait of Georgia (mid-Vancouver Island Areas 14-19), Johnstone Strait, WCVI and Fraser River Chum salmon-directed fisheries. Fishing opportunities for Chum salmon are evaluated at weekly meetings of the Chum Working Group that usually start in the first week of October. In-season data is reviewed on a weekly basis until the conclusion of the fishing season that usually occurs around the end of November. The Chum Working Group includes internal (CDFO staff) and external (First Nations, commercial and recreational) representation and has been in operation since 2004. Additionally, Chum salmon management in Area 18 is also guided by advice from the Cowichan Fisheries Roundtable.

Chum salmon fishing opportunities in the Strait of Georgia PFMA's are predominately shaped by in-season escapement information, test fishing data, commercial harvesting data from Johnstone Strait fisheries when available, and coast-wide allocations of Chum salmon. When escapement goals are achieved (Table 3-2) or when stock abundance allows, commercial gill net, purse seine and troll fisheries can be permitted. As mentioned, the exception is Area 14 where a pre-season forecast of returning Chum abundance is used in addition to the in-season information listed above, to guide management decisions on fishing opportunities (CDFO 2015).

Chum salmon stocks return to all areas of the Strait of Georgia (mid-Vancouver Island PFMA's 14-19) and are of enhanced and wild (natural) origin. Fisheries can target enhanced returns (e.g. Area 14: Puntledge, Big Qualicum and Little Qualicum rivers), wild returns (e.g. Area 16: main river systems of Tzoonie, Deserter and Skwawka rivers in the Jervis inlet area) and mixed returns (e.g. Area 19: directed primarily at enhanced Goldstream River population, although some natural Cowichan River Chum salmon are also harvested).

Table 3-2. Current spawning escapement goal, hatchery requirements and overall escapement goal for Chum salmon returning to the Strait of Georgia by PFMA's and system.

| PFMA | System | Spawning Escapement Goal | Hatchery Requirements | Overall Escapement Goal |
|---------|-----------------------|--------------------------|-----------------------|-------------------------|
| Area 14 | Puntledge River | 60,000 | 10,000 | 240,000 |
| | Big Qualicum River | 85,000 | | |
| | Little Qualicum River | 85,000 | | |
| Area 16 | Jervis Inlet Streams | n/a | n/a | 110,000 |
| Area 17 | Nanaimo River | n/a | n/a | 40,000 |
| Area 18 | Cowichan River | n/a | n/a | 160,000 |
| Area 19 | Goldstream River | n/a | n/a | 15,000 |

Fishing opportunities in the Strait of Georgia are constrained by concerns over bycatch of passing Chum salmon stocks and local Chinook and Coho salmon populations. These concerns are addressed by Area specific pre- and in-season management measures that may include:

- *area closures*: beach boundaries and limiting fisheries to terminal areas to minimize impacts on passing stocks; and,
- *non-retention*: of Chinook and Coho salmon.

As well, to reduce bycatch mortality and to improve the quality of catch data, additional pre-and in-season management measures may be implemented in the Strait of Georgia fisheries. These additional measures may include:

- *daylight fishing only*: in the gill net fishery;
- *maximum soak time*: in the gill net fishery;
- *mandatory brailing*: in the seine fishery;
- *barbless hooks*: in the troll fishery; and,
- *on-board observers*: if high bycatch occurs.

Pre- and in-season management measures aimed at reducing bycatch mortality are commonly implemented after consultation with the Chum Working Group.

First Nations FSC fisheries remain a priority and occur in terminal areas targeting individual populations from river systems within or adjacent to band territories. Harvest limits may be set when there are conservation concerns for the target population; however, First Nations FSC fisheries have priority over other fisheries. Generally, catch and effort in First Nations FSC Chum salmon fisheries is low compared to commercial Chum salmon fisheries.

Although tidal recreational fisheries are open throughout the Strait of Georgia, Chum salmon are not commonly targeted by recreational anglers. Nevertheless, the Strait of Georgia recreational Chum salmon catch is estimated annually through a Creel Survey conducted by CDFO. In most years, the Creel Survey operates from the beginning of May through to the end of September. Data from this survey is compiled and analyzed to produce catch and effort statistics by Area for each salmon species.

Non-tidal recreational fisheries vary by Area and are considered if escapement and FSC needs have been met.

3.3.2.1 Area 14 Chum Salmon Management Strategy

Chum salmon returning to Area 14 have been enhanced since the late 1960's, and terminal fisheries targeting the enhanced Chum salmon populations of Puntledge, Big Qualicum and Little Qualicum rivers have occurred in October and November since the early 1970's. Returning Area 14 Chum salmon abundance is forecasted pre-season using brood escapement, average survival, and age composition. In-season run strength is assessed from any early catch information, visual observations at river estuaries, and escapement counts to the three enhanced river systems.

The interim spawning escapement goals for Chum salmon to the three river systems are 60,000 to Puntledge River, 85,000 to Big Qualicum River, and 85,000 to Little Qualicum River; including enhancement facility requirements of 10,000, the overall escapement goal is 240,000 Chum salmon for Area 14 (Table 3-2). Spawning escapements are monitored by CDFO Stock Assessment and hatchery staff.

The recommended approach regarding Chum salmon-directed fishing opportunities in Area 14 is made at the first meeting of the Chum Working Group. This meeting is tentatively scheduled for the first week of October; Chum Working Group meetings are subsequently held every week until late November.

Commercial Chum salmon-directed fisheries have a specific harvest strategy, implemented since 1981 that consists of limited early harvest prior to any escapement occurring. The allowable early Chum salmon harvest is set at 65% of the predicted surplus. The predicted surplus is calculated by subtracting the overall escapement goal (240,000; Table 3-2) and a buffer (100,000) from the pre-season terminal run size forecast (CDFO 2015). The buffer safeguards against errors in the pre-season forecast of Area 14 stock abundance.

The limited early commercial harvest of Chum salmon is planned pre-season and executed in-season in the second or third week of October based on whether a surplus, using the formula described above, is identified. However, if catches in the commercial Chum salmon fisheries in Johnstone Strait indicate low Chum salmon abundance, a decision on the approach in Area 14 may be deferred until the following week.

The management objectives for Area 14 are:

- achieve Area 14 Chum salmon escapement requirements of 240,000;
- ensure adequate Chinook and Coho salmon escapements to Area 14 enhancement facilities;
- provide access to First Nations for FSC purposes;
- minimize the harvest of passing salmon populations;
- maximize economic return;
- work towards south coast Chum salmon allocation targets for gill net, seine net and troll sectors;
- attempt to manage initial fisheries in Area 14 to avoid large surpluses (e.g. greater than 100,000).

Starting in the second or third week of October, opportunities for gill net, seine net and troll fisheries are based on pre-season forecasts, in-season catch per unit effort (CPUE) information from commercial Chum salmon fisheries in Johnstone Strait, and escapement information. Escapement information

becomes increasingly important when considering further commercial opportunities in the latter part of October. These commercial opportunities are determined at the weekly Chum Working Group meetings and may include consideration for:

- *limited effort seine net fishery*: a limited effort seine fishery with a harvest target will be considered from late October to late November, based on Chum salmon escapement, abundance in the approach areas and South Coast allocation guidelines. Full fleet opportunities may also be available;
- *expanded gill net and troll fishery*: if gear counts indicate a modest fleet size of 50 vessels or less, gill net and troll openings may be expanded beyond one to two days per week; as well, additional fishing opportunities for gill net and troll may be considered following the seine net fisheries; and,
- *additional fishing days*: additional fishing days are considered if fishing time is lost due to poor weather conditions.

Once spawning escapement goals and hatchery requirements have been achieved, any returns above the overall escapement goal may be harvested (CDFO 2015).

Commercial fishing opportunities are constrained by concerns over local and passing Chinook, Coho and Chum salmon populations. Management measures to address these concerns can include:

- *area closures*: beach boundaries to protect Chinook and Coho salmon (boundaries may range from half a mile to one and a half miles depending upon bycatch concerns and time of year); French Creek radius boundary and Baynes Sound area closures to protect wild Chum and Coho salmon populations;
- *non-retention*: of Coho salmon;
- *maximum soak time*: in the gill net fishery;
- *mandatory brailing*: in the seine fishery;
- *barbless hooks*: in the troll fishery;
- *daylight fishing only*: in the gill net fishery if there are significant levels of bycatch; and,
- *outside boundaries*: designed to minimize impacts on passing salmon populations.

First Nations FSC fisheries are conducted in Area 14 and at the enhancement facilities prior to consideration of ESSR fisheries targeting Chum salmon. Tidal recreational fisheries are subject to the normal daily and possession limits for Chum salmon (daily limit four per day/possession limit eight) and are open throughout the Area.

Once escapements have been confirmed, non-tidal recreational fisheries for Chum salmon in the Puntledge and Big Qualicum rivers will be considered. These fishing opportunities may occur as early as the second to fourth week of October and are based, in part, upon in-season and past return timing of Chum salmon.

Chum salmon returning to Area 14 have been enhanced since the late 1960's, and terminal fisheries targeting the enhanced Chum salmon populations of Puntledge, (Big) Qualicum and Little Qualicum rivers have occurred in October and November since the early 1970's. Returning Area 14 Chum salmon abundance is forecast pre-season using brood escapement, average survival and age composition. In-

season run strength is assessed from any early catch information, visual observations at river estuaries, and escapement counts to the three enhanced river systems.

3.3.2.2 Area 16 Chum Salmon Management Strategy

This fishery targets wild Chum salmon stocks returning to river systems in the Jervis Inlet area. The main systems are Tzoonie, Deserted and Skwawka rivers. The overall escapement goal for Jervis Inlet streams is 110,000. These terminal fisheries occur when the individual or combined escapement goals have been assured.

Area 16 Chum salmon are managed as a component of “mixed-stock harvest strategy” for Chum salmon, and fishing opportunities are guided by coast-wide allocations of Chum salmon. Commercial fishing opportunities are evaluated at weekly meetings of the Chum Advisory Committee, usually starting in the first week of October. Assessment in the area is conducted by Fisheries and Oceans Canada Charter Patrol vessels, DFO Stock Assessment and Sechelt Indian Band staff. In-season data is reviewed on a weekly basis until the end of the season, which usually occurs around the end of November. Area 16 Chum fisheries are not planned based on pre-season forecasts alone. Fishing opportunities will be provided in an area when the escapement goal has been achieved. Achievement of the escapement goal includes the numbers of fish in-river plus the amount of fish inside a designated sanctuary area. The earliest potential fishing opportunity is anticipated near the end of October (CDFO 2015).

Chum salmon fishing opportunities are constrained by concerns over fishing impacts on Coho salmon and passing Chum salmon populations. To address these concerns, there is mandatory non-retention of Coho salmon and fishing is limited to terminal areas.

Historically, Chum salmon fishing opportunities in Area 16 do not occur on a regular basis. There have been no fisheries in Area 16 in recent years.

Area 16 Chum salmon-directed fisheries target wild Chum salmon populations returning to river systems in the Jervis Inlet area. The main systems are Tzoonie, Deserted, and Skwawka rivers.

3.3.2.3 Area 17 Chum Salmon Management Strategy

Chum salmon fisheries in Area 17 are directed primarily at the Nanaimo River population. On poor return years, the Nanaimo River Chum salmon stock is supplemented by production from the Nanaimo River Hatchery.

The overall escapement goal for the Nanaimo River is 40,000 Chum salmon (Table 3-2). Chum salmon escapement estimates are derived from joint DFO/Snuneymuxw in-river assessments.

Pre-season forecasts of Chum salmon abundance are helpful in defining possible opportunities, but decisions to open commercial fisheries are not based on pre-season information (CDFO 2015). Area 17 Chum salmon are managed as a component of the “mixed-stock harvest strategy” for Chum salmon, and in-season management is guided by advice from the Chum Working Group as outlined for Areas 14 and 16. Fishing opportunities are planned in-season based on escapement information and shaped by coast-wide allocations of Chum salmon. Escapements can fluctuate annually and fishery opportunities are evaluated during the weekly in-season review of Nanaimo escapement estimates within the Chum Working Group process.

Opportunities for gill net, troll and seine net fisheries are discussed once Chum salmon have started to enter the Nanaimo River and are present in terminal areas. Final decisions are made at the weekly

Chum Working Group meetings. If commercial opportunities are identified, management will be guided by the following considerations:

- gill nets open for one or two days; fishing days and opening duration subject to escapement levels;
- troll open seven days per week due to a demonstrated low catch rate;
- after initial opening, continued fishing opportunities depend upon information derived from CPUE in the commercial fisheries, and on-going approach area and in-river assessments;
- if harvest remains “good” and the escapement goal is reached, commercial fisheries can continue; and,
- additional fishing days will be considered if time is lost due to poor weather conditions.

Commercial fishing opportunities are constrained by concerns over domestic Chinook and Coho salmon populations and passing Chum salmon. Management measures to address these concerns can include:

- *area closures*: subarea boundaries protecting migrating Fraser River Chum salmon and confine the fishery to the Nanaimo River stock;
- *non-retention*: of Coho and Chinook salmon;
- *maximum soak time*: in the gill net fishery if Coho salmon encounters are high;
- *barbless hooks*: in the troll fishery; and,
- *daylight fishing only*: in the gill net fishery if there are significant levels of bycatch.

In-season management measures aimed at reducing bycatch mortality (e.g. maximum soak time and daylight fishing only in the gill net fishery) would be implemented after consultation with the Chum Working Group.

First Nations FSC fisheries as well as tidal/non-tidal recreational fisheries are conducted on these Area 17 populations. Local FSC opportunities are undertaken by Nanaimo First Nations in consultation with the DFO. Tidal recreational fisheries are subject to the normal daily and possession limits and there are no closed areas. There are no opportunities for non-tidal recreational fisheries in the Nanaimo River.

3.3.2.4 *Area 18 Chum Salmon Management Strategy*

The Area 18 fishery is directed primarily at Cowichan River stocks, although some Goldstream Chum salmon are also harvested. The outer Cowichan Chum salmon fishing boundary is situated to minimize encounters of Saanich Inlet Chum salmon. Chemainus River stocks are also impacted but likely to a lesser extent (CDFO 2015).

The escapement goal for the Cowichan River is 160,000 Chum salmon (Table 3-2). The target was revised from 110,000 in 2009, and is based on habitat area and Chum spawning densities in the Cowichan River. In-river Chum salmon escapement estimates are provided by the operation of a DIDSON Counter (Dual-frequency Identification Sonar) located in the lower river since 2006.

Pre-season forecasts of abundance are helpful in defining possible commercial opportunities, but decisions to open commercial fisheries are not based solely on pre-season information (CDFO 2015). In-season management is guided by advice from the Cowichan Fisheries Roundtable (the Roundtable) and the Chum Working Group. Commercial fishing opportunities are evaluated during the weekly in-

season meetings of the Roundtable and the Chum Working Group and are shaped by coast-wide allocations of Chum salmon.

Fishery openings are planned in-season based on escapement estimates from the DIDSON counter and information from a test fishery. Management is also guided by advice from the Cowichan Fisheries Roundtable (the Roundtable), the Chum Working Group and an in-season Chum Escapement Forecast Tool based on the DIDSON count by date (CDFO 2015)

Like other Strait of Georgia PFMA's, commercial fishing opportunities in Area 18 are constrained by concerns over domestic Chinook and Coho salmon populations and passing Chum salmon. Management measures to address these concerns can include:

- *area closures*: subarea boundaries to protect Coho salmon holding off Cherry Point; limiting commercial fisheries to Satellite Channel to minimize impact on the earlier timed Goldstream River Chum salmon populations; beach boundaries to protect Chinook and Coho salmon;
- *non-retention*: of Coho salmon;
- *maximum soak time*: in the gill net fishery if Coho salmon encounters are high;
- *mandatory brailing*: in the seine net fishery;
- *barbless hooks*: in the troll fishery; and,
- *daylight fishing only*: in the gill net fishery if there are significant levels of bycatch.

In-season management measures aimed at reducing bycatch mortality (e.g. maximum soak time and daylight fishing only in the gill net fishery) would be implemented after consultation with the Roundtable and the Chum Working Group.

First Nations FSC fisheries and tidal/non-tidal recreational fisheries are conducted on these Area 18 Chum salmon populations. Tidal recreational fisheries are subject to the normal daily and possession limits. Non-tidal recreational fisheries will also be considered if escapement targets and First Nations FSC needs are achieved.

3.3.2.5 *Area 19 Chum Salmon Management Strategy*

Chum salmon fisheries are directed primarily at the Goldstream River population although some Cowichan River Chum salmon are also harvested. The overall escapement goal for Goldstream River is 15,000 Chum salmon (Table 3-2). Goldstream River is a relatively small system with good viewing conditions; as such, escapement estimates are derived from stream walks.

Due to the inability of pre-season forecasts to reliably predict returns of Area 19 Chum salmon, pre-season fishing plans are not developed for Area 19. In-season management is guided by advice from the Chum Working Group. Commercial fishing opportunities are evaluated at the Chum Working Group's weekly meetings where in-season data is reviewed. In-season escapement information forms the basis for determining commercial fishing opportunities. Area 19 falls under the same management regime as other Strait of Georgia PFMA's; consequently, fishing opportunities are also shaped by coast-wide allocations of Chum salmon.

Commercial fishing opportunities in Area 19 are constrained by concerns over domestic Chinook and Coho salmon populations and passing Chum salmon. Management measures to address these concerns can include:

- *area closures*: subarea boundaries to protect Goldstream River Chinook and Coho salmon holding at Squally Reach and minimize encounters of Cowichan River Chum salmon; commercial fishery openings set for mid to late November are limited to portions of Saanich Inlet that are outside or to the north of Squally Reach;
- *non-retention*: of Coho and Chinook salmon;
- *maximum soak time*: in the gill net fishery if Coho salmon encounters are high;
- *mandatory brailing*: in the seine net fishery;
- *barbless hooks*: in the troll fishery;
- *daylight fishing only*: in the gill net fishery if Coho salmon encounters are high;
- *selective fishing techniques*: commercial fisheries will utilize selective fishing techniques to minimize bycatch impacts; and,
- *on-board observers*: on-board observers could be employed if high bycatch occurs.

3.3.3 Fraser River Chum Salmon Management Strategy

The management strategy for Fraser River Chum salmon includes management goals and harvest allocation objectives for Chum salmon fisheries occurring within the Fraser River. Fisheries are structured to ensure a spawning escapement of 800,000 Chum salmon using in-season information derived from the CDFO Albion Chum salmon test fishery (see section 3.3.3.1). Decision rules that guide the management of Fraser River Chum salmon fisheries are summarized in Table 3-3.

Additionally, management of in-river Chum salmon fisheries is guided by the need to minimize impacts on co-migrating stocks of concern, including Interior Fraser River Coho salmon and Interior Fraser River Steelhead. In order to limit incidental impacts on Interior Fraser River Coho salmon, fishing for salmon with non-selective gear is restricted for all sectors (First Nations, recreational, and commercial) from early September to mid-October within the Fraser River mainstem below Sawmill Creek. Conservation measures taken to protect Interior Fraser River Steelhead are developed in conjunction with the B.C. Ministry of Environment on an annual basis, and are primarily focused on reducing the impact of Fraser River commercial gill net fisheries. In recent years, these restrictions have resulted in the delay of commercial fishing opportunities for Chum salmon until late October, with the exception of those fisheries that operate with selective gear.

Table 3-3. Key decision points for Fraser River Chum salmon management.

| Run Size | Harvest Plan | First Nations | Commercial | Recreational |
|-----------------------------------|---|--|--------------------------|---|
| <500,000 in Fraser | <10% | Limited (reduced hours and days/week fishing) | Closed | Main stem Fraser River closed, restricted openings on tributaries |
| 500,000 to 800,000 in Fraser | Directed fisheries limited to FSC | Normal (72,000) | Closed | Main stem Fraser River closed, restricted openings on tributaries |
| 800,000 to 916,000 in Fraser | Catch not to exceed 81,000 (72,000 First Nations and 9,000 test fishing) | Normal (72,000) | Closed | Main stem Fraser River open, restricted openings on tributaries |
| 916,000 to 1,050,000 in Fraser | Commercial catch not to exceed 10% for Chum. | Normal (72,000) | Open (35,000-105,000) | Open |
| >1,050,000 in Fraser | Commercial catch not to exceed 15% for Chum. | Normal (72,000) | Open | Open |

3.3.3.1 Fraser River In-season Terminal Abundance Estimation

Terminal abundance of Fraser River Chum salmon is estimated using a Bayesian model that incorporates pre-season information on run size and migration timing, with in-season information on Chum salmon catch from the Albion Chum salmon test fishery (Gazey and Palermo 2000).

The Albion Chum salmon test fishery has operated annually since 1979 on the lower Fraser River in Area 29 near Fort Langley. The test fishery is conducted with a drifted gill net at a specific site near the old Albion ferry crossing. The test fishery begins in early September of each year, and usually fishes until the end of November. On each day of operation, the boat fishes two sets, timed to coincide with the daily high tide. The Albion Chum salmon test fishery normally fishes every other day from September 1st through October 20th, alternating days with the Albion Chinook salmon test fishery (which fishes an 8" mesh gill net during this period). Until 2013, the Chum salmon test fishery operated daily from October 21st through the end of November. Since 2013, the Chum salmon test fishery operates daily from October 21st through approximately November 10th and then every other day until November 23rd. The gill net used in the Albion Chum salmon test fishery is 150 fathoms long, constructed from uniform 6.75" mesh.

The first in-season estimate of terminal Fraser River Chum salmon abundance is typically provided in mid-October. Limited First Nations FSC fisheries for Fraser Chum salmon may be permitted prior to this date, unless a conservation concern has been identified.

3.3.4 West Coast Vancouver Island Chum Salmon Management Strategy

Chum salmon stocks return to all areas on the West Coast of Vancouver Island (WCVI). When stock abundance allows, commercial Chum salmon fisheries are conducted in the marine waters outside Nitinat Lake (Area 21), Barkley Sound (Area 23), Clayoquot Sound (Area 24) and Outer Nootka Sound/Thupana and Esperanza Inlet (Areas 25). Both commercial gill net and purse seine fleets target WCVI Chum salmon in these Areas.

A pre-season forecast of the return of WCVI Chum salmon is predicted for population aggregates corresponding to CDFO's Pacific Fishery Management Areas. Natural and enhanced returns are approximated through separate models and summed to produce a total return forecast for each Area. However, the relative inaccuracy of the pre-season forecasts resulted in the adoption of a fixed harvest rate strategy in the early 2000's for those fisheries targeting mixed stock aggregates.

With the exception of near-terminal areas, such as Nitinat (Area 21 & 22) or Thupana Inlet (Area 25), where hatchery stocks dominate, WCVI Chum salmon fisheries are currently managed to an exploitation rate of approximately 20% or less. Exploitation is controlled by limiting fishing effort to specific areas, specific times or to a specific number of vessel fishing days per week. Since 2012, revised fishery lower and target reference points (FLRPs and FTRPs) were developed based on the Sustainable Escapement Goal (SEG) method. The SEG approach uses the long-term escapement series to set fishery reference points. Conservative "SEGs" are defined as the 25% and 75% of a long-term escapement time series. Although the WCVI Chum forecast is highly uncertain, the forecast is used to inform pre-season fishery planning. Where the forecast is below the FLRP for an area, fisheries are curtailed. Where the forecast is below the FTRP, fisheries are more limited (CDFO 2015)

Purse seine opportunities generally occur when excess salmon to spawning requirements (ESSR) are identified in near terminal areas (e.g. hatchery stocks returning to Nitinat River or Thupana Inlet).

WCVI First Nations FSC fisheries remain a priority and occur in terminal areas targeting individual populations from river systems within or adjacent to band territories. Harvest limits may be set when there are conservation concerns for the target population; however, First Nations FSC fisheries have priority over other fisheries. Generally, harvest and effort in First Nations FSC Chum salmon fisheries is low compared to commercial Chum salmon fisheries.

Chum salmon are not commonly targeted by recreational anglers on the WCVI. Nevertheless, the WCVI recreational Chum salmon catch is estimated annually through a Creel Survey conducted by CDFO. Creel Survey information is augmented by logbook and manifest records of catch and effort submitted by lodges operating guided fishing trips on the WCVI. These data are compiled and analyzed to produce catch and effort statistics by Area for each salmon species.

3.3.4.1 Nitinat Chum Salmon Management Strategy

Of importance to the Pacific Salmon Treaty is the WCVI group of Chum salmon populations returning to the Nitinat watershed (Nitinat River and Nitinat Lake and tributaries). This Nitinat stock is the most significant WCVI group of populations and includes production from a major CDFO hatchery on the Nitinat River.

The Nitinat commercial Chum salmon fishery (Area 21), along with the ESSR fisheries (Area 22), is the largest fishery on the WCVI and targets returning Nitinat River hatchery Chum salmon. In the past, catch of non-WCVI Chum salmon has occurred in the outside seaward portion of the Nitinat fishing area. Management actions have been implemented to reduce this catch of non-WCVI Chum salmon by

restricting the Nitinat fishery to subareas closer to the terminal area (i.e. outlet of Nitinat Lake). More recently, First Nations' ESSR harvests have occurred within Nitinat Lake (Area 22). Historical Chum salmon harvest is shown in Table 3-10.

The spawning escapement target for Chum salmon returning to the Nitinat system (i.e. Nitinat River and other tributaries to Nitinat Lake) has been set at 175,000 fish (CDFO 2015). The target distribution of this spawning escapement between Nitinat River and the tributaries to Nitinat Lake has varied slightly over the last number of years; currently it is set at 125,000 into the Nitinat River and 50,000 into other Nitinat Lake tributaries. This spawning escapement requirement is set at a level to ensure that all run timing components are represented. Additional allocation for First Nations FSC requirements and hatchery requirements (i.e. broodstock) may total up to 50,000 Chum salmon. Consequently, the *gross escapement requirement* (spawning escapement + FN FSC requirements + hatchery requirements) for this Nitinat group of populations is 225,000 Chum salmon (CDFO 2015). Historical Chum salmon escapements are shown in Table 3-10.

A pre-season forecast of current year returns for the Nitinat stock is approximated based on broodyear escapements, estimated freshwater production from natural spawners, hatchery production, average age at return, environmental and ocean condition factors during brood ocean entry and estimated marine survival of each broodyear class.

A pre-season fishing plan is commonly developed by the end of the summer and is based on the forecast of current year returns. The fishery is managed through a fixed-harvest rate strategy to limit the harvest rate to approximately 25%. The allowable effort of the commercial fleets was determined from an analysis of the average exploitation rate per unit effort in Nitinat Chum salmon fisheries over the last decade.

In addition to concerns over catch of non-WCVI Chum salmon, early season opportunities are constrained by concerns over bycatch of Interior Fraser River Steelhead (Thompson River) and passing stocks of Coho salmon. Since 1995, these bycatch concerns in the commercial Nitinat fishery have been addressed by:

- *delayed opening dates*: the first commercial gill net fishery, if conducted, has been delayed until the beginning of October since 1998;
- *reduced fishing areas*: the traditional fishing area has been reduced to within a one-mile boundary between lines true south from Pachena and Dare Points (a portion of Area 21);
- *increased use of weed lines*: a 1.2m - 2.0m weed line for gill nets; and,
- *species selective fishing techniques*: non-retention of Coho salmon and Steelhead.

Additionally, to reduce bycatch mortality and to improve the quality of catch data, the following measures have been implemented in the Nitinat fisheries:

- *daylight fishing only*: gill net fishery;
- *mandatory functional revival tanks*: in purse seine and gill net fisheries;
- *on-board observers*: portions of purse seine and gill net fleet; and,
- *logbooks and weekly hail-ins*: purse seine and gill net fisheries.

The in-season management of the Nitinat fisheries (Area 21 and 22) are based on achieving the *gross escapement requirement* into the Nitinat watershed. Broodstock collection success, hatchery swim-ins,

and visual surveys of Nitinat River spawners are used to determine whether the weekly *escapement milestone levels* into Nitinat Lake are achieved (CDFO 2015). Weekly fisheries are generally scheduled in Area 21 and surrounding waters to harvest any identified surplus.

First Nations FSC fisheries remain a priority and occur in terminal areas of Nitinat (Areas 21 and 22). First Nations ESSR fisheries can also occur in Nitinat Lake (Area 22) targeting returns surplus to spawning and hatchery requirements.

3.4 Annual Fishery Descriptions

Fisheries targeting Chum salmon in Johnstone Strait, Strait of Georgia, Fraser River, and on the West Coast of Vancouver Island (WCVI) generally begin in late September and end in November. These fisheries are directed at fall Chum salmon.

In addition to these directed fisheries, a bycatch of Chum salmon may occur in fisheries directed at Sockeye and Pink salmon earlier in the season. This Chum salmon bycatch is assumed to be comprised mainly of summer Chum salmon. The summer Chum salmon harvest in boundary area fisheries are presented in Table 3-4.

The main components of the Inside South Coast (ISC) Chum return include both Fraser and non-Fraser stocks. These stocks are typically dominated by four year old fish which were from an average 2011 brood return that out-migrated to the ocean in 2012. It was quite apparent that other salmon species that also out-migrated in 2012 encountered improved survival conditions (i.e. Pink and Coho returns in 2013). One concern preseason was that the lower than average fish size of the 2011 brood would have some negative affect on the survival of the 4 year old returns in 2015. The preseason expectation for ISC Chum suggested near target returns to the area but was highly uncertain.

Table 3-4. Estimated harvest of summer Chum salmon by PFMA for commercial, research and test fishing vessels ¹ (July through the second week of September), 2006–2015.

| Year | Statistical Weeks | Pacific Fishery Management Area | | | | | Total |
|---------|-------------------|---------------------------------|----|----|----|-----------------|-------|
| | | 18 | 19 | 20 | 21 | 29 ² | |
| 2006 | 7/1–9/2 | 0 | 0 | 52 | 0 | 419 | 471 |
| 2007 | 7/1–9/2 | 0 | 0 | 0 | 0 | 49 | 49 |
| 2008 | 7/1–9/2 | 0 | 0 | 0 | 0 | 67 | 67 |
| 2009 | 7/1–9/2 | 0 | 0 | 0 | 0 | 27 | 27 |
| 2010 | 7/1–9/2 | 0 | 0 | 16 | 0 | 384 | 400 |
| 2011 | 7/1–9/2 | 0 | 0 | 0 | 0 | 883 | 883 |
| 2012 | 7/1–9/2 | 0 | 0 | 0 | 0 | 125 | 125 |
| 2013 | 7/1–9/2 | 0 | 0 | 0 | 0 | 61 | 61 |
| 2014 | 7/1–9/2 | 0 | 0 | 0 | 0 | 686 | 686 |
| 2015 | 7/1–9/2 | 0 | 0 | 0 | 0 | 1 | 1 |
| Average | | 0 | 0 | 7 | 0 | 270 | 277 |

(1) Does not include PSC test fisheries harvest.

(2) Majority of this harvest is fresh water based and likely of Fraser River origin.

The following sections describe the fisheries directed at fall Chum salmon in each of the Southern B.C. areas in 2015.

3.4.1 Johnstone Strait

The Johnstone Strait test-fishery provided timing and abundance information for the 2015 return, which are important in assessing the performance of the 20% fixed exploitation rate strategy. It also provided an index of abundance, used to determine the likelihood of the number of returning Chum salmon being over the 1.0 million critical threshold level (requirement for commercial openings). Catch per unit effort in the test fishery was higher than what was encountered in 2010 and it was determined that the ISC index of abundance was likely higher than the critical threshold required for continuation of commercial fisheries. All subsequent commercial openings in Johnstone Strait were prosecuted as planned. Age composition derived from the test-fishery and commercial samples was dominated by 4 year olds throughout the season. Escapements and catches in 2015 suggested returns were average to below average in many ISC Chum. Table 3-5 outlines the duration and Chum salmon harvest, of fishery openings during the 2015.

In 2015, there were two competitive commercial purse seine openings; three gill net openings in and one troll individual transferable quota (ITQ) fishery in Johnstone Straits (Area 12 and Area 13). The total 2015 commercial harvest for Johnstone Strait, including test fishery harvest, was 529,520 Chum salmon (Table 3-5). There were no directed commercial Chum salmon fisheries in the terminal areas of Johnstone Strait. Recreational harvest was low in Johnstone Strait (estimated at <2,000 Chum salmon; Table 3-5) with little monitoring during the month of October. First Nations FSC harvest was estimated at 22,472 in the Johnstone Strait area (Table 3-5). The total estimated harvest of Chum salmon in Johnstone Strait in 2015 was 553,569.

Table 3-5. Estimated Chum salmon harvest from Chum salmon-directed fisheries occurring in Johnstone Strait by PFMA, 2015.

| Gear Type | Area | Fishery Dates | | Fishery Duration (hours) | Estimated Harvest |
|----------------------------|-------|--------------------------------------|---------|--------------------------|-------------------|
| | | Start | End | | |
| Purse Seine (Test) | 12 | Sep. 15 | Oct. 29 | n/a | 36,679 |
| Purse Seine | 12 | Sockeye / Pink Directed ¹ | | -- | -- |
| | 13 | | | -- | -- |
| | 12 | Oct. 5 | Oct. 5 | 12 | 149,360 |
| | 13 | Oct. 5 | Oct. 5 | 12 | 89,055 |
| | 12 | Oct. 19 | Oct. 19 | 11 | 41,934 |
| | 13 | Oct. 19 | Oct. 19 | 11 | 72,153 |
| Gill Net | 12 | Oct. 2 | Oct. 4 | 41 | 29,408 |
| | 13 | Oct. 2 | Oct. 4 | 41 | 7,401 |
| | 12 | Oct. 16 | Oct. 18 | 41 | 21,802 |
| | 13 | Oct. 16 | Oct. 18 | 41 | 18,837 |
| | 12 | Oct. 25 | Oct. 27 | 41 | 4,967 |
| | 13 | Oct. 25 | Oct. 27 | 41 | 9,380 |
| Troll ² | 12 | Sep. 28 | Oct. 17 | n/a | 1,206 |
| | 13 | Sep. 28 | Oct. 17 | n/a | 40,369 |
| | 12 | Oct. 18 | Nov. 1 | n/a | 80 |
| | 13 | Oct. 18 | Nov. 1 | n/a | 6,889 |
| Recreational ³ | 11/12 | Jun. | Oct. | n/a | |
| | 13 | Sep. | Oct. | n/a | 1,577 |
| First Nations ³ | 12 | n/a | n/a | n/a | 7,565 |
| | 13 | n/a | n/a | n/a | 14,907 |
| Total | | | | | 553,569 |

(1) This is by-catch from Fraser Sockeye & Pink directed fisheries that took place during early September.

(2) The troll fishery was opened from Sep. 29 to Nov. 2. The fishery was an effort based quota over that time period. Statistical weeks 10/1 to 10/2 are grouped and weeks 10/3 to 11/1 are grouped.

(3) The recreational and First Nation fisheries time periods varied over the season.

3.4.2 Strait of Georgia

In 2015, the total return of Chum salmon to Area 14 systems was slightly below low end of the preseason forecasted range (280,000 and 420,000). Initially a small gillnet opening was initiated (Nov 2-4th for 12,812 Chum) but catches agreed with the below forecast returns and no further commercial opportunities occurred.

Escapements returning to the Nanaimo River in Area 17 indicated above goal returns and fisheries were initiated. Commercial gillnet fisheries took place from Oct 29th-Nov 18th harvesting a total of 18,133 Chum (Table 3-6). Commercial seine opening was also provided from Nov 5-18th harvesting a total of 176 Chum.

In-season assessments in Area 18 did identify sufficient escapement levels in the Cowichan River to initiate fisheries. Commercial Gillnet fisheries took place from Nov 7th harvesting a total of 22,117

Chum. (Table 3-6). Commercial Seine fisheries took place on Nov 11th harvesting a total of 182,857 Chum.

The recreational creel survey within the Strait of Georgia area was not conducted after September in 2015. There were 221 Chum estimated in the creel program in 2015. The harvest by First Nations in the Strait of Georgia in 2015 was estimated to be 6,275 Chum salmon (Table 3-6).

The total estimated harvest of Chum salmon in the Strait of Georgia in 2015 was 242,591.

Table 3-6. Estimated Chum salmon harvest from Chum salmon-directed fisheries occurring in the Strait of Georgia by PFMA, 2015.

| Gear Type | Area | Fishery Dates | | Fishery Duration (hours) | Estimated Harvest |
|----------------------------|------------------------------|---------------|---------|--------------------------|-------------------|
| | | Start | End | | |
| Purse Seine | 17 | Nov. 5 | Nov. 18 | 168 | 176 |
| | 18 | Nov. 11 | Nov. 11 | 12 | 182,857 |
| Gillnet | 14 | Nov. 2 | Nov. 4 | | 12,812 |
| | 17 | Oct. 29 | Oct. 30 | | 10,344 |
| | 17 | Nov. 4 | Nov. 18 | | 7,789 |
| | 18 | Nov. 7 | Nov. 7 | | 22,117 |
| Recreational ¹ | 14-19, 28 29 ³ | - | - | n/a | 221 |
| First Nations ² | 14 | n/a | n/a | n/a | - |
| | 15 | n/a | n/a | n/a | - |
| | 16 | n/a | n/a | n/a | - |
| | 17 | n/a | n/a | n/a | - |
| | 18 | n/a | n/a | n/a | 6,275 |
| | 19 | n/a | n/a | n/a | - |
| | 28 | n/a | n/a | n/a | - |
| Total | | | | | 242,591 |

(1) The recreational fishery was not monitored after September; harvest of Chum salmon was likely very low.

(2) The First Nation fisheries time periods varied over the season.

(3) Marine subareas of Area 29; i.e. does not include an estimate for Chum salmon harvest in the tidal portion of the lower Fraser River downstream of Mission BC (Appendix B-3).

3.4.3 Fraser River

Directed Chum salmon fisheries occur within the Fraser River for all fishing sectors, including First Nations, recreational, commercial, and test fisheries. In recent years, significant conservation measures have been implemented in-river during the Fraser River Chum salmon migration period in order to protect co-migrating stocks of concern, particularly Interior Fraser River Coho salmon and Interior Fraser River Steelhead. Depending on the fishery, these measures have included time and area closures, gear restrictions and release requirements.

The escapement objective for Fraser River Chum salmon is 800,000 spawners. As described in section 3.3.3 (*Fraser River Chum Salmon Management Strategy*), Chum salmon fishing opportunities are provided based on in-season assessments of run-size, using CPUE information from the CDFO Albion Chum salmon test fishery. The initial in-season estimate of terminal abundance provided to the US on

October 20, 2015 was 1.567 million Chum salmon. A subsequent in-season estimate of 1.780 million was provided on October 22.

Table 3-7. Estimated harvest of Chum salmon from salmon fisheries occurring in Area 29 (both marine and in-river) and in Region 2 (non-tidal) of the Fraser River, 2015.

| Fishing Sector | Fishery Description | Target Species | Fishery Dates | | Fishery Duration | Assessment Period ¹ | | Estimated Harvest |
|---------------------------------|-----------------------|----------------|---------------|--------|------------------|--------------------------------|--------|-------------------|
| | | | Start | End | | Start | End | |
| Test | Albion Chinook | Chinook | 26-Apr | 20-Oct | 153 days | n/a | n/a | 1,728 |
| | Albion Chum | Chum | 01-Sep | 23-Nov | 58 days | n/a | n/a | 8,603 |
| | Whonnock Sockeye | Sockeye | 22-Jul | 27-Sep | 68 days | n/a | n/a | 177 |
| | Cottonwood Sockeye | Sockeye | 13-Jul | 15-Sep | 65 days | n/a | n/a | 17 |
| Test Subtotal | | | | | | | | 10,525 |
| Commercial ² | Area E Gill Net | Chum | 23-Oct | 27-Oct | 2 days | n/a | n/a | 187,117 |
| | Area B Seine | Pink | 09-Sep | 10-Sep | 2 days | n/a | n/a | 0 |
| | | Chum | 27-Oct | 28-Oct | 2 days | n/a | n/a | 3,706 |
| Commercial Subtotal | | | | | | | | 190,855 |
| Recreational ^{3, 4, 5} | Fraser River Mainstem | Mixed | 03-Aug | 14-Aug | 12 days | 03-Aug | 14-Aug | 0 |
| | | Mixed | 29-Aug | 31-Dec | 125 days | 29-Aug | 30-Sep | 123 |
| | Chilliwack River | Mixed | 01-Jul | 31-Dec | 184 days | 21-Sep | 15-Nov | 2,466 |
| | Nicomen Slough | Mixed | 01-Jan | 31-Dec | 365 days | 09-Oct | 06-Nov | 45 |
| Recreational Subtotal | | | | | | | | 2,634 |
| First Nations ^{2, 6} | FSC | Mixed | 04-Apr | 30-Sep | n/a | n/a | n/a | 249 |
| | | Chum | 10-Oct | 22-Nov | n/a | n/a | n/a | 35,647 |
| | ESSR | Chum | n/a | n/a | n/a | n/a | n/a | 15,692 |
| | EO | Pink | 14-Sep | 25-Sep | 6 days | n/a | n/a | 0 |
| | | Chum | 22-Oct | 07-Nov | 17 days | n/a | n/a | 122,309 |
| First Nations Subtotal | | | | | | | | 173,897 |
| Total | | | | | | | | 377,911 |

(1) Assessment Period Start and End dates refer to the period of assessment for the recreational fisheries only.

(2) Fishery Start and End Dates for the commercial and First Nations fisheries reflect the first and last date the fisheries were permitted; the fisheries were not continuous between these dates.

(3) Fishery Start and End Dates for the recreational fisheries reflect the period the fishery was opened to the retention of Chum salmon; retention of other salmon species may have been allowed beyond these dates.

(4) The recreational estimates are for the periods of assessment; assessment periods may or may not coincide with Chum retention fishery opening and closing dates.

(5) Recreational estimates are preliminary.

(6) The Fishery Duration represents the maximum number of days provided to First Nations; individual First Nations may have fished for shorter periods.

First Nations FSC gill net fisheries targeting Chum salmon were initiated October 10 following a closure period to protect co-migrating Interior Fraser Coho salmon. These fisheries harvested a total of 35,647 Chum salmon. First Nations Economic Opportunity (EO) fisheries directed at Chum salmon in 2015 harvested 122,309 Chum (Table 3-7). In addition to these fisheries, certain First Nations groups were also provided access to Chum salmon that returned to hatchery facilities but that were not required for

broodstock. A total of 15,692 Chum salmon deemed to be surplus to spawning requirements (ESSR) were provided to First Nations from various hatchery facilities in 2015.

First Nations FSC Chinook and Sockeye-directed fisheries occurring in 2015 had minimal bycatch of Fraser Chum salmon, harvesting 249 Chum salmon (Table 3-7). Zero Chum were taken as bycatch in Pink-directed Economic Opportunity fisheries in 2015.

The total harvest of Chum salmon in all Fraser River First Nations fisheries (including hatchery ESSR fish) was 173,897 in 2015 (Table 3-7 and Table 3-8).

Table 3-8. Estimated harvest of Chum salmon from First Nations and commercial salmon fisheries occurring in Area 29 and Fraser River, 2006-2015.

| Year | First Nations | | | Commercial | | | Total |
|----------------|---------------|---------------|---------------|-------------------------------------|-------------------------------------|--|----------------|
| | FSC | EO | ESSR | Area E GN Area B SN Area H TR | Albion and PSC Test Fisheries | Other Scientific Licenses ¹ | |
| 2006 | 15,150 | 114,708 | 93,516 | 163,757 | 16,942 | n/a | 404,073 |
| 2007 | 13,344 | 77,490 | 29,884 | 30,400 | 7,650 | n/a | 158,768 |
| 2008 | 31,553 | 50,004 | 41,683 | 38,006 | 10,155 | n/a | 171,401 |
| 2009 | 12,991 | 68,150 | 8,458 | 42,116 | 9,249 | n/a | 140,964 |
| 2010 | 13,480 | 186 | 14,021 | 209 | 10,762 | n/a | 38,658 |
| 2011 | 22,331 | 4,886 | 50,867 | 36,058 | 3,553 | n/a | 117,695 |
| 2012 | 30,746 | 102,185 | 34,593 | 60,404 | 13,487 | n/a | 241,415 |
| 2013 | 37,435 | 107,959 | 43,176 | 101,890 | 12,270 | n/a | 302,730 |
| 2014 | 39,516 | 100,258 | 34,638 | 164,363 | 12,634 | n/a | 351,409 |
| 2015 | 35,896 | 122,309 | 15,692 | 190,855 | 10,525 | n/a | 375,277 |
| Average | 25,244 | 74,814 | 36,653 | 82,806 | 10,723 | n/a | 230,239 |

(1) Includes “experimental” selective fishing scientific licenses issued under the CDFO Selective Fishing Program.
(n/a) not applicable.

The major Fraser River watershed recreational salmon fisheries impacting Chum salmon in 2015 include significant fisheries occurring in the lower Fraser River mainstem, the Chilliwack River and the Stave River, and minor salmon fisheries occurring in the Harrison River and the Nicomen Slough/Norish Creek system. The latter four systems are tributaries to the Fraser River in the lower Fraser Valley. The recreational fisheries occurring in the Harrison and Stave rivers were not assessed in 2015.

The lower Fraser River mainstem was open to recreational angling and the retention of Chum salmon on two separate occasions in 2015. The initial opening occurred from August 3 to August 14, 2015 (daily retention limit of 2). This was followed by a closed period of 2 weeks before being re-opened to angling and the retention of Chum salmon from August 29 to December 31, 2015 (daily retention limit of 2). In 2015, this mainstem recreational fishery was assessed from August 3 to August 14 and August 29 to September 30, 2015; estimates of 123 and 111 Chum salmon were harvested and released, respectively (Table 3-7). The estimated harvest in 2015 was similar to the previous two years and notably less than in 2012 (Table 3-9). In part, this was due to the 2013, 2014 and 2015 assessments being truncated approximately two months earlier than 2012. In 2012, approximately 90% of the reported Chum salmon harvest in this fishery occurred from October 1 to November 30.

The Chilliwack River recreational fishery was open to the retention of Chum salmon from July 1 to December 31, 2015 (daily retention limit of 1). This fishery was assessed from September 21 to

November 15, 2015; estimates of 2,466 and 9,800 Chum salmon were harvested and released, respectively (Table 3-7). Similar to the lower Fraser River mainstem recreational fishery, the estimated Chum salmon harvest in 2015 for the Chilliwack fishery was equal in magnitude to the previous two years and notably less than in 2012 (Table 3-9); however, unlike the Fraser fishery, the Chilliwack fishery assessment was relatively consistent between 2012 and 2015.

The Harrison River, Stave River and Nicomen Slough/Norrish Creek recreational fisheries were open to the retention of Chum salmon year round (daily retention limit of 2). The Nicomen Slough/Norrish Creek fishery was assessed from October 9 to November 6, 2015. Estimates of 45 and 27 Chum salmon were harvested and released, respectively for this fishery (Table 3-7). Chum salmon catch in this fishery was similar to catch in 2014 and notably less than that estimated in 2013. This variation in catch may be explained, in part, by the 2015 assessment being truncated by 24 days in November. In 2013, approximately 85% of the Chum harvest and 90% of the Chum catch occurred from November 16 to 30. Recreational fisheries occurring in the Harrison and Stave rivers were not assessed in 2015.

In total, for assessed recreational fisheries occurring in the lower Fraser River watershed in 2015, estimates of 2,634 and 9,938 Chum salmon were harvested and released, respectively (Table 3-7 and Table 3-9).

Commercial fisheries in the lower Fraser River (i.e. Area E gill net) are closed annually from early September to mid-October during the Interior Fraser Coho migration period. In recent years, this closure has been extended into late October to provide additional protection to Interior Fraser Steelhead. As a result, commercial Area E gill net Chum salmon fisheries now typically occur during the last week of October and first week of November. In 2015, there were two 12-hour Area E commercial openings in the Fraser River on October 23 and October 27 that harvested a total of 187,117 Chum salmon.

The Area B seine fleet also mounted a limited-participation Chum salmon-directed fishery off the mouth of the Fraser River on October 27 that harvested 3,706 Chum salmon (Table 3-7).

Table 3-9. Estimated harvest of Chum salmon in assessed (i.e. surveyed) lower Fraser River recreational fisheries, 2006–2015.

| Year | Recreational Fishery | | | | | |
|----------------------|---------------------------|------------------|----------------|-------------|----------------|-------|
| | Fraser River ¹ | Chilliwack River | Nicomen Slough | Stave River | Harrison River | Total |
| 2006 | 900 | 5,375 | 2 | ns | ns | 6,277 |
| 2007 | 3,007 | 1,553 | 4 | ns | ns | 4,564 |
| 2008 * | 760 | 3,937 | 0 | ns | ns | 4,697 |
| 2009 * | 43 | 2,404 | 10 | ns | 742 | 3,199 |
| 2010 * | 1,549 | 1,142 | 10 | 3,578 | ns | 6,279 |
| 2011 * | 32 | 278 | 14 | ns | ns | 324 |
| 2012 * | 2,298 | 4,633 | 9 | ns | ns | 6,940 |
| 2013 * | 53 | 2,429 | 1,016 | ns | ns | 3,498 |
| 2014 * | 344 | 2,917 | 15 | ns | ns | 3,276 |
| 2015 * | 123 | 2,466 | 45 | ns | ns | 2,634 |
| Average ² | 911 | 2,713 | 113 | 3,578 | 742 | 4,169 |

(1) Mainstem portion of the lower Fraser River (downstream of Hope, B.C.; Appendix B).

(*) Preliminary estimates.

(ns) not surveyed.

In 2015, a total of 190,855 Chum salmon were harvested in commercial (excluding Economic Opportunity and test fisheries) fisheries in the Fraser River and the marine areas immediately off the Fraser River mouth (Table 3-7 and Table 3-8).

Four test fisheries operated in the Fraser River in 2015: the CDFO-operated Albion Chinook and Chum salmon gill net test fisheries and the PSC-operated Whonnock and Cottonwood Sockeye salmon gill net test fisheries. While Whonnock and Cottonwood test fisheries target Sockeye salmon, the Albion test fishery assesses abundance of both Chinook and Chum salmon, using different net configurations to target the two different species (section 3.3.3.1).

In 2015, the Albion Chinook salmon test fishery operated from April 26 through October 20, and the Albion Chum salmon test fishery operated from September 1 through November 23. From September 1 to October 20, the Chinook and Chum test fisheries operated on alternating days (section 3.3.3.1). The Albion test fishery did not operate on October 23 or October 27 due to gear conflicts with the Area E opening in the Fraser River. Chum salmon harvest in the Albion Chinook salmon test fishery, representing 153 fishing days, totaled 1,728 and harvest in the Albion Chum salmon test fishery, representing 58 days, totaled 8,603 Chum salmon (Table 3-7).

The PSC Whonnock Sockeye salmon test fishery operated for 68 days from July 22 through September 27, and harvested 177 Chum salmon. The PSC Cottonwood Sockeye salmon test fishery fished for 65 days from July 13 through September 15, harvesting only 17 Chum salmon (Table 3-7).

In total, an estimated 10,525 Chum salmon were harvested in the CDFO and PSC test fisheries occurring in the Fraser River in 2015 (Table 3-7 and Table 3-8).

3.4.4 West Coast Vancouver Island

The total catch of Chum in 2015 WCVI commercial fisheries was about 170,500. Most of that catch occurred in the Nitinat (Area 21/22) hatchery directed fishery. A limited entry commercial fishery in Nootka Sound (Area 25) caught about 1,200 Chum.

First Nations FSC harvest of Chum salmon occurred in all WCVI areas and totaled 2,246 for 2015.

3.4.4.1 Nitinat

The pre-season forecast was 150,000, which was below the lower reference point of 225,000 and resulted in a cautious approach to the management early in the season. In season information from the gillnet assessment fishery indicated abundance was well above forecast and fisheries were initiated. The return in 2015 was estimated at 502,000 Chum (Table 3-10), which was well above below the Lower Reference Point at which commercial fisheries can be conducted.

Gillnet and seine openings (Area 21) occurred for a total catch of 58,580 and 110,535 Chum respectively for a total commercial catch of 169,115 Chum. First Nations FSC (Area 21/22), ESSR (Area 22) harvest and hatchery brood stock of Chum salmon in 2015 totaled 93,000 Chum (Table 3-10).

Table 3-10. Nitinat area harvest, hatchery broodstock collection and spawning escapement estimates by PFMA, 2006–2015.

| Year | Area 21 | | Area 22 | |
|---------|-------------|---------------|----------------------------|------------------|
| | Seine Catch | Gillnet Catch | In-lake Catch & Broodstock | Natural Spawners |
| 2006 | 223,927 | 229,243 | 145,000 | 155,053 |
| 2007 | - | 180,111 | 22,268 | 115,789 |
| 2008 | 18,796 | 23,919 | 34,505 | 52,632 |
| 2009 | - | - | 32,434 | 54,000 |
| 2010 | - | - | 42,971 | 59,900 |
| 2011 | 65,469 | 211,968 | 136,641 | 248,655 |
| 2012 | 97 | 23,219 | 75,816 | 206,704 |
| 2013 | - | 15,730 | 25,800 | 25,066 |
| 2014 | - | - | 56,069 | 125,500 |
| 2015 | 58,580 | 110,535 | 93,177 | 240,000 |
| Average | 73,374 | 113,532 | 66,468 | 128,330 |

(1) includes Nitinat tributaries

3.5 Escapement

Chum salmon that escape First Nations, commercial (including test fisheries), and recreational fisheries, and any First Nations ESSR harvests, form the escapement totals of spawning Chum salmon for this section of the report. This *net escapement* can include Chum salmon that spawn in natural areas and constructed spawning channels, and those that are used as broodstock in enhancement facilities for hatchery production purposes. For the purposes of this report *spawning escapement* refers to salmon that spawn by their own volition in nature in either a natural (e.g. river) or manmade channel (e.g. controlled spawning channel), and does not include salmon used for broodstock.

Some river systems in southern B.C. support summer Chum salmon populations. However, because the summer aggregate is relatively minor when compared to the larger fall aggregate in southern B.C., and due to their distinctively early run timing (e.g. July to late August in Johnstone Strait) allowing for only a minimal bycatch in southern B.C. salmon fisheries (Table 3-4), summer Chum salmon escapement is not included in the escapement totals for this section of the report (i.e. escapement totals in this section refer to fall Chum salmon only).

3.5.1 Inside Southern

The stocks that are managed within the context of the ISC salmon plan are the fall Chum salmon. These Chum salmon enter Johnstone Strait during the September to November time period. Escapement estimates for these Chum salmon since 2006, are presented in Table 3-11.

Table 3-11. Inside Southern Chum salmon net escapement estimates (spawning escapement plus hatchery broodstock) for Fraser and Non-Fraser stock aggregates, 2006–2015¹.

| Year | Inside Southern Chum Escapement | | |
|---------|---------------------------------|------------------|-----------|
| | Fraser River | Non-Fraser River | Total |
| 2006 | 2,026,673 | 696,960 | 2,723,633 |
| 2007 | 1,026,701 | 902,149 | 1,928,850 |
| 2008 | 940,143 | 690,814 | 1,630,957 |
| 2009 | 619,363 | 853,501 | 1,472,864 |
| 2010 | 634,493 | 416,398 | 1,050,891 |
| 2011 | 1,084,652 | 912,293 | 1,996,945 |
| 2012 | 1,280,332 | 1,287,850 | 2,568,182 |
| 2013 | 799,573 | 1,224,131 | 2,023,704 |
| 2014 | 986,114 | 619,296 | 1,605,410 |
| 2015 | 808,219 | 659,847 | 1,468,066 |
| Average | 1,020,626 | 826,324 | 1,846,950 |

(1) Does not include Chum salmon which are surplus to spawning requirements (ESSR) and are removed from the spawning areas as they do not contribute to production in subsequent years. Those Chum salmon are assigned to harvest. Hatchery broodstock removals are included as they contribute to subsequent production.

The primary enhanced escapement areas for Inside Southern Chum salmon are presently limited to the mid-Vancouver Island, Squamish River, Burrard Inlet and Fraser River areas. The enhancement facilities in the mid-Vancouver Island consist of Big Qualicum, Little Qualicum and Puntledge rivers Hatcheries. There is one enhancement facility in the Burrard Inlet (Capilano River Hatchery: limited Chum salmon hatchery production), and the Squamish watershed (Tenderfoot Creek Hatchery) and four in the Fraser River watershed (Inch Creek, Chilliwack River, and Chehalis River Hatcheries and Weaver Creek Spawning Channel). Enhancement on the Capilano River is opportunistic, only using Chum salmon that return to the hatchery facility for broodstock.

3.5.2 West Coast Vancouver Island

In 2015, Nitinat Chum salmon spawning escapement was estimated at 240,000 (Table 3-10) which is a significant improvement compared to the 2014 return. In other WCVI Areas, Chum salmon spawning escapements varied from below average (Area 23 and 24) to above average (Area 25 and 26) in the areas monitored (Table 3-12). Spawning escapement estimates given for Areas 23-26 in Table 3-12 are derived from index system assessments in each Area. From historical surveys of Chum salmon streams in each Area, the proportion each stream contributed (on average) to the total spawning escapement for the Area is estimated. Spawning escapement estimates from index systems currently surveyed annually, are expanded to an Area spawning escapement to account for those streams not surveyed.

Table 3-12. WCVI Chum salmon spawning escapement estimates¹ by PFMA, 2006–2015.

| Year | WCVI Pacific Fishery Management Area | | | |
|---------|--------------------------------------|------------------------|----------------------------------|----------------------|
| | Area 23 (Barkley) | Area 24 (Clayoquot) | Area 25 ² (Nootka) | Area 26 (Kyuquot) |
| 2006 | 22,720 | 117,817 | 81,278 | 78,439 |
| 2007 | 58,012 | 82,272 | 43,178 | 91,197 |
| 2008 | 24,630 | 25,978 | 46,495 | 33,392 |
| 2009 | 27,920 | 35,392 | 82,138 | 32,532 |
| 2010 | 24,352 | 61,799 | 51,160 | 36,624 |
| 2011 | 121,692 | 149,164 | 91,989 | 125,989 |
| 2012 | 40,419 | 42,276 | 49,148 | 49,933 |
| 2013 | 35,302 | 44,174 | 69,877 | 79,812 |
| 2014 | 12,475 | 36,878 | 51,323 | 41,130 |
| 2015 | 32,559 | 33,299 | 108,154 | 82,277 |
| Average | 40,008 | 56,804 | 67,474 | 65,132 |

(1) Index system spawning escapements expanded to total spawning escapement for the Area.

(2) Nootka spawning escapements include hatchery river systems.

4 United States Chum Salmon

4.1 Washington Run Sizes, Spawning Escapements, and Catches

Tables 4-1 through 4-3 provide pre-season forecasts of run size, post-season estimates of run size and post-season estimates of spawning escapement for the Chum salmon runs returning to Puget Sound and coastal Washington areas. The tables present estimates for three major Chum salmon groupings, defined by their return timings.

Summer Chum Salmon

The Strait of Juan de Fuca summer Chum salmon post-season run size of 15,602 was 2.2 times greater than the pre-season forecast, as well as the 2006–2014 average post season run size. The Hood Canal summer Chum salmon run size of 36,055 was 60% larger than the pre-season forecast and 1.85 times larger than the 2006–2014 average post-season run size. Approximately 91% of the combined Strait of Juan de Fuca and Hood Canal summer Chum salmon run went to escapement. The summer Chum salmon pre-season forecast for South Puget Sound for 2015 was estimated at approximately 57,987 fish. The post-season run size estimate of South Puget Sound summer Chum salmon in 2015 was 32,570, which was 43% lower than the pre-season forecast and 42% lower than the 2006-2014 average post-season run size.

Table 4-1. Puget Sound summer Chum salmon pre-and post-season estimates of run size and spawning escapements, 2006–2015.

| Region | Type | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 |
|--------------------------------|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Strait of Juan de Fuca | Pre-Season | 8,238 | 8,566 | 5,969 | 5,198 | 3,991 | 5,308 | 5,915 | 6,603 | 9,040 | 7,105 |
| | Post-Season | 8,280 | 3,324 | 3,576 | 5,147 | 9,332 | 5,705 | 6,337 | 14,801 | 8,179 | 15,602 |
| | Escapement | 8,181 | 3,219 | 3,449 | 5,029 | 9,179 | 5,675 | 6,304 | 14,727 | 7,811 | 15,535 |
| Hood Canal | Pre-Season | 19,780 | 23,729 | 20,159 | 18,009 | 5,998 | 9,050 | 8,970 | 19,798 | 31,400 | 22,525 |
| | Post-Season | 29,590 | 12,838 | 17,618 | 9,200 | 12,957 | 7,170 | 31,134 | 24,325 | 30,408 | 36,055 |
| | Escapement | 26,418 | 10,539 | 15,112 | 7,236 | 12,533 | 6,914 | 29,855 | 22,618 | 27,396 | 32,644 |
| South Puget Sound ¹ | Pre-Season | 55,300 | 39,840 | 64,229 | 57,352 | 62,991 | 62,623 | 41,889 | 53,492 | 59,940 | 57,987 |
| | Post-Season | 78,797 | 57,786 | 32,065 | 25,010 | 49,098 | 39,823 | 38,936 | 38,332 | 25,317 | 32,570 |
| | Escapement | 61,405 | 52,668 | 20,872 | 21,660 | 43,739 | 38,515 | 24,521 | 35,444 | 20,593 | 32,521 |

(1) South Puget Sound estimates include only commercial harvest and escapement within Puget Sound.




Figure 4-1. Puget Sound summer Chum salmon abundance, 2006-2015, by region of origin (JDF: Strait of Juan de Fuca, HOOD: Hood Canal, SPS: South Puget Sound).

Fall Chum and Winter Salmon

The Puget Sound fall Chum salmon post-season assessment total was 1,450,298 fish, or 23% larger than the pre-season forecast and approximately 8% less than the 2006–2014 average run size. Regional post-season numbers varied in their deviation from the forecasted numbers, ranging from two times larger than the pre-season forecast (Skagit) to 91% below the pre-season forecast (Stillaguamish/Snohomish) (Table 4-2, Figure 4-2). The Washington coastal (Willapa Bay and Grays Harbor) fall Chum salmon

post-season assessment total was 96,805 fish, or approximately 41% above the pre-season forecast, and 2.3 times above the 2006-2014 average run size.




Figure 4-2. Puget Sound fall Chum salmon abundance, 2006-2015, by region of origin (JDF: Strait of Juan de Fuca, NOOK: Nooksack-Samish, SKAG: Skagit, STSNO: Stillaguamish/Snohomish, SPS: South Puget Sound, HOOD: Hood Canal).

The Puget Sound winter Chum salmon 2015 post-season run size assessment was 20,043, which was approximately 65% below the pre-season forecast and 72% below the 2006-2014 average run size (Table 4-3, Figure 4-3).




Figure 4- 1. South Puget Sound winter Chum salmon abundance, 2006-2015.

Table 4-2. Washington fall Chum salmon pre-and post-season estimates of run size and spawning escapements, 2006–2015.

| Region | Type | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 |
|-----------------------------|-------------|---------|-----------|---------|---------|---------|---------|---------|-----------|---------|---------|
| Willapa Bay | Pre-Season | 38,754 | 23,798 | 40,022 | 7,139 | 33,442 | 22,254 | 28,273 | 35,584 | 55,378 | 39,994 |
| | Post-Season | 23,486 | 17,668 | 12,989 | 17,444 | 26,701 | 69,802 | 43,069 | 27,564 | 30,276 | 48,756 |
| | Escapement | 15,250 | 17,392 | 9,511 | 12,743 | 25,204 | 66,871 | 26,640 | 24,979 | 26,508 | 45,409 |
| Grays Harbor | Pre-Season | 29,370 | 12,000 | 9,903 | 17,665 | 18,492 | 33,669 | 24,603 | 30,716 | 44,670 | 28,852 |
| | Post-Season | 15,786 | 12,136 | 8,879 | 19,776 | 43,582 | 50,494 | 40,561 | 41,524 | 28,728 | 48,049 |
| | Escapement | 11,268 | 11,342 | 6,249 | 15,216 | 34,644 | 30,101 | 27,876 | 22,519 | 15,378 | 34,587 |
| Strait of Juan de Fuca | Pre-Season | 3,830 | 2,143 | 2,745 | 2,587 | 2,222 | 1,618 | 1,203 | 813 | 4,369 | 9,305 |
| | Post-Season | 1,567 | 769 | 842 | 1,462 | 1,849 | 4,839 | 3,174 | 1,468 | 2,045 | 3,076 |
| | Escapement | 1,313 | 503 | 454 | 1,305 | 1,343 | 4,707 | 2,973 | 1,394 | 1,952 | 2,958 |
| Nooksack/ Samish | Pre-Season | 135,100 | 19,414 | 130,070 | 69,145 | 61,827 | 31,924 | 34,739 | 52,585 | 59,908 | 180,759 |
| | Post-Season | 66,610 | 48,632 | 27,471 | 32,007 | 48,892 | 152,176 | 39,100 | 68,176 | 95,440 | 112,704 |
| | Escapement | 29,289 | 21,572 | 16,257 | 26,268 | 30,510 | 100,748 | 21,703 | 48,506 | 60,058 | 68,690 |
| Skagit | Pre-Season | 164,094 | 90,481 | 132,036 | 26,828 | 50,226 | 26,834 | 59,167 | 15,325 | 16,505 | 47,292 |
| | Post-Season | 213,490 | 37,627 | 29,946 | 29,479 | 46,503 | 15,850 | 41,601 | 9,786 | 65,734 | 8,155 |
| | Escapement | 105,239 | 19,576 | 22,067 | 26,744 | 39,884 | 15,494 | 36,601 | 8,554 | 63,013 | 7,964 |
| Stillaguamish/ Snohomish | Pre-Season | 445,800 | 287,993 | 229,251 | 92,477 | 90,660 | 71,277 | 86,598 | 48,884 | 87,476 | 168,104 |
| | Post-Season | 479,987 | 235,799 | 125,064 | 37,689 | 110,050 | 61,598 | 40,003 | 35,145 | 64,272 | 13,669 |
| | Escapement | 272,925 | 43,664 | 38,553 | 29,378 | 46,628 | 60,792 | 35,450 | 30,317 | 51,599 | 8,169 |
| South Puget Sound | Pre-Season | 466,700 | 408,040 | 686,511 | 348,333 | 650,986 | 487,514 | 323,928 | 349,623 | 465,970 | 377,032 |
| | Post-Season | 939,638 | 1,036,772 | 464,193 | 344,854 | 627,131 | 548,484 | 611,657 | 597,762 | 589,869 | 503,531 |
| | Escapement | 369,035 | 368,848 | 152,549 | 150,732 | 201,992 | 179,094 | 190,817 | 186,457 | 200,660 | 160,260 |
| Hood Canal | Pre-Season | 668,400 | 587,155 | 668,397 | 374,929 | 477,409 | 352,019 | 426,675 | 323,597 | 442,308 | 394,246 |
| | Post-Season | 745,675 | 677,761 | 553,122 | 446,476 | 419,094 | 610,507 | 684,643 | 1,427,350 | 691,862 | 809,162 |
| | Escapement | 185,874 | 162,425 | 71,925 | 55,879 | 49,486 | 102,374 | 91,940 | 240,169 | 105,192 | 146,004 |

Table 4-3. Washington winter Chum salmon pre- and post-season estimates of run size and spawning escapements, 2006–2015.

| Region | Type | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 |
|-------------------------|-------------|---------|---------|--------|--------|---------|---------|--------|--------|--------|--------|
| South Puget Sound | Pre-Season | 149,200 | 81,065 | 98,922 | 83,380 | 89,293 | 55,923 | 51,599 | 61,113 | 84,263 | 56,709 |
| | Post-Season | 75,833 | 105,933 | 89,271 | 35,544 | 102,235 | 121,079 | 25,522 | 34,839 | 59,672 | 20,043 |
| | Escapement | 58,786 | 73,442 | 45,162 | 15,486 | 79,550 | 74,110 | 18,987 | 21,754 | 46,587 | 19,600 |

Chum Harvest

Paragraph 3 of the Chapter 6, Annex IV of the Treaty requires that Canada and the United States assess catch levels of summer Chum salmon caught during the August 1 through September 15 time period in boundary area fisheries. Table 4-4 provides Chum salmon harvest during this summer Chum salmon accounting period for U.S. boundary area fisheries. Table 4-5 provides Chum salmon harvest information from the Strait of Juan de Fuca (SJF) fisheries in Areas 4B, 5, and 6C and the San Juan Islands/Point Roberts (SJI/PR) fisheries in Areas 7 and 7A (Appendix B-1). Table 4-5 also includes annual Chum salmon harvest totals for the Puget Sound and the Washington Coastal areas.

Table 4-4. Harvest of Chum salmon in the Strait of Juan de Fuca (SJF) and the San Juan Islands (SJI) commercial fisheries during the HC/JDF summer Chum salmon accounting period (July 1–September 15) in years 2006–2015.

| Year | Time Period ¹ | | | | | Yearly Total |
|------------------------|--------------------------|-----------|----------|---------|----------|--------------|
| | 7/1–8/18 | 8/19–8/25 | 8/26–9/1 | 9/2–9/8 | 9/9–9/15 | |
| 95–97 GSI ² | 0.68 | 0.4 | 0.45 | 0.14 | 0.07 | |
| 2006 | 98 | 58 | 10 | 28 | 3 | 197 |
| 2007 | 0 | 0 | 0 | 0 | 9 | 9 |
| 2008 | 237 | 1 | 0 | 0 | 29 | 267 |
| 2009 | 4 | 0 | 1 | 14 | 25 | 44 |
| 2010 | 64 | 107 | 48 | 44 | 26 | 289 |
| 2011 | 29 | 0 | 3 | 118 | 30 | 180 |
| 2012 | 120 | 0 | 0 | 0 | 0 | 120 |
| 2013 | 44 | 21 | 123 | 131 | 556 | 875 |
| 2014 | 49 | 27 | 14 | 24 | 215 | 329 |
| 2015 | 97 | 0 | 27 | 0 | 0 | 124 |
| Averages | 74 | 21 | 23 | 36 | 89 | 243 |

(1) Indicates catch of Chum salmon during the Hood Canal/Juan de Fuca summer Chum period.

(2) Proportion of HC/JDF summer Chum in U.S. Areas 4b, 5, 6C, and 7 catch during the indicated periods, based on GSI proportions from 1995–1997.

Table 4-5. Harvest of summer, fall, and winter Chum salmon in SJF, SJI/PR, Puget Sound and Washington coastal areas, 2006–2015 ¹.

| Year | Region | | | |
|----------|--------------------------|-------------------------|--------------------------|-----------------------|
| | SJF (Areas 4B, 5, 6C) | SJI/PR (Areas 7, 7A) | Puget Sound ² | WA Coast ³ |
| 2006 | 4,483 | 106,634 | 1,568,576 | 13,532 |
| 2007 | 6,705 | 27,397 | 1,444,209 | 2,728 |
| 2008 | 5,695 | 75,374 | 961,036 | 7,067 |
| 2009 | 746 | 24,244 | 637,481 | 10,351 |
| 2010 | 1,941 | 23,652 | 949,870 | 12,593 |
| 2011 | 1,930 | 70,363 | 1,009,490 | 28,251 |
| 2012 | 637 | 73,255 | 1,083,578 | 30,697 |
| 2013 | 1,242 | 80,509 | 1,667,244 | 23,060 |
| 2014 | 5,253 | 147,022 | 1,018,649 | 18,069 |
| 2015 | 6,857 | 124,774 | 1,046,797 | 20,911 |
| Averages | 3,597 | 75,322 | 1,138,693 | 16,726 |

(1) Includes recreational harvest: 2015 recreational harvest estimates are preliminary.

(2) All other Puget Sound freshwater and marine harvest areas, except Strait of Juan de Fuca or San Juan Islands/Point Roberts Fisheries.

(3) WA Coast combines harvest from Areas 1–4 including Grays Harbor, Willapa Bay, Columbia River and Coastal Rivers.

4.2 *United States Strait of Juan de Fuca Fisheries (Areas 4B, 5, 6C)*

4.2.1 Management Intent

During the 2015 season, the management strategy for the Strait of Juan de Fuca (SJF) fishery consisted of maintaining the limited effort nature of this fishery and considering the needs of Puget Sound stocks of Chum salmon when making management decisions. This fishery has experienced minimal effort in recent years, resulting in relatively low harvest levels over the period of 2006–2015 (Table 4-5). Harvest and effort for the period of 2006–2015 were generally below levels observed in the late 1980s and early 1990s.

4.2.2 Fishery Description

As in previous years, the fall Chum salmon fishery in Areas 4B, 5 and 6C was restricted to Treaty Indian fishers using gill nets. The Treaty Chum salmon fishery opened the week of October 11, 2015 with a schedule of six days per week and continued through November 14, 2015. A total of 6,342 Chum salmon was harvested in this fishery, and there was a reported bycatch of 65 Coho salmon; 0 Steelhead; and 1 Chinook salmon.

In Areas 4B, 5, and 6C, the incidental harvests of Chum salmon prior to the Chum salmon-directed fishing season was slightly below the recent ten year average. A total of 95 Chum salmon, including those harvested in SJF test fisheries, were recorded during the summer Chum

salmon accounting period (July 1 – September 15). During the subsequent Coho salmon-directed fishery, 420 Chum salmon were harvested.

The total 2015 Chum salmon catch by all gears (including recreational) in the SJF was 6,857 Chum salmon (Table 4-5).

4.3 San Juan Islands / Point Roberts Fisheries (Areas 7 and 7A)

4.3.1 Management Intent

Chum salmon fisheries in Areas 7 and 7A are regulated to comply with a base harvest ceiling of 130,000 Chum salmon, unless a critically low level of abundance (<1,000,000) is identified for those populations migrating through Johnstone Strait (Inside Chum salmon). Chapter 6, Annex IV, specifies that Chum salmon-directed fishing is not allowed in Areas 7 and 7A before October 10. Paragraph 10 (a-b) specifies that for run sizes below the critical threshold, harvest of Chum salmon in Areas 7 and 7A will be limited to those taken incidentally to other species and in other minor fisheries, and shall not exceed 20,000.

4.3.2 Fishery Description

An estimated Fraser River Chum salmon run size of 1,567,000 was provided by Canada on October 21.

From October 1 through October 22, non-Tribal reef nets in Areas 7 and 7A were open daily with Chum salmon retention allowed. The reef net fishery was reopened from October 25 through November 7, but there was no effort during this period. Chum salmon catch in this fishery, between October 1 and November 7, was 5,851 fish (Table 4-6). Although the reef net fishery was re-opened, there was no reef net fishing effort after October 22.

Tribal and non-Tribal Chum salmon fisheries in Areas 7 and 7A for gill nets and purse seines began on October 10 and continued through October 23, with weekly catch monitoring. The U.S. catch for these gear types was 118,889 Chum salmon (Table 4-6). The Non-Treaty gillnet and purse seine fleets were open daily October 12, 13, 15 and then continuously from October 19 through October 22. The Treaty Indian gillnet and purse seine fisheries were opened on October 10 and ran continuously through October 23.

Table 4-6. Harvest of Chum Salmon in U.S. Areas 6, 7 and 7A Chum fisheries during 2015.

| Catch Area | Gill Net | Purse Seine | Reef Net |
|------------|----------|-------------|----------|
| 6 | 475 | | |
| 7 | 3,145 | 86,690 | 5,851 |
| 7A | 13,439 | 15,615 | |
| Total | 17,059 | 102,305 | 5,851 |

In the previous year, 2014, U.S. Chum salmon catches in Areas 7 and 7A had exceeded the U.S. catch ceiling by 16,571 Chum salmon (this value is updated from the previous annual report). Paragraph 10(h) of the Chum annex stipulates that *...catches in excess of 135,000 Chum salmon shall result in an overage being calculated by subtracting 130,000 from the total Chum catch.*

Overages will be accounted for by reducing the U.S. annual catch ceiling in up to two subsequent non-critical Inside Southern Chum salmon years. During 2015, the U.S. harvested 5,260 fewer Chum salmon in Areas 7 and 7A than the U.S. catch ceiling, thereby paying back in the first year 31.7% of the 2014 overage.

There were 34 Chum salmon reported as bycatch in Areas 7 and 7A during Fraser Panel approved Sockeye salmon fisheries in August and September. An additional 475 Chum salmon were harvested in U.S. commercial fisheries in Area 6, and 86 Chum salmon in recreational fisheries during the October 1 to October 31 time period (Table 4-6). Area 6 is not included in the PST Chum annex and the catch in this area does not count against the U.S. Chum salmon catch ceiling.

The total 2015 Chum salmon catch by all gears (including recreational) in Areas 6, 7 and 7A for the October 1 to November 7 time period was 125,215. Catch distribution, between Areas 6, 7 and 7A was 0.4%, 76.4% and 23.2%, respectively. During the fall Chum salmon fisheries in Areas 6, 7, and 7A, there was a reported bycatch of 19 Chinook and 3,510 Coho salmon and one Steelhead (Table 4-7).

Table 4-7. Bycatch of Chinook salmon, Coho salmon and Steelhead during U.S. commercial fisheries in Areas 6, 7 and 7A Chum salmon fisheries during 2015.

| Catch Area | Chinook | Coho | Steelhead |
|------------|---------|-------|-----------|
| 6 | | 415 | |
| 7 | 16 | 2,911 | 1 |
| 7A | 3 | 184 | |
| Total | 19 | 3,510 | 1 |

5 Stock Identification

5.1 Tagging of Adult Chum Salmon

No adult tagging projects were reported by either Canada or the United States in marine or nearshore areas in 2015.

5.2 Otolith, Fin Marking and Coded Wire Tagging (CWT)

5.2.1 Canada

Thermal marking of otoliths (Volk et al. 1987) has been used to mass mark hatchery raised salmonids in B.C. since the early 1990's (Hargreaves et al. 2001). The method involves manipulating the temperature of the rearing water by at least 2°C to induce lighter or darker deposition (density) of otolith layers. By systematically altering water temperature over a period of time, banding patterns (i.e. a "mark") unique to each hatchery can be created.

Thermal marking was first used on Chum salmon at Nitinat Hatchery on the West Coast of Vancouver Island in 1993 and continues to be used as a means of estimating hatchery contribution to both fisheries and escapement. Thermal marking is currently the only mark being applied to Chum salmon from Nitinat hatchery (Table 5-1). Transplanting thermally-marked

Chum salmon into Klanawa River started with the 2001 brood and ended with the 2006 broodyear (Table 5-1).

Table 5-1. Releases of Chum salmon with thermally-marked otoliths from WCVI based Canadian hatchery facilities, 2006–2015

| Broodyear | Nitinat Hatchery Release Sites | | Conuma Hatchery Release Sites | | | | |
|-----------|--------------------------------|---------------|-------------------------------|----------------|---------------|--------------|--------------|
| | Nitinat River & Lake | Klanawa River | Conuma River | Conuma Estuary | Tlupana River | Sucowa River | Canton River |
| 2006 | 26,323,943 | 2,585,428 | 0 | 1,455,119 | 1,139,028 | 181,333 | 133,731 |
| 2007 | 13,004,189 | 0 | 0 | 271,820 | 109,922 | 15,701 | 47,846 |
| 2008 | 7,631,058 | 0 | 0 | 1,011,562 | 254,905 | 292,430 | 418,655 |
| 2009 | 5,252,749 | 0 | 18,847 | 1,814,475 | 685,465 | 551,365 | 854,548 |
| 2010 | 14,182,582 | 0 | 0 | 1,794,205 | 1,015,355 | 311,828 | 646,203 |
| 2011 | 25,303,286 | 0 | 0 | 1,270,286 | 37,241 | 223,882 | 365,682 |
| 2012 | 15,813,031 | 0 | 0 | 785,682 | 813,957 | 41,196 | 116,757 |
| 2013 | 9,738,442 | 0 | 0 | 1,705,633 | 820,612 | 269,435 | 1,009,478 |
| 2014 | 17,767,521 | 0 | 410,214 | 1,467,870 | 0 | 0 | 200,646 |
| 2015 | 18,236,300 | 0 | 534,334 | 723,867 | 452,375 | 0 | 576,649 |

A second hatchery on the West Coast of Vancouver Island, Conuma Hatchery, first began thermally marking otoliths for several populations of Chum salmon beginning with the 1998 broodyear. This continued in 2015 with extensive releases of thermally-marked Chum salmon in various river systems of Nootka Sound (Table 5-1). Similar to Nitinat releases, comprehensive adult sampling programs aimed at detecting (i.e. recapturing) the thermally-marked Chum salmon have occurred during the fall Chum salmon fisheries. Information from these programs has permitted a better understanding of timing and catch distribution for the Chum salmon populations in Nootka Sound.

Table 5-2. Releases of marked Chum salmon from southern B.C. based Canadian hatchery facilities, 2006 to 2015

| Population | Broodyear | Fin Clip Type ¹ | Clipped | Poor Clips + Unclipped | Total |
|--------------------|-----------|----------------------------|---------|------------------------|---------|
| Big Qualicum River | 2006 | ADRV | 252,016 | 6,853 | 258,869 |
| | 2007 | ADRV | 251,329 | 9,471 | 260,800 |
| | 2008 | ADRV | 204,801 | 5,409 | 210,210 |
| | 2009 | ADRV | 250,692 | 8,381 | 259,073 |
| | 2010 | ADRV | 201,825 | 5,463 | 207,288 |
| | 2011 | AD | 250,337 | 4,011 | 254,348 |
| | 2012 | AD | 249,655 | 3,636 | 253,291 |
| | 2013 | AD | 251,108 | 15,932 | 267,040 |
| | 2014 | AD | 191,823 | 7,248 | 199,071 |
| | 2015 | AD | 250,646 | 4,423 | 255,069 |

(1) Clip Types: AD-adipose fin; LV-left ventral fin; RV-right ventral fin.

Not all release groups were represented by a mark. Contributions to fisheries and escapement for those groups were estimated by associating them with a marked release group having a similar size and release timing. Table 5-2 provides a summary of mark/tag combinations applied from 2006-2015 broodyears. Big Qualicum River is the only population that had any fin-clipped Chum salmon releases in 2015 (Table 5-2).

There were no coded wire tagged Chum released from Canadian facilities during the 2006-2015 time period.

5.2.2 United States

Otolith marking (Volk et al. 1987) was used in several hatchery supplementation and re-introduction programs for summer Chum salmon in the Hood Canal and Strait of Juan de Fuca regions (Table 5-3), and for fall Chum salmon in Puget Sound and the Lower Columbia River (Table 5-4). Otolith marking was used as a tool to assess program success and determine proportions of hatchery- and natural-origin fish (and identify hatchery strays) on spawning grounds.

Table 5-3. Numbers of Hood Canal summer Chum salmon released with thermally-marked otoliths or an adipose fin clip by broodyear, 2006–2015 (Point No Point Treaty Tribes and Washington Department of Fish and Wildlife 2015).

| Hood Canal Summer Chum Salmon | | | | |
|-------------------------------|-----------------|----------------|-----------|---------|
| Broodyear | Jimmycomelately | Hamma Hamma | Lilliwaup | Tahuya |
| 2006 | 79,428 | 151,550 | 88,800 | 133,826 |
| 2007 | 73,840 | 48,530 | 0 | 53,632 |
| 2008 | 88,766 | 208,450 | 68,810 | 97,142 |
| 2009 | 92,200 | 0 | 140,210 | 69,711 |
| 2010 | 85,630 | 0 | 139,816 | 27,706 |
| 2011 | 0 | 0 | 41,006 | 19,600 |
| 2012 | 0 | 0 | 157,760 | 110,000 |
| 2013 | 0 | 0 | 169,440 | 52,778 |
| 2014 | 0 | 0 | 109,680 | 120,000 |
| 2015 | 0 | 0 | 54,930 | 0 |

Table 5-4. Numbers of Puget Sound and Lower Columbia River fall Chum salmon released with thermally-marked otoliths or an adipose fin clip by broodyear, 2006–2015.

| Broodyear | Puget Sound | | | | | Lower Columbia River | |
|-----------|----------------|----------------|------------------|------------------|------------------|---|-----------------------|
| | Elwha River | Tulalip Bay | Kendall Creek | Terrell Creek | Whatcom Creek | Grays River Hatchery ¹ | Washougal Hatchery |
| 2006 | 0 | 0 | 0 | 0 | 0 | 129,427 | 0 |
| 2007 | 0 | 0 | 0 | 0 | 450,000 | 77,609 | 0 |
| 2008 | 0 | 0 | 0 | 0 | 400,000 | 104,600 | 0 |
| 2009 | 0 | 0 | 0 | 0 | 0 | 300,000 | 70,000 |
| 2010 | 0 | 0 | 0 | 0 | 2,000,000 | 250,000 | 57,455 |
| 2011 | 0 | 0 | 0 | 0 | 1,360,000 | 206,000 | 74,893 |
| 2012 | 0 | 0 | 0 | 0 | 1,547,900 | 157,800 | 58,004 |
| 2013 | 450,000 | 40,000 | 650,895 | 85,000 | 3,048,000 | 151,567 | 46,083 |
| 2014 | 450,000 | 8,000,000 | 1,094,383 | 1,000,000 | 2,283,000 | 192,156 | 400,000 |
| 2015 | 450,000 | 8,000,000 | 1,000,000 | 0 | 2,000,000 | 250,000 | 500,000 |

(1) Includes Grays River broodstock released in Big Creek, OR.

5.3 Genetic Stock Identification

In 2010, the Committee completed a joint recommendation to the Southern Panel concerning the type of stock identification methodology to be used for immediate and future application of

Canada and United States (U.S.) Chum salmon stock assessment (Appendix C). The Pacific Salmon Treaty (Annex IV, Chapter 6, Section 2) states:

“When the Parties provide stock composition information for fisheries, the Committee shall evaluate and report its conclusions using bilaterally agreed upon methods.”

Fisheries of interest include Johnstone Strait, Qualicum River, Fraser River, and Nitinat in Canada and Point Roberts, San Juan Islands and the Strait of Juan de Fuca in the U.S.

5.3.1 Fishery Sample Collection for Genetic Stock Identification

5.3.1.1 *Canada*

GSI samples were collected during the Johnstone Straits purse seine fisheries (commercial and test fishery) in 2015 (Table 5-5). In 2015, these samples were analysed for microsatellite markers using the first year of four years of funding supported by the Southern Endowment Funds (Araujo et al. 2016).

Table 5-5. Chum salmon tissue samples collected in Johnstone Strait fisheries and analyzed for GSI, 2006–2015.

| Year | Commercial Samples | | Test Fishery Samples | |
|------|--------------------|----------|----------------------|----------|
| | Collected | Analysed | Collected | Analysed |
| 2006 | 716 | 714 | 0 | 0 |
| 2007 | 879 | 869 | 0 | 0 |
| 2008 | 995 | 868 | 830 | 449 |
| 2009 | 1,418 | 1,101 | 999 | 0 |
| 2010 | 400 | 400 | 1,015 | 1,114 |
| 2011 | 865 | 863 | 1,404 | 905 |
| 2012 | 987 | 935 | 1,275 | 1,261 |
| 2013 | 1,185 | 973 | 1,311 | 1,291 |
| 2014 | 1,087 | 990 | 1,369 | 1,342 |
| 2015 | 1,348 | 982 | 1,395 | 1,326 |

5.3.1.2 *United States*

Among focused studies, WDFW used GSI to distinguish supplementation-origin and natural-origin adult summer Chum salmon in Hood Canal as part of the effort to monitor the Hood Canal summer Chum salmon recovery program. For instance, from 2006-2009 mSAT DNA data were used to clarify uncertain hatchery assignments when otoliths were broken or thermal marks were ambiguous. Each supplementation program for summer Chum salmon had a unique otolith mark and spawners could be identified as supplementation-origin or natural-origin, but not identified to their program of origin if their otolith mark was ambiguous.

These ambiguous supplementation program fish were genotyped and identified by assigning them to populations in the Hood Canal summer Chum salmon genetic baseline. Genetic analysis

was used to evaluate impacts of supplementation on the Hood Canal summer Chum salmon metapopulation (Small et al. 2009). WDFW also conducted juvenile stock assignment studies in the Strait of Juan de Fuca and Hood Canal during 2009 to 2015. The juvenile studies documented emigration timing and relative production of summer and fall Chum salmon in key populations. Further, 732 tissue samples were collected near Apple Cove Point in Puget Sound Area 9 in the eleventh year of Chum salmon DNA collection from this test fishery and an additional 357 tissues were collected in Area 9.

Fisheries in Area 7 and Area 7a were sampled and tissues analyzed (Table 5-6) from the fourth year of Southern Endowment Funding project using microsatellites (Araujo et al. 2016).

Table 5-6. Chum salmon tissue samples from Areas 7 and 7a analyzed for GSI in 2007-2015.

| Year | Area 7 | Area 7A |
|------|--------|---------|
| 2007 | 675 | 700 |
| 2008 | 711 | 651 |
| 2012 | 1,328 | 829 |
| 2013 | 419 | 423 |
| 2014 | 121 | 285 |
| 2015 | 192 | 204 |

5.3.2 Baseline Collection for Genetic Stock Identification

5.3.2.2 *Canada*

In 2015, CDFO continued to collect baseline GSI tissue samples of Southern British Columbia Chum salmon populations (Table 5-7).

Table 5-7. Fall Chum salmon GSI tissue collections from Southern B.C. by Conservation Unit, collection site, life stage, and influence¹ of enhancement, 2015.

| Conservation Unit | Collection Site | Life Stage | Enhancement Level ¹ | Sample Size |
|--------------------------|-----------------|------------|--------------------------------|-------------|
| Lower Fraser | Blaney_Cr | Adult | Natural | 66 |
| Lower Fraser | Chilqua_Cr | Adult | Natural | 1 |
| Lower Fraser | Silverhope_Cr | Adult | Natural | 242 |
| Lower Fraser | Hicks_Cr | Adult | Natural | 195 |
| Lower Fraser | Hopedale_Slough | Adult | Natural | 77 |
| Fraser Canyon | Hunter Cr | Adult | Natural | 43 |
| Fraser Canyon | Kawkawa Cr | Adult | Natural | 45 |
| Lower Fraser | Mcintyre Cr | Adult | Natural | 67 |
| Georgia Strait | Okeover | Adult | Natural | 223 |
| Lower Fraser | Railroad_Cr | Adult | Natural | 8 |
| Lower Fraser | Street Cr | Adult | Natural | 27 |
| Georgia Strait | Theodosia R | Adult | Natural | 148 |
| Lower Fraser | Whonnock R | Adult | Natural | 206 |
| Lower Fraser | Worth_Cr | Adult | Natural | 74 |
| Lower Fraser | Big Silver Cr | Adult | Natural | 1 |
| Lower Fraser | Cogburn Cr | Adult | Natural | 1 |
| Lower Fraser | Siddle Cr | Adult | Natural | 2 |
| Lower Fraser | Worth Cr* | Adult | Natural | 108 |
| Lower Fraser | Peach Cr | Adult | Natural | 163 |
| Lower Fraser | Widgeon | Adult | Natural | 135 |
| Loughborough | Phillips R | Adult | Natural | 37 |
| Howe Sound-Burrard Inlet | Cheakamus R | Adult | Natural | 100 |
| Total | | | | 1,969 |

- (1) Enhancement level refers to the degree to which the CDFO Committee members believed hatchery production contributed to the returns to the collection site. Based partly on the enhancement history in the CDFO hatchery production database, enhancement level is subjectively classified as either *hatchery* (insignificant contribution of natural production to returns), *mixed* (both enhanced and natural production contribute significantly to returns) or *natural* (insignificant contribution of hatchery production to returns).

5.3.2.1 *United States*

During 2015, WDFW and Washington Treaty Tribes continued baseline sampling of Washington State populations of Chum salmon (Table 5-8). Tissues were archived for future genotyping and will be added to the genetic baseline.

Table 5-8. Chum salmon GSI tissue collections from WA State, 2015.

| Year | Location | Tributary and Collection Name | N | WDFW code |
|------|---------------------|-------------------------------|----|-----------|
| 2015 | Central Puget Sound | Chico Creek | 66 | 15TJ |

REFERENCES CITED

- Araujo, A., P. Van Will, J. Candy and J. Zischke. 2016. Joint US and CA mixed stock Chum fisheries sampling and design and analysis 2015. Report to the Southern Fund Project 57536.
- Beattie, W., G. Kirby, L. LeClair. 1996. The Stock Composition of 1995 Chum Salmon Fisheries in Washington: Strait of Juan de Fuca (Catch area 5) and Northern Puget Sound (Catch Areas 7&7a). Northwest Indian Fisheries Commission and WDFW Joint document.
- CDFO [Canadian Department of Fisheries and Oceans]. 2015. Pacific Region Integrated Fisheries Management Plan, Salmon, Southern B.C., June 1, 2015 to May 31, 2016. Fisheries Management Branch, Fisheries and Oceans Canada. Vancouver, British Columbia.
- Gazey, W.J. and R.V. Palermo. 2000. A preliminary review of a new model based on test fishing data analysis to measure abundance of returning Chum stocks to the Fraser River. DFO Canadian Stock Assessment Secretariat Research Document 2000/159. 30 p.
- Hargreaves, B., W. Luedke, and J. Till. 2001. Application of otolith thermal mass marking in British Columbia, Canada. NPAFC Technical Report No. 3. 2001 International Workshop on Salmonid Otolith Marking. March 21, 2001, Seattle, WA, U.S.A.
- Point No Point Treaty Tribes and Washington Department of Fish and Wildlife. 2014. Five-year review of the Summer Chum Salmon Conservation Initiative for the period 2005 through 2013: Supplemental Report No. 8, Summer Chum Salmon Conservation Initiative – An Implementation Plan to Recover Summer Chum in the Hood Canal and Strait of Juan de Fuca Region, September 2014. Wash. Dept. Fish and Wildlife. Olympia, WA. 237 pp., including Appendices.
- Small, M.P., K. Currens, T. H. Johnson, A. E. Frye and J. F. Von Bargen. 2009. Impacts of supplementation: Genetic diversity in supplemented and unsupplemented populations of summer Chum salmon (*Oncorhynchus keta*) in Puget Sound (Washington, USA). Canadian Journal of Fisheries and Aquatic Sciences 66(8):1216-1229.
- Volk, E.C., Schroder, S.L., and Fresh, K.L. 1987. Inducement of banding patterns on the otoliths of juvenile Chum salmon (*Oncorhynchus keta*). Proceedings of the 1987 northeast Pacific Pink and Chum salmon workshop. Alaska Department of Fish and Game: 206–212.

APPENDICES

Appendix A. ANNEX IV, CHAPTER 6, OF THE PACIFIC SALMON TREATY

Southern British Columbia and Washington State Chum Salmon


The provisions of this Chapter shall apply for the period 2010 through 2018.

1. The Parties shall maintain a Joint Chum Technical Committee (“the Committee”) reporting, unless otherwise agreed, to the Southern Panel and the Commission. The Committee will undertake to, *inter alia*:
 - a) maintain and present historical catch and escapement information for stocks relevant to the Treaty;
 - b) utilize available information to estimate and document stock composition and exploitation rates in fisheries of concern to the Treaty;
 - c) review annually the Parties’ assessment of stock status and fisheries activities for Chum fisheries of concern to the Treaty;
 - d) identify high priority research and information needs for the Parties, including fishery and escapement monitoring and assessment, stock identification, and enhancement; and
 - e) periodically and/or when requested;
 - i. Exchange available information on the productivity and escapement requirements of stocks relevant to the treaty;
 - ii. Identify and document stocks of concern (with respect to conservation) relevant to the treaty;
 - iii. Evaluate the effectiveness and performance of management strategies; and
 - iv. Evaluate the effectiveness of alternative regulatory and production strategies recommended by the Parties.
2. When the Parties provide stock composition information for fisheries, the Committee shall evaluate and report its conclusions using bilaterally agreed upon methods.
3. Canada and the United States shall assess catch levels and make attempts to collect additional genetic samples from any Chum salmon caught during the July 1 through September 15 time period in the Boundary Area fisheries (U.S. Areas 4B, 5, 6C, 7 and 7A; Canadian Areas 18, 19, 20, 21, and 29).
4. During the period from July 1 through September 15, Canada will require the live release of Chum salmon from all purse seine gear fishing in the Strait of Juan de Fuca (Canadian Area 20) and the United States will require the same for the non-Indian seine fisheries in Areas 7 and 7A. Note: By U.S. regulation, purse seine fisheries are not permitted in U.S. Areas 4B, 5 and 6C.


5. Canada will manage its Johnstone Strait, Strait of Georgia, and Fraser River Chum salmon fisheries to provide continued rebuilding of depressed naturally spawning Chum salmon stocks, and, to the extent practicable, not increase interceptions of U.S. origin Chum salmon. Terminal fisheries conducted on specific stocks with identified surpluses will be managed to minimize interception of non-targeted stocks.
6. Canada will manage its Johnstone Strait mixed stock fishery as follows:
 - a) Inside Southern Chum salmon levels of less than 1.0 million as estimated by Canada are defined, for purposes of this chapter, as critical;
 - b) For run sizes above the critical threshold, Canada will conduct fisheries with an exploitation rate of up to 20% in Johnstone Strait of Inside Southern Chum salmon; and
 - c) When run sizes are expected to be below the critical threshold, Canada will notify the United States and will only conduct assessment fisheries and non-commercial fisheries. Commercial fisheries targeting Chum salmon will be suspended.
7. Canada will manage its Fraser River fisheries for Chum salmon as follows:
 - a) For Fraser River terminal area run sizes, identified in-season, at abundance levels lower than 900,000 Chum salmon, the Canadian commercial Chum salmon fisheries within the Fraser River and in associated marine areas (Area 29), will be suspended; and
 - b) For Fraser River terminal area run sizes, identified in-season at levels greater than 900,000 Chum salmon, Canadian commercial Chum salmon fisheries within the Fraser River, shall be guided by the limits of the in-river Total Allowable Catch set by Canada.
8. Canada will manage the Nitinat gill net and purse seine fisheries for Chum salmon to minimize the harvest of non-targeted stocks.
9. Canada shall conduct a genetic sampling program of Chum salmon taken in the West Coast Vancouver Island troll fishery if early-season catch information indicates that catch totals for the July 1 through September 15 season may reach levels similar to 1985 and 1986. Sampling, should it occur, will include catches taken from the southern areas (Canadian Areas 121-124).
10. The United States will manage its Chum salmon fishery in Areas 7 and 7A as follows:
 - a) Inside Southern Chum salmon levels of less than 1.0 million as estimated by Canada are defined, for purposes of this chapter, as critical;
 - b) For run sizes below the critical threshold, the U.S. catch of Chum salmon in Areas 7 and 7A shall be limited to Chum salmon taken incidentally to other species and in other minor fisheries, but shall not exceed 20,000, provided that catches for the purpose of genetic stock identification sampling shall not be included in the aforementioned limit;

- c) For run sizes above the critical threshold, the base catch ceiling for the U.S. Chum salmon fisheries in Areas 7 and 7A will be 130,000 Chum salmon;
 - d) Canada will provide a run size estimate of Chum salmon entering the Fraser River no later than October 22. If the estimate is less than 900,000, the U.S. will limit its fishery impacts on Fraser River Chum salmon by restricting catch in Areas 7 and 7A to not exceed 20,000 additional Chum from the day following the date the U.S. is notified. The total catch is not to exceed the catch ceiling of 130,000 Chum salmon;
 - e) U.S. commercial fisheries for fall Chum salmon in Areas 7 and 7A will not occur prior to October 10;
 - f) The U.S. will manage the Areas 7 and 7A fisheries for Chum salmon with the intent to minimize the harvest of non-targeted species;
 - g) No U.S. catch shortfalls may be accrued; however any overages shall be carried forward as indicated in (h) and (i);
 - h) Due to management imprecision, a catch in the U.S. of up to 135,000 Chum salmon will not result in an overage calculation. Catches in excess of 135,000 Chum salmon shall result in an overage being calculated by subtracting 130,000 from the total Chum catch. Overages will be accounted for by reducing the U.S. annual catch ceiling in up to two subsequent non-critical Inside Southern Chum salmon years; and
 - i) From the day following the date the U.S. is notified of a run size below the critical threshold as defined in 10(b) or (d), any catches in excess of 20,000 Chum salmon will result in an overage. Overages will be accounted for by reducing the U.S. annual catch ceilings in up to two subsequent non-critical Inside Southern Chum salmon years.
11. The United States shall conduct its Chum salmon fishery in the Strait of Juan de Fuca (United States Areas 4B, 5, and 6C) so as to maintain the limited effort nature of this fishery, and, to the extent practicable, not increase interceptions of Canadian origin Chum salmon. The United States shall continue to monitor this fishery to determine if recent catch levels indicate an increasing level of interception.
 12. All information concerning by-catch of other salmon species from the Chum salmon fisheries covered by this chapter will be shared between the Parties in the annual Post Season Report.
 13. Should circumstances arise that are inconsistent with either Parties understanding of the intent of the chapter, the Southern Panel will discuss the matter postseason and explore options for taking the appropriate corrective actions.

Appendix B-1. United States commercial salmon catch areas for the Strait of Juan de Fuca and Northern Puget Sound.




Appendix B-2. Canadian Pacific Fishery Management Areas (PFMA) for Southern B.C. - Marine areas.



G:\projects\science\PieterVanWill\BC Albers\Southern Chum CU's map\Working\southern bc chum salmon - Marine areas.mxd

Appendix B-3. Canadian Pacific Fishery Management Areas (PFMA) for Southern B.C. - Lower Fraser River and surrounding areas.



G:\projects\science\PieterVanWill\BC Albers\Southern Chum CU's map\Final\southern bc chum salmon - lower fraser river and surrounding areas.mxd

2015 POST-SEASON MEETING BRIEFING FOR THE COMMISSIONERS

Chum Technical Committee

MEETING PLAN

Session Objectives:

- Review and discuss preliminary post-season 2014 fisheries information
- Collate and review report items for 2012 and 2013 final post-season report
- Finalize 2012 annual report for submittal
- Initiate drafting of 2013 final report
Review and discuss research and analysis activities essential to the Committee tasks
- Continue work on existing and possible Southern Fund related projects associated with Southern Chum.

Outstanding and Upcoming Issues:

- Status report required on 1st phase of Chum GEM project
- Review and Summarize 3rd year of Mixed stock fishery project

Noteworthy highlights from the 2014 Post-Season Reports:

Canadian Post-Season Report

- Varied Chum returns throughout the Inner South Coast
- Inner South Coast Chum aggregate abundance below average but an improvement over the main 2010 brood year and above critical threshold
- Harvest occurred in Johnstone Strait and Fraser River

US Post-Season Report

- Harvest summaries for US catch areas Straits of Juan de Fuca, San Juan Islands & Point Roberts regions
- Tissue collections from boundary area fisheries for future genetic stock ID.

Appendix D. Concept proposals (see work plans) submitted to Pacific Salmon Commission Southern Endowment Fund

Project Title: S15-102A Joint US and CA Mixed-stock Chum Fisheries Sampling Design, continued year 4 of 4 (Araujo et al. 2016)

In order to facilitate management responses to Southern Chum stock strength, in accordance with Annex IV, Chapter 6 of the Pacific Salmon Treaty (Treaty) it is necessary to provide the catch composition in fisheries targeting Southern origin Chum populations. We are proposing to sample Southern BC and US mixed stock Chum fisheries to determine stock composition to the Canadian Conservation Unit (CU) and United States Management Unit (MU) level using genetic mixed stock analysis. In addition, run timing, distribution and diversion by CU and MU will be explored for run reconstruction. This proposal is informed by the results from the Southern fund project (Southern Study Area Chum stock distribution assessment in Washington and San Juan Island, Point Roberts and BC southern Gulf fisheries) that occurred 2007–2010. The goal is to sample the mixed stock Chum fisheries for four consecutive years in a multi-agency sampling effort.

Project Title: S15-108 Chum salmon stock identification in southern British Columbia and Puget Sound using microsatellites, assessment of the baseline expansion with an additional 53 populations. (Beacham et al. 2016)

Chum salmon (*Oncorhynchus keta*) from 53 populations in southern British Columbia and Puget Sound were surveyed for variation at 28 microsatellite loci was completed as part of a baseline expansion for microsatellites. Initially 20 populations were analyzed for a SNP microsatellite baseline comparison in 2014. Significant genetic differentiation was observed among Chum salmon populations sampled in the different geographic regions surveyed. The F_{ST} value over all populations and loci was 0.012 (SD=0.001), with individual locus values ranging from 0.004 (*Omm1070*, *One102*) to 0.039 (*Oki100*). Populations clustered into geographic regions. Microsatellites with larger numbers of alleles were generally provided more accurate estimates of stock composition of single-population samples than did loci with smaller numbers of alleles. The average accuracy of estimates of stock composition of the single-population samples was 48.8%, although estimates of individual populations ranged from 0.0% for the Tsouwwin River population to 100.0% for the Salmon River population in Hood Canal. The average regional accuracy of estimates of stock composition of the single-population samples was 85.2%, with estimates of individual populations ranging from 10.3% for the Demamiel Creek population to 100.0% for the Salmon River population in Hood Canal. Standard deviations about stock composition estimates were higher for population-specific estimates compared with regional estimates of stock composition, and the utilization of baselines with larger numbers of microsatellites for estimation of stock composition resulted in estimates with higher precision (lower standard deviations). Increased numbers of sampled fish per population also resulted in estimates of stock composition with higher precision.

Project Title: Adapting benchmarks of biological status for persistent changes in productivity and variability in exploitation history with a focus on data-limited populations (Conservation Units) of Chum Salmon in southern BC. (Holt et al.)

The PST Chum Annex requires biological benchmarks to inform the development of fishery reference points for PST related fisheries; including the lower fishery reference point for the

Johnstone Strait fisheries as well as subsequent terminal fisheries. Biological benchmarks for data-limited populations have been proposed and are currently being applied to Conservation Units (CUs, population units of biological assessment under Canada's Wild Salmon Policy) of Chum Salmon in southern BC. However, recent simulation modelling has shown that those benchmarks, which are derived from time-series of spawner abundances alone, are associated with high probabilities of extirpation under low and/or declining stock productivity (Holt and Folkes, in review). In addition, previous unpublished studies suggest these benchmarks may be higher than necessary to achieve conservation objectives when exploitation rates are low (cited in Fair et al. 2010). For this project, we will (1) identify candidate benchmarks for data-limited CUs and revised versions that account for persistent changes in productivity and variability in exploitation rate history, (2) evaluate them against benchmarks based on data-intensive methods, and (3) apply them to Chum management units (through component CUs) within the context of the PST Chum Annex.

Project Title: Chum salmon southern area genetic baseline enhancement Part 2, Year 2 of 2: Expanded SNP collections, genotyping, and amplicon development (Small et al.)

This project is a continuation of the Southern Boundary Restoration and Enhancement Fund 2012-2014 project, "Chum Salmon Southern Area Genetic Baseline Enhancement." This project expands the nascent Chum SNP baseline, developed in part during the 2012-2014 phases I and II of this project, by increasing and expanding the geographic coverage of collections for better stock assessment, management and research, and by increasing sample sizes, where needed, from collections already genotyped. In addition to field collection of new samples and SNP genotyping, we are developing amplicon sequencing assays from the existing and additional SNPs discovered from the RAD sequencing conducted during Parts 1 and 2 of this project. Amplicon sequencing assays will enable us to expand the number of SNPs genotyped from the current 192 TaqMan assays to several hundred SNPs, at a reasonable cost. Increasing the number of SNPs will increase power to identify stocks that are now poorly resolved with the current set of microsatellite or SNP assays. Some of the SNPs are at loci under selection which will provide information on Chum salmon response to environmental variation, data that will be useful for the ChumGEM model. We will make available these amplicon sequencing panels to any genetic laboratory and will also make available the Chum salmon SNP genetic baseline at a publicly accessible website.

To expand geographic coverage, tissue samples from representative stocks (roughly 20 stocks from Washington (WA) and southern British Columbia (SBC)), will be genotyped at SNP loci and combined with existing SNP baseline data (10 SBC and 10 WA stocks under Southern Fund plus 3 SBC and 9 WA stocks under Salton-Stall Kennedy Fund, see Appendix I for population list) generated in Part 1 of this study. This study enhances the existing SNP baseline, particularly in Canada, and thus will provide more accurate estimates of stock composition for boundary area mixed stock fisheries (e.g. Strait of Juan de Fuca and Strait of Georgia (Figure 1), see Rawson et al. 2014). We will subject the resulting SNP baseline to cross-validation simulations to test the accuracy of the baseline in estimating stock contributions in boundary area mixed stock fisheries.