

PSC Northern Fund Final Report

Project Number: NF-2016-I-59

Project Title: Mixed Stock Analysis of US Districts 108 and 111 Chinook Salmon Fisheries 2016

Project Manager: Sara Gilk-Baumer, Alaska Dept. of Fish and Game - Gene Conservation Laboratory, (907) 267-2535, sara.gilk@alaska.gov

Duration of Project: Start date: May 1, 2016 End Date: June 30, 2017

Abstract:

The Stikine and Taku rivers in Southeast Alaska support Chinook salmon runs important for various commercial, sport, and aboriginal fisheries in both Alaska and Canada. This project continues the use of genetic stock identification (GSI) of Chinook salmon harvested in the 2016 commercial gillnet and sport fisheries in districts 108 and 111 by screening 13 microsatellite genetic markers in 1,296 salmon. The mixed stock analysis results indicated that the stock composition of commercial and sport fisheries in districts 108 and 111 varied by fishery. The District 108 directed gillnet fishery primarily targeted enhanced Andrew Creek Chinook salmon (64%), while the District 108 directed troll fishery primarily harvested fish that originated from outside Alaska (60%). The District 108 sport fishery harvested a variety of stocks, but was also dominated by enhanced Andrew Creek fish (31%). In District 111, harvest was primarily Taku-origin and enhanced Andrew-origin fish (58% and 36%, respectively). There was no directed gillnet fishery in District 111 in 2016.

Introduction:

The Stikine and Taku rivers in Southeast Alaska (SEAK) support Chinook salmon runs important for various commercial, aboriginal, and recreational fisheries in both the United States (U.S.) and Canada. Included in these are U.S. commercial gillnet fisheries in Alaskan Districts 108 and 111, as well as sport fisheries near Wrangell, Petersburg, and Juneau. U.S. fisheries in these areas harvest stocks of Chinook salmon bound for SEAK and for tributaries in the transboundary Stikine and Taku rivers. Catches of Stikine and Taku river Chinook salmon stocks are subject to a harvest sharing agreement, in which the U.S. and Canada are each given an Allowable Catch (AC) according to provisions outlined in the Transboundary Annex (Annex IV) of the Pacific Salmon Treaty (PST). Allowable catches are specified by the Pacific Salmon Commission (PSC) and rely on catch, escapement, recruitment information, and stock composition estimates to forecast indices of abundance in PST fisheries.

Until 2006, stock composition of harvests was estimated primarily using coded-wire tags, however, only a small portion of the out-migrating Chinook salmon are annually coded wire tagged leading to subsequent harvest estimates that lack the level of precision needed for management. Genetic stock identification (GSI) provides a complementary set of accurate, precise, and reliable stock composition estimates necessary to meet the needs of the abundance-based management regime for Chinook salmon in these fisheries. Since 1999, GSI has been successfully used to

estimate the composition of the commercial troll fishery harvest (Crane et al. 2000; Gilk-Baumer et al. 2013, 2017a, 2017b; Templin et al. 2011), gillnet and seine harvest (e.g. Gilk-Baumer and Carlile 2012), and sport fishery harvests (Gilk-Baumer et al. 2017c). In addition, the Transboundary Technical Committee has been using GSI estimates from Districts 108 and 111 for post-season analyses calculating harvest-share estimates for more than five years.

This project extended GSI of Chinook salmon harvested in drift gillnet fisheries and sport fisheries in Districts 108 and 111 through 2016. These data are also used to determine Chinook salmon exclusions from the all-gear limits in place for the SEAK AABM fishery, and to estimate actual contributions of above-border Stikine and Taku Chinook salmon to the sport and commercial fisheries in Districts 108 and 111.

Objectives:

The objective of this project is to estimate the stock composition of Southeast Alaska Chinook salmon fisheries in Districts 108 and 111 in 2016 such that the estimates are within 5% of the true value 90% of the time. This was accomplished through the following tasks:

- Representatively sample Chinook salmon from commercial and sport fishery harvests relevant to Annex IV, Chapter 1 (3) (Transboundary Rivers) of the Pacific Salmon Treaty.
 - Directed fisheries – sample harvests from the commercial drift gillnet fisheries operating in Districts 108 and 111 between May and August 2016.
 - Sport fishery – sample the sport fishery harvests in Districts 108 and 111 between April and August 2016.
- Assay up to 1,600 individual genotypes from sampled Chinook salmon at the loci in the current PSC baseline of genetic markers.
- Estimate the relative stock composition of the Southeast Alaska fisheries using genetic stock identification.

Approach:

Fishery Sampling

Chinook salmon were collected from commercial gillnet and directed troll landings at processors in Southeast Alaska, and in the sport fishery by onboard participants and by creel census samplers. During sampling, Chinook salmon were selected without regard to size, sex, adipose fin-clip, or position in the hold. Axillary process tissue was dissected from sampled fish and dried onto Whatman paper. Along with each individual sampled, matched information was recorded such as size, sex, date, vessel, and age (from scale samples). At the end of the fishery, samples were transported back to the ADF&G Gene Conservation Laboratory, Anchorage, for analysis. Associated data is archived as part of the ASL database maintained by ADF&G.

Representative tissue collections of individuals for mixture analysis was created by subsampling up to 800 large (≥ 660 mm mid-eye-to-fork length) individuals harvested between statistical weeks 17–29 from the collected samples. Because the PST applies to large Chinook salmon for transboundary management, only large Chinook salmon were included in the analysis. Target mixture sample sizes was 200 individuals per mixture to achieve acceptable levels of accuracy and precision. Due to the vagaries of fisheries and fishery sampling, target sample sizes were not available for every stratum. Sample sizes smaller than the target were analyzed, but strata represented by fewer than 100 individuals would be pooled into larger groups for analysis. Since

directed Chinook salmon gillnet fisheries did not occur in 2016 in District 111, commercial fishery samples were obtained by sampling Chinook salmon caught incidentally in sockeye gillnet fisheries that district.

Laboratory Analysis

Samples were assayed for DNA loci developed by the Genetic Analysis of Pacific Salmon (GAPS) group funded by the Pacific Salmon Commission (PSC) for use in PST fisheries (Seeb et al. 2007). Laboratory methods are well established and have been described in previous proposals and reports. Briefly, DNA was extracted from fin and muscle tissue and the polymerase chain reaction (PCR) was used to label DNA fragments at specified locations in the genome. PCR fragment analysis was done on an AB 3730 capillary DNA sequencer and PCR bands were visualized and separated into bin sets using AB GeneMapper software v4.0. All laboratory analyses followed protocols accepted by the CTC of the PSC. The data collected were individual genotypes for each locus. Genotype data are stored in an Oracle database (LOKI) on a network drive maintained by ADF&G computer services.

Mixture Analysis

Stock composition estimates for stock groups were generated using the BAYES (Pella and Masuda 2001) software package. The estimation was run using five chains without thinning with MCMC sample sizes of 40,000. Inference was based on the combined distributions of the last 20,000 samples of each chain. We defined prior parameters for each stock group to be equal to results from the corresponding estimates generated for the 2015 fisheries, with the prior for each stock group subsequently divided equally to populations within that stock group. Individual population or stock estimates were calculated and then summed into reporting regions. The mean of the regional compositions in the posterior distribution were reported as the best estimate and the 90% credibility intervals for all group contribution estimates were computed from the posterior distribution. The goal was to report estimates that fall within the precision and accuracy guidelines set by the TTC in April 2013 (to estimate the proportions of stocks within 10% of the true mixture proportions 90% of the time).

A total of 4 seasonal stock composition estimates were made for the 2016 fisheries: District 108 gillnet, District 108 directed troll, District 108 sport, and District 111 sport. Insufficient samples were obtained from Chinook salmon collected in the District 111 sockeye gillnet fisheries to produce an estimate within the precision and accuracy guidelines specified by the TTC.

Results/Findings:

Fishery sampling

A total of 1,296 Chinook salmon were sampled in districts 108 and 111 commercial and sport fisheries. In District 108, 484 fish were sampled in the directed commercial gillnet fishery, 241 fish were sampled in the directed commercial troll fishery, and 256 fish were sampled in the sport fishery during the treaty period. In District 111, 94 Chinook salmon were sampled as part of the commercial sockeye gillnet fishery and 221 were sampled in the sport fishery.

Laboratory analyses

Of the 1,296 total samples collected in District 108 and 111 commercial gillnet and sport fisheries, all samples were genotyped at 13 microsatellite genetic markers. During quality control procedures

a total of 143 samples were reanalyzed at all 13 markers for a total of 1,859 comparisons. The average failure rate was ~3.6%. No inconsistencies were found across all comparisons.

Mixture analysis

Mixtures of fish representing catches by fishery and district were analyzed. Stock composition estimates can be found in Table 2. All estimates meet the minimum criteria for precision and accuracy accepted by the Pacific Salmon Commission (PSC) Transboundary Technical Committee (within 10% of the true mixture 90% of the time), except for the District 111 gillnet fishery. This estimate was not provided.

Mixture composition varied by fishery and location (Table 2). The largest component of the District 108 directed gillnet and sport fisheries was the *Andrew* reporting group (64% and 31%, respectively), almost all of which are likely enhanced (i.e. hatchery-origin). The *Other*, *Stikine*, and *Southern Southeast Alaska (SSEAK)* groups were also important contributors to the District 108 sport fishery (21%, 19%, and 18%, respectively), and the *Stikine* group was an important contributor to the directed gillnet fishery (15%). In contrast, the *Other* reporting group was the largest contributor to the District 108 directed troll fishery (58%), followed by the *Stikine* (17%) and *SSEAK* (17%) groups. In District 111, the *Taku* reporting group was the largest contributor to the sport fishery at 58%, followed by the *Andrew* group (36%).

Evaluation:

We accomplished the following:

- A total of 1,296 samples were collected from large (> 600mm MEF) Chinook harvested between statistical week 17–29 in commercial gillnet, troll, and sport fisheries in Districts 108 and 111.
- All samples from were assayed for genotypes for the 13 microsatellite loci and quality control procedures revealed a low rate of inconsistencies.
- Mixture analyses estimated the contributions of 5 reporting groups including *Taku*, *Andrew* (mostly hatchery-origin), *Stikine*, *SSEAK*, and *Other*.
- Mixture analyses estimate the seasonal stock composition for the District 108 commercial gillnet and troll fisheries, the District 108 sport fishery, and the District 111 sport fishery. Insufficient samples were collected to generate an estimate for the District 111 commercial gillnet fishery.
- Results were incorporated into harvest estimates for PST purposes by the Transboundary Technical Committee (TTC 2019).

Project Products:

Results from this project have been presented both to ADF&G Commercial Fisheries management staff and to the bilateral PSC Transboundary Technical Committee.

Date Prepared: October 21, 2019

References:

Crane, P. A., W. D. Templin, D. M. Eggers, and L. W. Seeb. 2000. Genetic stock identification of Southeast Alaska Chinook salmon fishery catches. Final report of the Alaska Department of Fish and

- Game to US Chinook Technical Committee, U.S. Letter of Agreement Award No. NA87FPO408. Regional Information Report 5J00-01, Alaska Department of Fish and Game, Anchorage.
- Gilk-Baumer, S., and J. Carlile. 2012. Estimating the Chinook salmon stock composition of Southeast Alaska District 108 and 111 fisheries, 2011. Final report of the Alaska Department of Fish and Game to the Pacific Salmon Commission Northern Fund, Award No. NF-2011-I-12.
- Gilk-Baumer, S., W. D. Templin, and L. W. Seeb. 2013. Mixed stock analysis of Chinook salmon harvested in Southeast Alaska commercial troll fisheries, 2004–2009. Alaska Department of Fish and Game, Fishery Data Series No. 13-26, Anchorage.
- Gilk-Baumer, S., D. F. Evenson, and W. D. Templin. 2017a. Mixed stock analysis of Chinook salmon harvested in Southeast Alaska commercial troll fisheries, 2010-2014. Alaska Department of Fish and Game, Fishery Data Series No. 17-37, Anchorage.
- Gilk-Baumer, S., D. F. Evenson, K. Shedd, and W. D. Templin. 2017b. Mixed stock analysis of Chinook salmon harvested in Southeast Alaska commercial troll fisheries, 2015. Alaska Department of Fish and Game, Fishery Data Series No. 17-41, Anchorage.
- Gilk-Baumer, S., W. D. Templin, D.F. Evenson, and E. L. Jones. 2017c. Mixed stock analysis of Chinook salmon harvested in the Southeast Alaska sport fishery, 2004-2015. Alaska Department of Fish and Game, Fishery Data Series No. 17-43, Anchorage.
- Pella, J. and M. Masuda. 2001. Bayesian methods for analysis of stock mixtures from genetic characters. *Fisheries Bulletin* 99(1):151-167.
- Seeb, L. W., Antonovich, A., Banks, M., Beacham, T., Bellinger, R., Blankenship, S., Campbell, M., Decovich, N., Garza, J. C., Guthrie, C., Lundrigan, T., Moran, P., Narum, S., Stephenson, J., Supernault, J., Teel, D., Templin, W. D., Wenburg, J. K., Young, S., and Smith, C. T. 2007. Development of a standardized DNA database for Chinook salmon. *Fisheries* 32(11):540-552.
- Templin, W. D., J. M. Berger, and L. W. Seeb. 2011. Genetic stock identification of Chinook salmon harvested in the Southeast Alaska troll fishery, 1999–2003. Alaska Department of Fish and Game, Fishery Manuscript No. 11-03, Anchorage.
- TTC (Transboundary Technical Committee). 2019. Final estimates of Transboundary River salmon production, harvest and escapement and a review of joint enhancement activities in 2016. Pacific Salmon Commission Report TCTR(19)-1, Vancouver.

Table 1. Sample sizes for each mixture indicating the number of fish genotyped and samples removed in the quality assurance process (missing > 20% of genotypes or duplicate individuals).

District	Fishery	Sample Size			Final
		Genotyped	Missing	Duplicate	
108	Gillnet	484	23	0	461
108	Sport	256	2	0	254
108	Troll	241	15	0	226
111	Gillnet	94	0	0	94
111	Sport	221	2	0	219

Table 2. Stock composition for each mixture, including the mean estimate, standard deviation and 90% credibility interval (Lo = 5% and Hi = 95%).

District	Fishery	Sample Size		5 Reporting Groups				
				Taku	Andrew	Stikine	SSEAK	Other
108	Gillnet	293	Estimate	0.074	0.644	0.145	0.098	0.038
			SD	0.027	0.033	0.033	0.023	0.014
			Lo	0.036	0.590	0.090	0.063	0.018
			Hi	0.122	0.697	0.200	0.137	0.065
108	Sport	251	Estimate	0.116	0.306	0.187	0.182	0.208
			SD	0.031	0.037	0.038	0.030	0.028
			Lo	0.068	0.246	0.127	0.134	0.163
			Hi	0.169	0.368	0.251	0.235	0.257
108	Troll	197	Estimate	0.011	0.068	0.174	0.169	0.579
			SD	0.019	0.023	0.037	0.036	0.045
			Lo	0.000	0.033	0.115	0.112	0.506
			Hi	0.051	0.109	0.236	0.231	0.655
111	Sport	211	Estimate	0.584	0.355	0.003	0.005	0.052
			SD	0.037	0.035	0.008	0.005	0.016
			Lo	0.523	0.298	0.000	0.000	0.029
			Hi	0.644	0.414	0.020	0.015	0.080