

**Kilbella/Chuckwalla Chinook Salmon
Stock Recovery Enhancement, 2015-2016
Progress Report**



Prepared for:

Pacific Salmon Commission
1155 Robson St, Vancouver, BC V6E 1B5

Prepared by:

LGL Limited environmental research associates
9768 Second Street, Sidney, BC V8L 3Y8

September 2016



**Kilbella/Chuckwalla Chinook Salmon
Stock Recovery Enhancement, 2015-2016
Progress Report**

Prepared by:

Jason J. Smith, Karl K. English¹, and Sandie MacLaurin²

Prepared for:

Pacific Salmon Commission
1155 Robson St, Vancouver, BC V6E 1B5

September 2016

¹ LGL Limited environmental research associates, 9768 Second Street, Sidney, BC V8L 3Y8

² Fisheries and Oceans Canada, Box 340, Hagensborg, BC V0T 1H0

INTRODUCTION

Stock assessment and enhancement priorities for Rivers Inlet have been examined and discussed extensively over the past four years. This project was one of the top priorities for work on Chinook and other salmon species in Rivers Inlet identified by a Rivers Inlet Salmon Steering Committee (RISSC) comprised of representatives from the Wuikinuxv First Nation, local lodge owners, Fisheries and Oceans Canada (DFO), Rick Hansen Foundation (RHF), Pacific Salmon Foundation (PSF), and independent scientists. The RISSC was established in 2011 to guide the development of an immediate action plan for Wannock River Chinook salmon and long-term plans for other Rivers Inlet stocks and salmon species.

Historically, Chinook salmon from the Kilbella/Chuckwalla rivers (Figure 1) represented a significant portion of the Chinook salmon caught in the Rivers Inlet recreational fishery (Nelson et al. 2000). Annual escapement monitoring through 2010 indicated a 5-10 fold decline in spawners from the numbers observed in the 1998-2002 period when these stocks were being augmented through small-scale enhancement. In 2010, the escapement estimates for Chinook salmon were only 150 and 75 for the Kilbella and Chuckwalla rivers, respectively. Results from surveys conducted in September 2013 indicated that the number of Chinook salmon spawners in these rivers was similar to the very low numbers observed in 2010. Chinook salmon escapement estimates derived from the 2014 aerial surveys and broodstock-collection efforts were 200-300 and 100 for the Kilbella and Chuckwalla rivers, respectively.

PROJECT GOALS AND OBJECTIVES

The purpose of this project was to increase Chinook salmon returns to the Kilbella/Chuckwalla watersheds through a small-scale, targeted, enhancement program. Eggs and milt were collected from Kilbella/Chuckwalla Chinook salmon and reared at the Snootli Hatchery near Bella Coola. Fed fry were returned to net pens located near the Kilbella River estuary for further rearing prior to release. Information on the contribution of these fish to coastal fisheries will be obtained by applying coded-wire tags (CWT) to a large portion of the smolts released, and recovering marked fish from ongoing sampling programs in BC and Alaska.

The enhancement goal for 2014 and 2015 was to obtain 50,000 eggs and adequate milt from each stock in each year. This equates to 10 mature Chinook salmon of each sex from each river. The enhancement goal for 2016-2018 will be to collect 75,000 eggs and adequate milt annually from each stock. The ultimate goal is to increase number of annual spawners in the Kilbella and Chuckwalla rivers to an initial target of 1,000 adult Chinook salmon.

In the first year of this program (2014-15), Snootli Hatchery personnel and Wuikinuxv Fisheries technicians collected eggs from 4 female fish (3 Kilbella, 1 Chuckwalla), transported approximately 17,500 eggs to the Snootli Hatchery, transferred 16,486 fry to net pens in Rivers Inlet (12,207 from Kilbella; 4,279 from Chuckwalla), and released a total of 16,450 5.4-g smolts in the Spring of 2015 (English and MacLaurin 2015). Survival rates from egg take to release averaged 87.4% (95% for Kilbella, 71.4% for Chuckwalla; English and MacLaurin 2015).

The following sections document results from 2015-16, the second consecutive year of this program. Plans for 2016-17 activities are also discussed.

2015 PROJECT LOGISTICS

Personnel from Wuikinuxv Fisheries, DFO, Good Hope Cannery, and LGL Limited contributed to the 2015-16 project. In August and September, out-of-town crew members stayed at the Eagle's Nest Bed and Breakfast in Wuikinuxv Village. Helicopter support to conduct broodstock collection and aerial/swim surveys was provided by West Coast Helicopters (Port McNeill, BC). Eggs were transported from the Wuikinuxv Village airport to Bella Coola by West Coast Helicopters and Bella Coola Air (Hagensborg, BC). A 16-ft, flat-bottomed boat (Lowe 1648-MT) equipped with a 30-HP outboard jet was used to conduct reconnaissance trips and swim surveys in the lower Chuckwalla River.

RESULTS OF AERIAL AND SWIM COUNTS IN 2015

Kilbella River: In the Kilbella River, aerial counts of live and dead Chinook salmon were conducted via helicopter on 25 August and 4 September, 2015 (Table 1). Counts were stratified into 3 river sections: Mallon Creek (rkm 40.0) to Cascades (rkm 29.5), Cascades to 9-Mile Bridge (rkm 17.3), and 9-Mile Bridge to the mouth (rkm 0.0). A total of 4 live Chinook salmon were counted on 25 August, and 1 live Chinook salmon was counted on 4 September. Water clarity was very poor on 25 August despite extremely low water levels, likely due to run-off of glacial silt. Water clarity was better on 4 September; however the river still contained glacial silt making it difficult to count fish in deep water.

Chuckwalla River: In the Chuckwalla River, two helicopter counts (25 August and 3 September) and one swim count (6 September) were conducted in 2015 (Table 1). Counts were stratified into 4 river sections: Johns Creek (rkm 23.0) to Cascades (rkm 20.0), Cascades to the Canyon (rkm 14.0), the Canyon to the Bridge Pool (rkm 6.0), and the Bridge Pool to the mouth (rkm 0.0). On 25 August, a total of 21 live and 1 dead Chinook salmon were counted; and on 3 September, 6 live Chinook salmon were counted. Only 1 dead Chinook salmon was observed during the swim count on 6 September from the Bridge Pool to the mouth. Similar to the Kilbella River, water clarity on the 25 August survey was poor despite extremely low water levels. Water clarity was significantly better for the 3 September aerial count and 6 September swim count.

In 2015, 67% (4 of 6 fish on 3 September) and 100% (21 of 21 fish on 25 August) of all Chinook salmon counted during helicopter surveys of the Chuckwalla River were observed upstream of the Canyon Pool (Table 1). In comparison, only 7-8% of Chinook salmon counted in 1999 (4 surveys from 1-17 September) were observed upstream of the Canyon Pool (see Table 7 in Nelson et al. 1999).

RESULTS OF BROODSTOCK COLLECTION EFFORTS IN 2015

Kilbella River: In 3 days of effort (27 August, 4-5 September), 38 gillnet sets (100 ft long, 8.25" mesh) were made between rkm 40.0 and rkm 30.5 in the upper Kilbella River (Table 2; Figure 2). A total of 33 live Chinook salmon were captured (Figure 3), including 32 unmarked fish (10 female, 22 male) and 1 recapture (male). In addition, 3 carcasses (all female) were encountered on 5 September. Six females were spawned, including three that were full of eggs and three that were partially full of eggs (Table 3). Milt was taken from 9 males. Females averaged 74.2 cm POH (range: 68.0-88.0 cm POH) and males averaged 65.0 cm POH (range: 53.0-76.0 cm POH; Figure 4). Scales were collected from 10 females and 18 males; however, age results were not available at the time this report was prepared.

Chuckwalla River: In 3 days of effort (26 August, 3 September, 11 September), 43 gillnet sets were made between rkm 23.3 and rkm 17.0 in the upper Chuckwalla River (Table 2; Figure 5). A total of 28 live Chinook salmon were captured (Figure 6), including 25 unmarked fish (14 female, 11 male) and 3 recaptures (1 female, 2 male). Two carcasses (female) were encountered on the Chuckwalla River, one of which was found below the Bridge Pool (rkm 6.0) during the swim count on 6 September. Two females were spawned, including one that was full of eggs and one that was partially full of eggs (Table 3). Milt was taken from 8 males. Females averaged 74.3 cm POH (range: 69.0-85.0 cm POH) and males averaged 69.2 cm POH (range: 59.0-83.0 cm POH; Figure 4). Scales were collected from 16 females and 11 males; however, age results were not available at the time this report was prepared.

As in 2014, all unripe females were marked and released with the hope that they would be captured later in the season. Unripe fish captured on 26-27 August were not retained/held in fish tubes because of concerns over helicopter availability and expected high water levels in the following days. On 11 September, the crew was prepared to transport all unripe fish encountered to a holding area in the lower Chuckwalla River, however, no green fish were captured.

Despite seeing an increase in catches in 2015 compared to 2014, Chinook salmon abundance in both rivers continued to be very low, and the crew was not able to collect eggs from 10 ripe female Chinook salmon in each river.

RESULTS OF 2015-16 EGG INCUBATION AND FRY RELEASE

In total, 32,853 eggs (21,252 Kilbella, 11,601 Chuckwalla) were planted at the Snootli Hatchery for incubation in August/September 2015 (Table 4). This was a 74% increase in the number of eggs planted compared to 2014.

On 20 May 2016, 19,518 Kilbella and 10,793 Chuckwalla smolts (5 g, CWT-marked) were released from a net pen in Kilbella Bay, meeting the target time and size of release goals. Egg-to-fry survival was excellent for both stocks and there were few mortalities through rearing. Overall egg-to-release survival was 92% for the Kilbella and 93% for the Chuckwalla (Table 4).

AGE COMPOSITION

Of the 7 scale samples successfully aged for Chinook salmon caught in the Kilbella River in 2014, 3 fish were age 5₂ and 4 fish were age 6₂ (Table 5). Only two samples were successfully aged from the Chuckwalla River in 2014, one was age 4₂ and the other was age 5₂.

For the 2015 brood year, successfully aged scale samples for Chinook salmon caught in the Kilbella River consisted of fish of ages 3₁ (2) 4₁ (1), 3₂ (1), 4₂ (2), and 5₂ (18; Table 6). Chinook salmon caught in the Chuckwalla River in 2015 were comprised of fish of ages 5₁ (1), 4₂ (5), 5₂ (15), and 6₂ (2).

PLANS FOR 2016 BROODSTOCK COLLECTION EFFORTS

The experience gained from 2014 and 2015 has been used to develop plans for 2016 that will address the challenges encountered to date related to helicopter access, fish holding, and broodstock transport. The four project components for 2016 will include:

- 1) additional survey effort to improve the reliability of annual escapement estimates and determine the best locations and approach for obtaining broodstock;
- 2) broodstock collection in late August and early September;
- 3) egg incubation, fry ponding/rearing, and CWT application at Snootli Hatchery; and
- 4) transport of fry to net pens in Rivers Inlet for further rearing from April until release in May of 2017.

These activities will be coordinated with the enhancement efforts for Wannock River Chinook salmon to ensure there is capacity to accommodate both groups of fish at the Snootli Hatchery and in the net pens. Wuikinuxv Fisheries personnel will assist with escapement monitoring and broodstock collections, and will be responsible for all tasks associated with rearing Chinook salmon fry in the net pens prior to release.

As in 2015, escapement numbers and broodstock collection locations will be determined using a combination of aerial, snorkel, and ground-based survey techniques in 2016. Broodstock collection will be conducted using gill nets, similar to the methods used in 2014 and 2015, which have also been used successfully for Wannock River Chinook salmon. The use of holding pens rather than holding tubes should minimize the potential for bear predation on female broodstock while they ripen in the deep pool near the old bridge site on the Chuckwalla River. The proposed 2016 escapement monitoring and broodstock collection program will include a full-time field program supervisor (Jason Smith, LGL) who will be on-site for the entire broodstock collection period. Fishing effort in 2016 will be focused in the upper reaches of both rivers (i.e., from the canyon to John's Creek on the Chuckwalla River [rkm 14–23], and from the cascades to Mallon Creek on the Kilbella River [rkm 30–40]). Relatively few Chinook salmon were observed in the lower reaches of either river in 2015. Any Chinook salmon that were present in these lower reaches were co-mingled with hundreds of pink and chum salmon which would make it difficult to target Chinook salmon with a gillnet.

Similar to 2014 and 2015, one 3-4 person field crew will be used initially in 2016. If necessary however, a second crew will be deployed to increase fishing effort. All costs associated with the addition of a second crew would be covered by in-kind contributions from Duncanby Lodge, and the crew would be directed by the LGL field program supervisor. A holding site in the lower Chuckwalla River will be used to retain pre-spawn Chinook salmon for short periods of time until they are ripe and ready for egg collection. During the 2015 season, potential holding sites were identified in the lower Chuckwalla River that were accessible by both helicopter and small jet boat. It is important that these sites are accessible so that crews can regularly monitor the health of fish being held and easily transport eggs to the airstrip in Wuikinuxv Village. It had been previously proposed to operate a temporary field camp in the lower Chuckwalla River in 2015, as well as reconstruct a riverside ATV trail that had been used in 1999 (Nelson et al. 2000). However, reconnaissance in 2015 showed that it was not feasible to reconstruct the ATV trail as part of this project. Also, crews based in Wuikinuxv Village were able to access the lower 6 km of the Chuckwalla River in 2015 using a 15-ft jon boat equipped with a 30-HP outboard jet, even at low river flows.

PLANS FOR 2016-17 EGG INCUBATION AND 2017 FRY RELEASE

Eggs and sperm will be collected on the rivers and the gametes flown to Snootli Hatchery. Eggs will be fertilized, rinsed, and placed in Heath-type, vertical incubators with a 100 ppm iodine solution. The antiseptic bath will be timed for 10 minutes, and then the trays will be placed into flowing water. At the eyed stage of development, eggs will be shocked, picked, enumerated, and placed back in Heath trays. Ponding will be done to two Capilano Type troughs arranged in lines of two. Fish will be split equally between the two troughs with 50,000 fry held in the upper section of each trough. Once densities reach 32 kg/m³, half of the fish will be transferred to the lower section of each trough. When fry are over 1.5 g, all fish will be CWT marked, which will also verify fish numbers. Rearing will continue until fry reach ~3 g, at which point they will be transported to net pens in Rivers Inlet for final rearing and release. One float with two nets (15' long x 15' wide x 15' deep per net) will be used. This will ensure density at maximum size will not exceed the recommended 5 kg/m³. Fish will be feed a daily ration until they reach an average weight of 5 g (target date to reach this size is 15 May). Records of mortality will be kept throughout rearing so an accurate number of fish released can be reported.

PROJECT FUNDING AND COST FOR 2015-16

The PSC funding available for activities associated with the second year (2015-16) of Kilbella-Chuckwalla Chinook recovery enhancement efforts was \$98,000. The current contract is slated to end on 30 June 2016, with a final technical report and expenditure report due within 45 days of project completion. As in July 2015, we anticipate requesting a contract amendment to extend the end date to 30 September 2016 to ensure that costs associated with the preparation of the project reports, preparations for the 2016 field program and complete project accounting and deliverables are covered by the 2015-16 project. A detailed accounting of all

personnel time, expenses, and in-kind contributions related to this project will be submitted to the PSC with an updated version of this report on or before 30 September 2016.

ACKNOWLEDGEMENTS

This project would not have been possible without the support from the Pacific Salmon Commission, Wuikinuxv First Nation, and DFO. We are grateful for their leadership and guidance throughout this project. This project is a key component of a 5-year business plan prepared for Rivers Inlet salmon by the Rivers Inlet Salmon Steering Committee. This committee includes: Rick Hansen (Chair), Sid Keay, Ted Walkus, George Cuthbert, John McCulloch, Fred Helmer, Dave Rolston, and Sandie MacLaurin. We thank each of these individuals for the continued support, guidance, and fundraising efforts for the Rivers Inlet Salmon Initiative. We thank Dave Rolston, Wuikinuxv Fisheries Manager, and Billie Johnson, Wuikinuxv Fisheries Technician, for their assistance with field logistics, broodstock collection and fry rearing in the net pens. We thank Marshall Hans (DFO), Dwayne Walkus, Tyler Mills, and Jessica Lyford (Good Hope Cannery) for their assistance with broodstock capture. Logistical support in Wuikinuxv Village was provided by Frank Johnson and Fred Smith. Air support was provided by West Coast Helicopters (Port McNeill) and Bella Coola Air (Hagensborg).

Funding for this project was provided by the Pacific Salmon Commission, with in-kind contributions from DFO, LGL Limited, the Rick Hansen Foundation, Duncanby Lodge, and Good Hope Cannery.

LITERATURE CITED

- English, K. K., and MacLaurin, S. 2015. Kilbella/Chuckwalla Chinook – Stock recovery enhancement, 2014-15. Report prepared for the Pacific Salmon Commission, Vancouver, BC. 6 p.
- Nelson, T. C., R. C. Bocking, D. E. Miller, 2000. Chinook escapement to the Chuckwalla River 1999. A comparison of mark-recapture and area-under-the-curve estimates based on biotelemetry, aerial surveys, swim counts, and carcass recoveries. Report prepared for Fisheries and Oceans Canada and the Rivers Inlet Restoration Society. 116 p.

Table 1. Aerial and swim counts of Chinook salmon in the Kilbella and Chuckwalla rivers, 2015.

River/Date/Section	River km		Chinook Salmon		
	From:	To:	Live	Dead	Total
KILBELLA RIVER					
<u>25 August (Helicopter)</u>					
Mallon Cr to Cascades	40.0	29.5	3	0	3
Cascades to 9-Mile Br	29.5	17.3	1	0	1
9-Mile Br to Mouth	17.3	0.0	0	0	0
Total			4	0	4
<u>4 September (Helicopter)</u>					
Mallon Cr to Cascades	40.0	29.5	1	0	1
Cascades to 9-Mile Br	29.5	17.3	0	0	0
9-Mile Br to Mouth	17.3	0.0	0	0	0
Total			1	0	1
CHUCKWALLA RIVER					
<u>25 August (Helicopter)</u>					
Johns Cr to Cascades	23.0	20.0	16	0	16
Cascades to Canyon	20.0	14.0	5	0	5
Canyon to Bridge Pool	14.0	6.0	0	0	0
Bridge Pool to Mouth	6.0	0.0	0	1	1
Total			21	1	22
<u>3 September (Helicopter)</u>					
Johns Cr to Cascades	23.0	20.0	3	0	3
Cascades to Canyon	20.0	14.0	1	0	1
Canyon to Bridge Pool	14.0	6.0	2	0	2
Bridge Pool to Mouth	6.0	0.0	0	0	0
Total			6	0	6
<u>6 September (Swim Survey)</u>					
Bridge Pool to Mouth	6.0	0.0	0	1	1

Table 2. Number of Chinook salmon encountered during broodstock collection and swim counts on the Kilbella and Chuckwalla rivers, 2015.

River/Date	Number of Sets	Live - Unmarked		Live - Recaps		Total Live Catch				Dead - Unmarked		Total (Live + Dead)		
		Female	Male	Female	Male	Female	Male	Total	CPUE	Female	Male	Female	Male	Total
Kilbella River^a														
27-Aug	11	4	7	0	0	4	7	11	1.0	0	0	4	7	11
4-Sep	15	4	7	0	1	4	8	12	0.8	0	0	4	8	12
5-Sep	12	2	8	0	0	2	8	10	0.8	3	0	5	8	13
Total	38	10	22	0	1	10	23	33	0.9	3	0	13	23	36
Chuckwalla River^b														
26-Aug	18	9	6	1	1	10	7	17	0.9	0	0	10	7	17
3-Sep	10	3	2	0	1	3	3	6	0.6	0	0	3	3	6
6-Sep ^c	-	0	0	0	0	0	0	0	n/a	1	0	1	0	1
11-Sep	15	2	3	0	0	2	3	5	0	1	0	3	3	6
Total	43	14	11	1	2	15	13	28	0.7	2	0	17	13	30
Grand Total	81	24	33	1	3	25	36	61	0.8	5	0	30	36	66

^a All fish were encountered between Mallon Cr and the Cascades.

^b 2 live fish (1F, 1M) and 1 carcass (F) were encountered between Cascades and Canyon, 1 carcass (F) was encountered below the Bridge Pool, and all other fish were encountered between Johns Cr and Cascades.

^c No broodstock collection occurred on 6 September, the carcass was found during a swim count.

Table 3. Number of live Chinook salmon encountered during broodstock collection in 2015 that were spawned, green, or spent.

River/Date	Female				Male - All released				Total
	Spawned - Ripe		Released		Spawned		Unspawned		
	Full	Partial	Green	Spent	Ripe	Ripe	Green	Spent	
Kilbella River									
27-Aug	2	1	1	0	3	3	0	1	11
4-Sep	0	2	1	1	3	3	0	2	12
5-Sep	1	0	1	0	3	4	0	1	10
Total	3	3	3	1	9	10	0	4	33
Chuckwalla River									
26-Aug	1	0	7	2	4	1	2	0	17
3-Sep	0	1	1	1	3	0	0	0	6
11-Sep	0	0	0	2	1	0	0	2	5
Total	1	1	8	5	8	1	2	2	28
Grand Total	4	4	11	6	17	11	2	6	61

Table 4. Number of female Chinook salmon spawned, eggs planted at the Snootli Hatchery, and fry reared in net pens and released in Rivers Inlet for broods 2014 and 2015.

River/Year	Females Spawned	Eggs Planted at Snootli (Est.)	Number of Fry		Egg-to-Release Survival
			Transferred to Sea Pens	Released in Rivers Inlet	
Kilbella River					
2014-15	3	12,859	12,224	12,122	94.3%
2015-16	6	21,252	19,523	19,518	91.8%
Total	9	34,111	31,747	31,640	92.8%
Chuckwalla River					
2014-15	1	5,998	4,297	4,258	71.0%
2015-16	2	11,601	10,810	10,793	93.0%
Total	3	17,599	15,107	15,051	85.5%
Both Rivers					
2014-15	4	18,857	16,521	16,380	86.9%
2015-16	8	32,853	30,333	30,311	92.3%
Total	12	51,710	46,854	46,691	90.3%

Notes:

2014-15: 1 female from the Kilbella was partially spent.

2015-16: 3 females from the Kilbella and 1 from the Chuckwalla were partially spent.

Table 5. Age composition of Chinook salmon sampled in the Kilbella/Chuckwalla rivers in 2014.

River	European	1.2	1.3	1.4	M2	M3	Total
	Gilbert-Rich	4 ₂	5 ₂	6 ₂	2M	3M	
Kilbella	Male	0	0	1	2	0	3
	Female	0	3	3	0	2	8
	Total	0	3	4	2	2	11
Chuckwalla	Male	1	0	0	0	1	2
	Female	0	1	0	0	1	2
	Total	1	1	0	0	2	4

Notes:

M2 (2M) = 2 marine annuli; M3 (3M) = 3 marine annuli

Table 6. Age composition of Chinook salmon sampled in the Kilbella/Chuckwalla rivers in 2015.

River	European	0.2	0.3	0.4	1.1	1.2	1.3	1.4	M2	M3	M4	Total
	Gilbert-Rich	3 ₁	4 ₁	5 ₁	3 ₂	4 ₂	5 ₂	6 ₂	2M	3M	4M	
Kilbella	Male	2	1	0	1	2	11	0	1	0	1	19
	Female	0	0	0	0	0	7	0	0	4	1	12
	Total	2	1	0	1	2	18	0	1	4	2	31
Chuckwalla	Male	0	0	0	0	5	5	1	0	0	0	11
	Female	0	0	1	0	0	10	1	0	3	0	15
	Total	0	0	1	0	5	15	2	0	3	0	26

Notes:

M2 (2M) = 2 marine annuli; M3 (3M) = 3 marine annuli; M4 (4M) = 4 marine annuli



Figure 1. Map of the study area showing the Kilbella and Chuckwalla rivers and location of the net pens at the head of Rivers Inlet.



Figure 2. Location of gillnet sets made in the upper Kilbella River, 2015. Labels indicate the month/day and set number (e.g., “9/5-1” is the first set made on 5 September). Note that some set sites may not be shown if their location overlaps with other sets. The blue line indicates the GPS tracks made during broodstock collection on each day.



Figure 3. Location of Chinook salmon captured in the upper Kilbella River, 2015. Data labels indicate the month/day, fish ID, sex, and fish condition (e.g., “9/5-47-F-R” is fish #47, female, ripe, caught on 5 September). Note that some fish captured may not be shown if their capture location overlaps with other fish. The blue line indicates the GPS tracks made during broodstock collection on each day.




Figure 4. Length-frequency distribution of Chinook salmon sampled in 2015, by river and sex. Post-orbital hypural (POH) length was used, and both live and dead fish were included.



Figure 5. Location of gillnet sets made in the upper Chuckwalla River, 2015. Labels indicate the month/day and set number (e.g., “9/5-1” is the first set made on 5 September). Note that some set sites may not be shown if their location overlaps with other sets. The blue line indicates the GPS tracks made during broodstock collection on each day.



Figure 6. Location of Chinook salmon captured in the upper Chuckwalla River, 2015. Data labels indicate the month/day, fish ID, sex, and condition (e.g., “9/5-47-F-R” is fish #47, female, ripe, caught on 5 September). Note that some fish captured may not be shown if their capture location overlaps with other fish. The blue line indicates the GPS tracks made during broodstock collection on each day.



Photo 1. Gillnet deployed in the upper Chuckwalla River on 26 August, 2015.



Photo 2. Eggs being collected from a ripe female Chinook salmon on the upper Kilbella River, 27 August 2015.



Photo 3. Milt being collected from a ripe male Chinook salmon on the upper Kilbella River, 27 August 2015.



Photo 4. Aerial photograph showing the difference in water clarity between the Kilbella (top) and Chuckwalla (bottom) rivers on 25 August 2015. River flow is from right to left in the photograph.



Photo 5. Sockeye salmon captured in the upper Kilbella River, 5 September 2015. This was the only sockeye salmon observed in either river in 2015.