

Increased north-migrating Chinook salmon indicator stock coded-wire tagging to improve the quality of Chinook indicator stock analyses

Final Report to the Northern Endowment Fund Committee

2017

**Jason Mahoney
Carolyn Churchland
Jennifer Sandher**

Fisheries and Oceans Canada
Salmonid Enhancement Program
200-401 Burrard St.
Vancouver, B.C.
V6C 3S4

Report Prepared For:

Pacific Salmon Commission
Northern Boundary Restoration and Enhancement Fund
600-1155 Robson Street
Vancouver, BC.
Canada V6E 1B5

INTRODUCTION

The United States and Canada have recognized the importance of developing and maintaining a coded-wire tag (CWT) program to estimate exploitation rates of Chinook salmon stocks and better define their time-area distributions to develop management options at least since the August 13, 1985 Memorandum of Understanding (PSC 2004: March 2004 Annexes, P. 96). In 1999, government-to-government negotiations culminated in the successful renewal of a long-term fishing agreement under the Pacific Salmon Treaty (PST). With this agreement CWTs became one of the key methods to assess harvest rate reduction compliance. Furthermore, in the 2009 agreement, CWT-based individual stock-based management (ISBM) indices are used to monitor relative exploitation rate reductions from the base period (para. 8(b&c), 9(b&c)). CWT data and analyses are also important for developing stock abundance forecasts used in the Pacific Salmon Commission (PSC) Chinook Technical Committee (CTC) Coast wide model calibration. In 2005, the PSC convened an Expert Panel to review the utility of the CWT system for future PST implementation (PSC Tech. Report No. 18). The Panel reported that the CWT program must be relied upon as the primary fishery and stock assessment tool for at least the next 5-10 years (Hankin et al. 2005). No accepted alternative technology currently exists that is capable of providing the data necessary for the implementation of the PST. In 2006, the PSC convened a CWT Work Group to review and recommend a plan to implement the recommendations of the PSC Expert Panel (PSC Tech. Report No. 25). This report states that the principal factors influencing the uncertainty surrounding CWT-based estimates of exploitation rates are those affecting precision and those causing bias. The major factors affecting precision are the number of CWTs released and sample rates for fisheries and escapements. As increased tagging is the most cost effective way to increase precision of CWT-based statistics for these indicator stocks, this project would maintain increased tagging beyond base tagging levels funded by Canadian Department of Fisheries & Oceans (CDFO) to the release group size standards based on expected marine survivals for the 2015 brood.

PROJECT OBJECTIVES

The objective of this project is the application of CWTs on Adipose Fin-Clipped (AFC) juvenile Chinook salmon incremental to the current tagging levels already funded by CDFO for nine British Columbia (BC) Chinook indicator stocks to meet the CWT release group size standards as outlined in PSC Tech. Report No. 25.

METHODS

Adult Chinook salmon are captured for brood by CDFO staff upon return to their spawning rivers in the summer or fall. Exact capture methods differ by location, but they include a variety of strategies such as: weir, fish ladder, beach seine, angling and tangle net. Adult Chinook are held at a hatchery, either in concrete ponds or in circular fiberglass tubs until they are ready to be spawned. This determination is

made by the fish culturists, who check the females to ensure that the eggs are loose, the belly is soft, and the ovipositor is distended. Eggs are gathered by incising the belly of the female and collecting them in a disinfected container. Milt is then added from one or two males to fertilize the eggs. Water is then added to the fertilized eggs, after which they are disinfected in a solution of Ovadine and water for 10 minutes. It is at this stage that fish culturists must conduct bulk fecundity sampling to try to ensure that egg targets are met.

Fertilized eggs are placed into the incubation container, which may be a Heath Tray, Atkins cell, or bulk box. Fungal treatments are conducted on eggs, typically using Parasite-S. Chinook eggs typically require approximately 500-525 accumulated thermal units prior to hatching (Billard & Jensen, 1996). Swim up fry are ponded into rearing containers where they are reared until they are of suitable size for coded wire tag application and adipose fin-clipping. Fish health monitoring occurs continuously throughout the early rearing period, with prophylactic and antibiotic treatments used as required. The Salmonid Enhancement Program (SEP) veterinarian is available to diagnose any fish health or well-being issues that may arise and works closely with all hatcheries to ensure that fish are healthy prior to marking and release.

The procedures used to implant the CWTs into juvenile Chinook are documented in detail by Nichols & Hillaby (1990). Marking and tagging of sub-yearling Chinook (fish that have hatched in the spring or winter preceding marking, and that will be released shortly after) occurs when fish begin to reach 3-4 grams (g), with a typical release size of 6 g. The juveniles must not be fed for 48 hours prior to marking and tagging, as this reduces the output of ammonia and excretory by-products associated with stressful fish handling. Juvenile Chinook are transported to the tagging area in small batches into a holding tank prior to being anaesthetized using Tricaine methanesulfonate (TMS). Following anaesthetization, the adipose fin of each juvenile salmon is excised using a set of surgical scissors, after which it is placed nose-first into a Mark IV CWT machine for tag insertion in the nasal tissue. Fish size-grading will occur at fin-clipping to ensure that the appropriate sized head mold is used for fish size. Typically, there are 2 or 3 Mark IVs operating simultaneously, often with different sized head molds. Tagged fish are passed through a quality control device to ensure successful tag implantation.


Tag placement and retention is monitored in 3 ways. A small group of tagged fish will be retained at the end of each tagging day for a 24 hour retention check the following day. In many instances, small checks will be conducted on a more immediate basis to ensure quality control. In addition to the 24 hour retention check, a larger group of at least 500 fish is kept for up to 30 days to conduct a longer term retention check (Table 2). Finally, to ensure proper tag placement, one tagged smolt is euthanized and dissected every hour, with the tag placement observed (Figure 1).

Detailed operational procedures may vary slightly by facility, but generally follow the practices as described by Nichols & Hillaby (1990).

Following a holding period of about two weeks after tag application, juvenile Chinook are released from the hatchery back into their river of origin. Yearling juvenile Chinook (fish to be released the following year) are retained at the hatchery for further rearing. Hatcheries that have swim-in infrastructure will

release directly from the hatchery to the river, while stocks from hatcheries without this mechanism will be transported to the river and force released. Juvenile releases typically occur when Chinook are smolting. Some juveniles may stay in the river for a short period of time prior to migrating to saltwater.

Figure 1 - Proper coded wire tag placement (Nichols & Hillaby, 1990)



RESULTS

Coded wire tagging began on schedule at all sites in 2016, as water temperatures during the incubation and rearing period were relatively normal. All tagging project operations were completed at or before the expected date, and there were no significant fish health issues during the tagging process.

Table 1 - Tag application schedule by hatchery.

Hatchery	Stock	Tagging Period
Chilliwack	Chilliwack River	April - May
Cowichan	Cowichan River	March - April
Deep Creek	Kitsumkalum River	May - June
Harrison	Harrison River	April - May
Quinsam	Quinsam River	March – April
Robertson	Robertson River	April - May
Shuswap	Lower Shuswap River	April – May
Snootli	Atnarko River	May – June
Spius	Nicola River	Sept - Oct

Table 2 - Estimated tag loss rate by hatchery stock during 2016 tag application.¹

Hatchery	Stock	Tag Loss
Chilliwack	Chilliwack River	2.65%
Cowichan	Cowichan River	0.48%
Deep Creek	Kitsumkalum River	4.54%
Harrison	Harrison River	4.00 %
Quinsam	Quinsam River	1.24%
Robertson	Robertson River	0.22%
Shuswap	Lower Shuswap River	0.44%
Snootli	Atnarko River	0.37%
Spius	Nicola River	2.02%

¹Updated from MRP database - March 2017.

Table 3 – Tag targets and actuals (base level and incremental) for the 2016 tag application (2015 Brood).¹

Stock	Base Level CWT Target	Additional CWT Target	Total CWT Target	Total CWT Applied in 2016
Atnarko R	150,000	250,000	400,000	409,088
Chilliwack R	100,000	100,000	200,000	199,355
Cowichan R	200,000	400,000	600,000	708,370
Harrison R	100,000	200,000	300,000	277,330
Kitsumkalum	60,000	200,000	260,000	266,732
Nicola R	140,000	60,000	200,000	172,698
Quinsam R	200,000	300,000	500,000	727,624
Robertson Cr	200,000	250,000	450,000	692,685
Shuswap R Low	250,000	300,000	550,000	496,044
Total	1,400,000	2,060,000	3,460,000	3,949,926

¹Updated from MRP database - March 2017.

Table 4 - Base tagging level and percentage increase due to expanded tagging, by brood year and stock.

Stock	Base Tag Level	2009	2010	2011	2012	2013	2014	2015
Atnarko R	150,000	246%	239%	194%	166%	163%	170%	173%
Chilliwack R	100,000	90%	98%	96%	100%	212%	213%	99%
Cowichan R	200,000	110%	119%	175%	199%	12%	307%	254%
Harrison R	100,000	113%	95%	176%	190%	172%	192%	177%
Kitsumkalum	60,000	344%	354%	249%	113%	175%	186%	345%
Nicola R	140,000	38%	34%	52%	36%	24%	23%	23%
Quinsam R	200,000	227%	217%	185%	224%	200%	229%	264%
Robertson Cr	200,000	126%	124%	122%	126%	139%	161%	246%
Shuswap R Low	250,000	93%	96%	101%	104%	85%	76%	98%

Table 5 - Total observed fishery CWTs, by brood year and stock.¹

Stock	2009	2010	2011	2012	2013	2014
Atnarko R	690	720	857	340	22	6
Chilliwack R	393	1289	826	280	171	21
Cowichan R	291	418	395	160	44	10
Harrison R	74	256	203	27	57	3
Kitsumkalum	53	224	167	17	9	0
Nicola r	48	9	41	25	0	0
Quinsam R	88	60	100	359	100	8
Robertson Cr	56	325	64	992	155	21
Shuswap R Low	325	1170	458	308	290	8

¹Updated from MRP database - March 2017.

Table 6 - Total observed fishery CWTs directly attributable to PSC funded expanded tagging, by brood year and stock.

Stock	2009	2010	2011	2012	2013	2014
Atnarko R	491	508	566	212	14	4
Chilliwack R	186	638	405	140	116	14
Cowichan R	152	227	251	106	5	8
Harrison R	39	125	129	18	36	2
Kitsumkalum	41	175	119	9	6	0
Nicola R	13	2	14	7	0	0
Quinsam R	61	41	65	248	67	6
Robertson Cr	31	180	35	553	90	13
Shuswap R Low	157	573	230	157	133	3

DISCUSSION

Tag application numbers exceeded targets on 5 of 9 stocks, with 4 of the 9 stocks coming in a little under target. The total project overall goal of 3.46 million CWTs applied was exceeded by 489,926. It is common to exceed CWT targets as the spools of wire that the tags are printed on often have 5-10% more tags than is stated. Increases in tagging numbers help to increase the number of observed and estimated CWTs, which will result in increased precision in the estimated survival and exploitation rate.

Actual tag application numbers are subject to variability for several reasons, including, but not limited to: insufficient broodstock available for egg target, lower than expected in-hatchery survival, or unresolvable tagging equipment malfunctions. Hatcheries that have large production targets to support fisheries will rarely fail to reach their tag target. For example, Robertson Creek hatchery has a production target of 6M smolts, of which only 450K are required for tagging (base level + incremental). Thus, even with a very weak adult return and a fraction of their egg target, the tagging target can still be met. Conversely, stocks that are enhanced purely for stock assessment purposes (Lower Shuswap and Kitsumkalum) have less flexibility in their targets. If there are surplus juveniles available they will typically all be tagged; however, if there is any issue obtaining the release target (such as at Nicola in 2016) the tag target will be compromised.

It must be noted that Kitsumkalum tag target is unique compared to all other stocks included in this project in that it is comprised of two different year classes; fed fry that are tagged and released the spring following emergence, and yearling smolts that are tagged at the same time as fed fry but that are held for an additional year prior to release.

Although the direct results of the tagging completed in 2016 (2015 brood) will not be apparent until those fish begin to recruit to the fishery and escapement as jacks in 2017, it can be assumed with certainty that the number of observed tags in catch and escapement will have increased as a function of the increase in tagging over the base level. Incremental tagging has been occurring at sites since 2009,

with some sites starting earlier. This work has been previously funded through the Coded Wire Tag Improvement Fund of the PSC, and the results of this earlier work can be used to illustrate the future benefits of this 2016 NEF funded project. Note that at the time of this report, brood years 2010-2014 are considered preliminary as there are still cohorts that will likely recruit to catch and escapement in upcoming years. Table 5 shows the total observed fishery tags, by brood year and stock (data current as of March 2017), while Table 6 shows the observed tags that are directly attributable to the expanded tagging projects. It is important to note that only the 2009 brood can be considered a complete brood year.

It is too early to be able to assess the ultimate success of this project, as well as those that preceded it. This project represents the first step in a complex process that requires fishery and escapement sampling to recover CWTs. Even upon completion of the 2016 spawning and catch year, there are still cohorts that have yet to return from the majority of the years of expanded tagging.

APPENDIX 1

Financial Expenditure Summary

Details of expenditures registered in the DFO financial system at fiscal year-end.

Funding Total	\$ 239,061.87
----------------------	----------------------

DFO Casual Hire Salary	\$ 41,601.17
Snootli Hatchery Contracting Costs	\$ 25,000.00
Spius Hatchery Contracting Costs	\$ 36,945.11
Kitsumkalum Hatchery (Deep Creek) Contracting Costs	\$ 26,250.00
Cowichan Hatchery Contracting Costs	\$ 49,080.00
Robertson Hatchery Contracting Costs	\$ 29,475.00
Chehalis hatchery Contracting Costs	\$ 22,000.00
Equipment & Supplies	\$ 8,315.11
Broker fees	\$ 395.48
Total Costs	\$ 239,061.87

Balance	\$ -
----------------	-------------

REFERENCES

Hankin, D.G. (Chair), J.H. Clark, R.B. Deriso, J.C. Garza, G.S. Morishima, B.F. Riddell, and C. Schwarz. 2005. Report of the expert panel on the future of the coded wire tag recovery program for Pacific salmon. Pacific Salmon Commission Technical Report No. 18. 230 pp

Pacific Salmon Commission Coded Wire Tag Workgroup. 2008. An action plan in response to Coded Wire Tag (CWT) Expert Panel Recommendations. Pacific Salmon Comm. Tech. Rep. No. 25: 170 p.

Billard, R., and J.O.T. Jensen. 1996. Gamete removal, fertilization and incubation. Pages 291- 363 In: W. Pennell and B.A. Barton, Editors. Developments in Aquaculture and Fisheries Science V. 29: Principles of Salmonid Culture. Elsevier, Amsterdam.

Nichols, T.L., and J.E. Hillaby. 1990. Manual for Coded-Wire Tagging and Fin-Clipping of Juvenile Salmon at Enhancement Operations Facilities. Prepared under contract #90SB.FP501-7-0060/A to Supply and Services Canada by Streamline Consulting Services Limited