

**2013 Alsek Sockeye Salmon Run
Reconstruction Using Genetic Stock
Identification**
(A project funded through the Northern Fund of the
Pacific Salmon Commission)

Prepared by:

Bonnie Huebschwerlen
Treaties and Fisheries
Fisheries and Oceans Canada
Whitehorse, Yukon
Y1A 3V1

July 2014

TABLE OF CONTENTS

List of Figures	ii
List of Appendices	ii
Introduction.....	3
Objectives	6
Methodology.....	6
DNA Collection and Analysis	6
Run Reconstruction.....	6
Results.....	7
DNA Collection and Analysis	7
Run Reconstruction.....	7
Discussion and Recommendations	7
Literature Cited	7
Appendices.....	8
Financial Summary	10

List of Figures

Figure 1. Alsek River drainage map.	5
--	---

List of Appendices

Appendix 1. Population estimate for the 2013 Alsek sockeye salmon using GSI analysis and expansion of the Klukshu weir count. Model developed by W.J. Gazey Research, 2010.....	9
---	---

Introduction

The Alsek River originates in the Yukon Territory, Canada, and flows in a southerly direction into the Gulf of Alaska, southeast of Yakutat, Alaska (Figure 1). Alsek River sockeye salmon (*Oncorhynchus nerka*) are caught primarily in U.S. commercial and subsistence set gillnet fisheries in the lower Alsek River and in aboriginal and recreational fisheries in Canada. Small harvests of Alsek sockeye are probably taken in marine commercial gillnet fisheries near Yakutat. These populations are managed jointly by Canada and the U.S. through a sub-committee of the Pacific Salmon Commission (PSC) as part of the Canada/U.S. Pacific Salmon Treaty (PST) adopted in 1985 (TTC 1999). Historically, the status of sockeye salmon has been evaluated by monitoring escapement trends of what were assumed to be two principal sockeye stocks within the drainage: Klukshu River and Nesketahen Lake sockeye salmon.

Prior to 2000, the Alsek River sockeye salmon return was largely unknown (a mark-recapture study was conducted in 1983 by ADF&G) because stock assessment projects to determine system-wide escapements had not yet been fully developed. Escapements were known only for the Klukshu River. An escapement goal was originally developed for the Klukshu River in 2000 but very little else was known about the magnitude of run sizes and system wide production capacity. In 2000, a pilot project was initiated to determine the feasibility of assessing the drainage wide escapement for sockeye salmon using mark-recapture. After achieving the objectives of the 2000 study, the program was continued from 2001 through to 2004. Since then, the only other assessments of the total sockeye return to the Alsek were made in 2005 - 2006 and 2011 and 2012 using genetic stock identification (GSI). In 2013, a revised escapement goal for Klukshu and a system wide goal for the Alsek was adopted by the Transboundary Technical Committee.

Since 1976, weir counts by the Fisheries and Oceans Canada (DFO) in co-operation with the Champagne-Aishihik First Nation (CAFN) have been made on the Klukshu River. Recently, sockeye salmon weir counts at Klukshu, in conjunction with sockeye catch rate information from the commercial fishery at Dry Bay plus GSI of tissue samples taken in the Dry Bay commercial fishery, were used to estimate the Alsek sockeye population in 2005 and 2006 (Waugh and Stark 2008a&b). The results were encouraging; a project recommendation was made by the Northern Fund Committee (PSC) in 2008 to develop a sampling strategy (statistically valid) which would include methods of calculations and the precision expected based on various sample sizes and stock contribution levels for the apportionment of Alsek sockeye abundance into the requisite stocks (i.e. Klukshu and others) which, in conjunction with the Klukshu weir counts, would provide the foundation for reconstructing the sockeye return to the Alsek River. The project was completed by W. J. Gazey Research with funding from the Northern Fund ("*GSI Sample Size Requirements for In-river Run Reconstruction of Alsek Chinook and Sockeye Stocks*, W. J. Gazey, April 2010"). Gazey's analysis provided a model with which to determine the required sample sizes needed to reconstruct the Alsek sockeye returns to achieve a desired precision at a prescribed confidence level.

The 2009-2018 Transboundary chapter of the Pacific Salmon Treaty (PST) tasked the Parties and the Transboundary Technical Committee (TTC) to explore methods to determine in-river abundance for Alsek sockeye salmon. Further, the Transboundary Panel, as noted in the "*Pacific Salmon Commission Transboundary Panel Strategic Salmon Plan, March 2009*", has committed to develop and implement abundance-based management regimes for Alsek River sockeye which includes estimates of total abundance. The project proposed for 2013 was designed to meet the

objectives set out in the PST and the Panel's strategic salmon plan. Subsequently, funding was secured through the Northern Fund of the Pacific Salmon Commission to analyze tissue samples collected from the U.S. commercial fishery in Dry Bay for the purpose of reconstructing the 2013 Alek sockeye salmon return.




Figure 1. Alsek River drainage map.

Objectives

The following objectives are expected to be achieved from the collection and analysis of sockeye salmon tissue samples taken in the U.S. Dry Bay commercial fishery for 2013:

- Collection of fishery performance data and tissue samples (up to 750) from the Alsek sockeye salmon commercial fishery (Dry Bay, Alaska);
- GSI analysis of tissue samples collected in the Dry Bay commercial fishery;
- Run reconstruction of the 2013 Alsek sockeye return at the desired precision (25%) within a 95% confidence level;
- Assessment of the Klukshu River sockeye salmon contribution to the drainage wide escapement within the Alsek River.

Methodology

DNA Collection and Analysis

Approximately 750 Alsek River sockeye salmon tissue samples (severed axillary appendages) were to be collected from the 2013 U.S. commercial fishery catch located in Dry Bay, Alaska from the beginning of June to the middle of August. The samples were to be sent to Terry Beacham (DFO Salmon Genetics Laboratory) for GSI analysis to develop weekly contributions of Klukshu River and other sockeye stocks to the commercial sockeye catch. The DNA analysis was to be conducted as per Withler et al (2000).

Run Reconstruction

Run reconstruction methodology was to be followed using the model detailed in “*GSI Sample Size Requirements for In-river Run Reconstruction of Alsek Chinook and Sockeye Stocks*, W.J. Gazey, April 2010”. Sockeye salmon weir counts at Klukshu, in conjunction with sockeye catch rate information from the commercial fishery at Dry Bay and GSI of tissue samples taken in the Dry Bay commercial fishery, were to be used to estimate the 2013 Alsek River sockeye salmon population.

Results

DNA Collection and Analysis

A total of 734 sockeye tissue samples (axillary appendages) from the Dry Bay commercial fishery were received from the Alaska Department of Fish and Game and shipped to Terry Beacham (DFO – Genetics Lab) for analysis in the fall of 2013. A total of 733 samples were analyzed and results were received from the DFO genetics lab in January of 2014. Sample analysis was roughly apportioned based on the run timing observed during the 2013 U.S. commercial sockeye fishery. Samples were collected beginning in statistical week 23 (starting June 2nd) and completed in week 33 (ending August 17th).

Run Reconstruction

The total Alesek sockeye run for 2013 was estimated to be 91,876 (above Dry Bay escapement plus the Dry Bay catches; Appendix 1). The above Dry Bay escapement was estimated to be 84,279 (95% CI = 16,466 – 152,091, SD = 34,599). This resulted in an estimated 8% harvest rate in the combined fisheries located in Dry Bay (catch = 7,517). The Klukshu stock contributed a total of 4,308 sockeye (approximately 5%) to the total Alesek sockeye escapement.

Discussion and Recommendations

As detailed in Gazey 2010, the major assumptions for the estimation of aggregate escapement, associated variance and subsequent sample size requirements at specified levels of precision and confidence were:

1. Equal vulnerability of capture. Migration behaviour and size differences of returning stocks may cause unequal vulnerability of stocks to the test or commercial fisheries. Estimates are likely sensitive to violation.
2. Minimum in-river mortality. Computationally, straight-forward to adjust the weir counts for any suspected mortality. Impacts the aggregate escapement estimate and fraction to be sampled.
3. Functional relationship (linear assumed here) between sample size adjustment to obtain effective sample size and the Klukshu stock proportion. New baseline GSI data may alter or change the functional relationship (e.g., non-linear or non-stationary).

Literature Cited

Gazey, W. J. 2010. GSI Sample Size Requirements for In-river Run Reconstruction of Alesek Chinook and Sockeye Stocks. Pacific Salmon Commission, Vancouver, British Columbia.

Withler, R.E, Le, K.D., Nelson, R.J., Miller, K.M., and Beacham, T.D. 2000. Intact genetic structure and high levels of genetic diversity in bottlenecked sockeye salmon, *Oncorhynchus nerka*, populations of the Fraser River, British Columbia, Canada. *Can.J. Fish. Aquatic Sci.* 57: 1985–1998.

Appendices

Appendix 1. Population estimate for the 2013 Alesek sockeye salmon using GSI analysis and expansion of the Klukshu weir count. Model developed by W.J. Gazey Research, 2010.

GSI Run Reconstruction for the Alesek River - Sockeye 2013									
Required input values are in red font									
Output of required GSI sample using the "EstSample" macro in blue font									
Week	Run Weight	Klukshu Prop.	Run Standard	Escapement Above Dry Bay	"Actual" Sample	Effective Sample	SD Prop	SD wp	SD Escape
1 (23-24)	0.059	0.001	0.059	4,956	180	45.8	0.004	0.000	2,035
2 (25)	0.068	0.006	0.068	5,736	80	21.1	0.017	0.001	2,355
3 (26)	0.074	0.016	0.074	6,261	84	23.5	0.026	0.002	2,570
4 (27)	0.098	0.009	0.098	8,234	96	25.7	0.018	0.002	3,380
5 (28)	0.097	0.022	0.097	8,156	80	23.2	0.031	0.003	3,348
6 (29)	0.166	0.055	0.166	13,994	80	27.6	0.043	0.007	5,745
7 (30)	0.227	0.127	0.227	19,117	39	18.2	0.078	0.018	7,848
8 (31-33)	0.211	0.040	0.211	17,824	94	30.1	0.036	0.008	7,317
Total	1.000		1.000	84,279	733			0.021	34,599
Klukshu Above Dry Bay		4,308						Confidence Interval	
Desired Confidence		95%			% Sample =	0.9%		Low	High
Desired Precision		25%			Precision =	80.5%			
Effect. Sample Coeff.		0.253	1.677					16,466	152,091
Effective sample size coefficients based on 2005-2007 studies:									
Sockeye		0.253	1.677						
Chinook		0.342	-0.005						
							5.1% Klukshu comp.		
							Total Run: 91,876		

Financial Summary

Project Budget Form

Name of Project: 2013 Alek Sockeye Run Reconstruction Using GSI

ELIGIBLE COSTS

**TOTAL
PROJECT
BUDGET**

**OTHER
FUNDING**

**PSC
N. FUND
GRANT
AMOUNT**

**Labour
Wages & Salaries**

Position	# of crew	# of work days	hrs per day	rate per hour	Total (In-kind & cash + PSC Amount)	In-Kind & Cash	PSC Amount	Actual Expenditures	Variance
Project Authority (DFO Bi-\$)	1	1	7.5	42	316	316			
Biologist (DFO Bi-3)	1	1	7.5	39	293	293			
Senior Technician (DFO EG-5)	1	10	7.5	35	2,618	2,618			
Financial Officer (DFO As-2)	1	1	7.5	30	225	225			
Fisheries Technician	1	10	7.5	32	2,400	2,400			
Person Days (# of crew x work days)		13		sub total	5,852	5,852			

Labour - Employer Costs (percent of wages subtotal amount)

rate	20%	sub total	1,170	1,170		
------	-----	------------------	-------	-------	--	--

Subcontractors & Consultants

# of crew	# of work days	hrs per day	rate per hour			
Insurance if applicable	rate	0%		sub total		

Volunteer Labour

# of crew	# of work days	hrs per day				
Skilled						
Un-skilled						
Insurance if applicable	rate	0%		sub total		

Total Labour Costs

7,022

7,022

Site / Project Costs

Provide details in the space below (use an additional page if needed)

Travel (do not include to & from work)						
Small Tools & Equipment						
Site Supplies & Materials	lab suples (to cover 500 samples)			12,750	12,750	12,750
Equipment Rental						
Work & Safety Gear						
Repairs & Maintenance						
Permits						
Technical Monitoring						
Other site costs						
Total Site / Project Costs				12,750	12,750	12,750

Project Budget Form (continued)

ELIGIBLE COSTS				BUDGET	OTHER FUNDING	CONTRIBUTION FUNDING		
				Total (PSC + In-kind + cash)	In-Kind & Cash	PSC Amount	PSC Amount	PSC Amount
Training (e.g Swiftwater, bear aware, electrofishing, etc).								
Name of course	# of crew	# of days						
Total Training Costs								
Overhead / Indirect Costs								
Office space; including utilities, etc.				800	800			
Insurance								
Office supplies				100	100			
Telephone & long Distance				100	100			
Photocopies & printing								
Indirect/overhead costs								
(If the PSC contribution to Indirect costs exceeds 20% of the total PSC grant you will be required to submit back-up documentation justifying the expense).								
Other overhead costs								
Total Overhead Costs				1,000	1,000			
Provide details in the space below (use an additional page if needed)								
Capital Costs / Assets								
Assets are things of value that have an initial cost of \$250 or more and which can be readily misappropriated for personal use or gain or which are not, or will not be, fully consumed during the term of the project.								
Total Capital Costs								
Project Total Costs				20,772	8,022	12,750	12,750	-
						Project Balance	12,750	
						Hold Back (not deposited)	1,275	
						Owed to DFO as of Aug 31/	1,275	
Budget Summary								
(PSC + in-kind + cash)								
Total Labour Costs			7,022					
Total Site / Project Costs			12,750					
Total Training Costs			-					
Total Overhead Costs			1,000					
Total Capital Costs			-					
Project Total			20,772					